



Full wwPDB X-ray Structure Validation Report ⓘ

May 27, 2020 – 06:10 pm BST

PDB ID : 1QOQ
Title : CRYSTAL STRUCTURE OF WILD-TYPE TRYPTOPHAN SYNTHASE
COMPLEXED WITH INDOLE GLYCEROL PHOSPHATE
Authors : Weyand, M.; Schlichting, I.
Deposited on : 1999-11-15
Resolution : 1.80 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.11
buster-report : 1.1.7 (2018)
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.11


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.80 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	5950 (1.80-1.80)
Clashscore	141614	6793 (1.80-1.80)
Ramachandran outliers	138981	6697 (1.80-1.80)
Sidechain outliers	138945	6696 (1.80-1.80)
RSRZ outliers	127900	5850 (1.80-1.80)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	268	<div> <div>4%</div> <div> <div></div> <div>80%</div> <div>13%</div> <div>• 5%</div> </div> </div>
2	B	396	<div> <div>4%</div> <div> <div></div> <div>85%</div> <div>13%</div> <div>••</div> </div> </div>

2 Entry composition [i](#)

There are 5 unique types of molecules in this entry. The entry contains 5459 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called TRYPTOPHAN SYNTHASE ALPHA CHAIN.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	254	Total	C	N	O	S	0	1	0
			1927	1227	331	361	8			


- Molecule 2 is a protein called TRYPTOPHAN SYNTHASE BETA CHAIN.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	394	Total	C	N	O	S	0	3	0
			2982	1873	521	567	21			

There are 2 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
B	34	SER	ARG	cloning artifact	UNP P0A2K1
B	175	ALA	ARG	conflict	UNP P0A2K1

- Molecule 3 is INDOLE-3-GLYCEROL PHOSPHATE (three-letter code: IGP) (formula: $C_{11}H_{14}NO_6P$).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
3	A	1	Total	C	N	O	P	0	0
			19	11	1	6	1		

- Molecule 4 is PYRIDOXAL-5'-PHOSPHATE (three-letter code: PLP) (formula: $C_8H_{10}NO_6P$).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
4	B	1	Total	C	N	O	P	0	0
			15	8	1	5	1		


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	192	Total	O	0	0
			192	192		
5	B	324	Total	O	0	0
			324	324		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: TRYPTOPHAN SYNTHASE ALPHA CHAIN



• Molecule 2: TRYPTOPHAN SYNTHASE BETA CHAIN



4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, α , β , γ	182.74Å 60.05Å 67.10Å 90.00° 94.48° 90.00°	Depositor
Resolution (Å)	20.00 – 1.80 31.15 – 1.80	Depositor EDS
% Data completeness (in resolution range)	94.0 (20.00-1.80) 94.1 (31.15-1.80)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	0.08	Depositor
$\langle I/\sigma(I) \rangle$ ¹	4.20 (at 1.79Å)	Xtriage
Refinement program	REFMAC	Depositor
R, R_{free}	0.171 , 0.210 0.169 , 0.206	Depositor DCC
R_{free} test set	3245 reflections (5.10%)	wwPDB-VP
Wilson B-factor (Å ²)	15.3	Xtriage
Anisotropy	0.746	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.41 , 62.6	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	5459	wwPDB-VP
Average B, all atoms (Å ²)	21.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.92% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality ⓘ

5.1 Standard geometry ⓘ

Bond lengths and bond angles in the following residue types are not validated in this section: IGP, PLP

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.73	0/1970	1.40	21/2674 (0.8%)
2	B	0.89	0/3056	1.54	38/4128 (0.9%)
All	All	0.83	0/5026	1.49	59/6802 (0.9%)

There are no bond length outliers.

All (59) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	341	ARG	NE-CZ-NH2	-18.64	110.98	120.30
2	B	394	ARG	NE-CZ-NH2	-14.86	112.87	120.30
2	B	379	ARG	NE-CZ-NH2	12.69	126.64	120.30
2	B	202	ARG	NE-CZ-NH1	12.33	126.47	120.30
2	B	394	ARG	NE-CZ-NH1	11.39	126.00	120.30
1	A	60	ASP	CB-CG-OD1	11.05	128.24	118.30
2	B	70	ARG	NE-CZ-NH1	10.04	125.32	120.30
1	A	225	ARG	NE-CZ-NH2	-9.73	115.43	120.30
1	A	15	ARG	NE-CZ-NH1	9.20	124.90	120.30
2	B	55	ARG	NE-CZ-NH1	8.89	124.75	120.30
2	B	341	ARG	NE-CZ-NH1	8.35	124.48	120.30
2	B	329	ASP	CB-CG-OD1	8.07	125.56	118.30
1	A	3	ARG	NE-CZ-NH1	8.02	124.31	120.30
2	B	381	ASP	CB-CG-OD2	-7.98	111.12	118.30
1	A	14	ARG	NE-CZ-NH1	7.73	124.17	120.30
1	A	225	ARG	CD-NE-CZ	7.71	134.40	123.60
2	B	297	SER	N-CA-CB	-7.50	99.25	110.50
2	B	168	ASP	CB-CG-OD1	-7.42	111.62	118.30
2	B	55	ARG	NE-CZ-NH2	-7.37	116.62	120.30
1	A	171	ARG	NE-CZ-NH2	-7.33	116.64	120.30
2	B	70	ARG	CD-NE-CZ	6.93	133.31	123.60
1	A	15	ARG	NE-CZ-NH2	-6.87	116.86	120.30

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	329	ASP	CB-CG-OD2	-6.86	112.13	118.30
2	B	168	ASP	CB-CG-OD2	6.47	124.12	118.30
1	A	89	ARG	NE-CZ-NH2	-6.37	117.12	120.30
2	B	368	LYS	N-CA-CB	-6.25	99.34	110.60
2	B	226	ALA	N-CA-CB	6.11	118.65	110.10
2	B	275	ARG	N-CA-CB	-6.06	99.70	110.60
2	B	141	ARG	CD-NE-CZ	6.02	132.03	123.60
2	B	331	GLU	OE1-CD-OE2	-5.89	116.24	123.30
2	B	187	MET	CA-CB-CG	5.86	123.26	113.30
1	A	16	GLU	N-CA-CB	-5.80	100.17	110.60
1	A	70	ARG	NE-CZ-NH1	5.73	123.17	120.30
1	A	54	PHE	CB-CG-CD2	-5.72	116.79	120.80
2	B	330	ASP	CB-CG-OD1	5.71	123.44	118.30
2	B	222	ARG	NE-CZ-NH2	-5.67	117.47	120.30
1	A	3	ARG	NE-CZ-NH2	-5.62	117.49	120.30
2	B	222	ARG	CD-NE-CZ	5.57	131.40	123.60
1	A	134	GLU	OE1-CD-OE2	-5.52	116.67	123.30
2	B	323	ASP	CB-CG-OD1	5.42	123.18	118.30
2	B	275	ARG	CA-CB-CG	5.28	125.02	113.40
2	B	287	MET	CA-CB-CG	-5.27	104.34	113.30
1	A	15	ARG	CD-NE-CZ	5.24	130.94	123.60
1	A	164	ARG	CD-NE-CZ	5.24	130.93	123.60
2	B	133	TYR	CB-CG-CD1	-5.18	117.89	121.00
2	B	379	ARG	NH1-CZ-NH2	-5.18	113.70	119.40
2	B	321	ARG	NE-CZ-NH1	-5.17	117.71	120.30
2	B	52	TYR	CB-CG-CD1	5.13	124.08	121.00
2	B	240[A]	MET	CG-SD-CE	5.12	108.40	100.20
2	B	240[B]	MET	CG-SD-CE	5.12	108.40	100.20
2	B	247	ASP	CB-CG-OD2	-5.12	113.70	118.30
1	A	112	ASP	CB-CG-OD2	-5.11	113.70	118.30
2	B	226	ALA	CB-CA-C	-5.07	102.50	110.10
1	A	115	TYR	CB-CG-CD1	5.05	124.03	121.00
1	A	93	PRO	O-C-N	5.03	130.75	122.70
1	A	158	ALA	CB-CA-C	-5.01	102.58	110.10
2	B	206	ARG	NE-CZ-NH1	5.01	122.81	120.30
1	A	97	ILE	C-N-CA	-5.01	111.77	122.30
2	B	275	ARG	CG-CD-NE	5.00	122.31	111.80

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1927	0	1933	15	0
2	B	2982	0	2952	25	0
3	A	19	0	12	0	0
4	B	15	0	6	0	0
5	A	192	0	0	0	0
5	B	324	0	0	4	0
All	All	5459	0	4903	40	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (40) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:209:GLY:HA2	2:B:240[A]:MET:HE3	1.48	0.93
2:B:65:ILE:HD11	2:B:73:LEU:HD23	1.57	0.84
2:B:143:SER:OG	2:B:144:PRO:HD3	1.84	0.77
2:B:209:GLY:HA2	2:B:240[A]:MET:CE	2.18	0.73
2:B:240[A]:MET:HE1	2:B:244:PHE:HE2	1.53	0.72
2:B:240[A]:MET:CE	2:B:244:PHE:HE2	2.07	0.68
2:B:337:LYS:HE2	2:B:391:LEU:HD11	1.78	0.66
2:B:143:SER:HG	2:B:144:PRO:HD3	1.67	0.59
2:B:65:ILE:CD1	2:B:73:LEU:HD23	2.29	0.58
2:B:136:ALA:O	2:B:140:GLU:HG3	2.06	0.56
2:B:240[A]:MET:HE1	2:B:244:PHE:CE2	2.39	0.56
2:B:297:SER:HB3	2:B:305:ASP:OD1	2.06	0.55
1:A:178:SER:HA	1:A:193:LEU:HD11	1.90	0.54
1:A:6:ASN:O	1:A:10:GLN:HG3	2.07	0.54
2:B:109:GLU:HG3	2:B:170[B]:CYS:SG	2.48	0.54
2:B:30:GLU:HG2	5:B:2051:HOH:O	2.09	0.53
1:A:112:ASP:OD1	1:A:146:HIS:HE1	1.91	0.53
1:A:211:GLY:O	1:A:212:PHE:CB	2.57	0.52
2:B:284:ALA:O	2:B:286:MET:HE3	2.10	0.51
1:A:58:LEU:HD21	5:B:2035:HOH:O	2.11	0.49
2:B:61:LYS:HE3	2:B:63:GLN:NE2	2.27	0.49

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:260:HIS:HD2	5:B:2146:HOH:O	1.95	0.49
2:B:337:LYS:HE2	2:B:391:LEU:CD1	2.44	0.47
1:A:194:HIS:O	1:A:198:GLU:HG3	2.15	0.46
1:A:178:SER:HB2	1:A:210:GLN:HE21	1.81	0.46
2:B:257:PRO:HG3	2:B:304:LEU:HB3	1.96	0.46
1:A:87:LEU:O	1:A:87:LEU:HD12	2.17	0.45
1:A:27:ASP:HA	1:A:28:PRO:HA	1.76	0.44
1:A:133:VAL:HG12	1:A:152:PHE:CD1	2.53	0.43
2:B:11:GLU:HG2	2:B:11:GLU:O	2.19	0.43
2:B:193:GLY:HA2	2:B:280:PHE:O	2.19	0.42
1:A:95:ILE:HA	1:A:96:PRO:HD3	1.95	0.42
1:A:239:LYS:O	1:A:243:LYS:HG3	2.20	0.42
1:A:16:GLU:HG3	1:A:267:ARG:HA	2.01	0.41
1:A:58:LEU:HD11	5:B:2035:HOH:O	2.20	0.41
2:B:379:ARG:HD3	2:B:379:ARG:HH11	1.67	0.41
2:B:300:ILE:HD11	2:B:390:ILE:CD1	2.51	0.41
2:B:103:LYS:NZ	2:B:181:TYR:O	2.43	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	251/268 (94%)	247 (98%)	3 (1%)	1 (0%)	34	21
2	B	395/396 (100%)	389 (98%)	6 (2%)	0	100	100
All	All	646/664 (97%)	636 (98%)	9 (1%)	1 (0%)	47	33

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	212	PHE

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	200/208 (96%)	200 (100%)	0	100	100
2	B	310/309 (100%)	306 (99%)	4 (1%)	69	62
All	All	510/517 (99%)	506 (99%)	4 (1%)	81	78

All (4) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
2	B	109	GLU
2	B	196	PRO
2	B	279	TYR
2	B	297	SER

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	194	HIS
2	B	63	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

5.6 Ligand geometry

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
4	PLP	B	500	2	15,15,16	1.50	3 (20%)	20,22,23	3.12	9 (45%)
3	IGP	A	300	-	20,20,20	2.86	7 (35%)	24,29,29	2.32	5 (20%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	PLP	B	500	2	-	0/6/6/8	0/1/1/1
3	IGP	A	300	-	-	1/10/14/14	0/2/2/2

All (10) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	300	IGP	C2-C3	9.02	1.62	1.53
3	A	300	IGP	CG-CD2	4.63	1.46	1.40
3	A	300	IGP	CZ3-CE3	3.82	1.45	1.36
3	A	300	IGP	CD1-NE1	2.82	1.42	1.36
3	A	300	IGP	C1-C2	2.59	1.55	1.51
3	A	300	IGP	CE2-NE1	-2.49	1.30	1.38
4	B	500	PLP	C5-C4	-2.36	1.37	1.40
4	B	500	PLP	P-O3P	-2.28	1.46	1.54
4	B	500	PLP	C3-C2	-2.12	1.38	1.40
3	A	300	IGP	CH2-CZ2	2.04	1.41	1.36

All (14) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	300	IGP	CG-C3-C2	-8.96	98.16	112.14

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	B	500	PLP	O4P-C5A-C5	7.37	123.39	109.35
4	B	500	PLP	C3-C4-C5	6.70	125.97	118.74
4	B	500	PLP	C6-C5-C4	-5.81	113.58	118.16
4	B	500	PLP	C4A-C4-C5	-3.78	117.04	120.94
4	B	500	PLP	C3-C2-N1	-3.44	116.32	120.77
4	B	500	PLP	O4P-P-O1P	-3.32	97.16	106.47
3	A	300	IGP	O2-C2-C3	-3.22	104.94	109.26
3	A	300	IGP	P-OP4-C1	-2.64	111.03	118.30
3	A	300	IGP	OP1-P-OP4	2.25	112.71	106.73
3	A	300	IGP	CG-CD1-NE1	-2.24	104.52	108.79
4	B	500	PLP	O3-C3-C4	2.08	123.59	118.10
4	B	500	PLP	C4A-C4-C3	-2.07	116.99	120.50
4	B	500	PLP	C6-N1-C2	2.02	122.91	119.17

There are no chirality outliers.


All (1) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	A	300	IGP	C2-C1-OP4-P

There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	254/268 (94%)	0.08	10 (3%) 39 33	11, 20, 36, 62	0
2	B	394/396 (99%)	-0.03	15 (3%) 40 35	9, 15, 35, 58	0
All	All	648/664 (97%)	0.01	25 (3%) 39 33	9, 17, 36, 62	0

All (25) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	195	HIS	4.4
2	B	162	GLY	4.3
2	B	393	ALA	4.0
2	B	392	LYS	3.8
2	B	160	HIS	3.8
1	A	194	HIS	3.4
2	B	140	GLU	3.2
2	B	136	ALA	2.9
2	B	390	ILE	2.8
1	A	49[A]	GLU	2.7
1	A	249	LYS	2.6
2	B	394	ARG	2.6
2	B	161	SER	2.5
2	B	385	PHE	2.4
1	A	198	GLU	2.3
2	B	68	GLY	2.3
2	B	164	ALA	2.3
2	B	159	VAL	2.2
1	A	250	GLN	2.2
1	A	100	LEU	2.1
2	B	395	GLY	2.1
1	A	109	ASN	2.1
2	B	137	LYS	2.1
1	A	13	ASP	2.0

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	243	LYS	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
3	IGP	A	300	19/19	0.95	0.20	18,20,24,24	19
4	PLP	B	500	15/16	0.99	0.10	10,12,14,15	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

Electron density around IGP A 300:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers [i](#)

There are no such residues in this entry.