



# wwPDB X-ray Structure Validation Summary Report ⓘ

May 28, 2020 – 08:21 pm BST

PDB ID : 1QRK  
Title : HUMAN FACTOR XIII WITH STRONTIUM BOUND IN THE ION SITE  
Authors : Fox, B.A.; Yee, V.C.; Pederson, L.C.; Le Trong, I.; Bishop, P.D.; Stenkamp, R.E.; Teller, D.C.  
Deposited on : 1999-06-14  
Resolution : 2.50 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Xtriage (Phenix)	:	1.13
EDS	:	2.11
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac	:	5.8.0158
CCP4	:	7.0.044 (Gargrove)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.11


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*



The reported resolution of this entry is 2.50 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	141614	5346 (2.50-2.50)
Ramachandran outliers	138981	5231 (2.50-2.50)
Sidechain outliers	138945	5233 (2.50-2.50)
RSRZ outliers	127900	4559 (2.50-2.50)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	731	 60% 32% . .
1	B	731	 56% 34% 6% .

## 2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 11492 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called PROTEIN (COAGULATION FACTOR XIII).

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	700	Total	C	N	O	S	0	0	0
			5610	3557	966	1061	26			
1	B	705	Total	C	N	O	S	0	0	0
			5650	3583	973	1068	26			

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	651	GLU	GLN	CONFLICT	UNP P00488
B	651	GLU	GLN	CONFLICT	UNP P00488

- Molecule 2 is STRONTIUM ION (three-letter code: SR) (formula: Sr).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	B	1	Total	Sr	0	0
			1	1		
2	A	1	Total	Sr	0	0
			1	1		


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	115	Total	O	0	0
			115	115		
3	B	115	Total	O	0	0
			115	115		


### 3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

#### • Molecule 1: PROTEIN (COAGULATION FACTOR XIII)



#### • Molecule 1: PROTEIN (COAGULATION FACTOR XIII)



D722	T639	I549	M474	W392	D271	V161	R77
V723	Q640	Y552	I477	S397	E272	T165	S81
I725	V641	A555	Q484	T398	L275	G168	I86
Q726	V642	R556	Q487	P399	N281	V169	I87
R727	G643	I557	E488	Q401	P289	L170	F88
ARG	W646	T561	E489	E401	D297	R171	S89
PRO	T647	K565	E490	S403	I300	T172	R90
SER	T653	K568	E491	G405	E301	R174	P91
MET	P654	F568	L492	Y406	L182	R95	
	P655	K569	E495	Y407	L183	R96	
	L656	D574	L498	R408	F184	D97	
	R657	L577	A502	C409	E306	L98	
	T659	L578	P505	G410	W187	V101	
	L660	F579	L506	P411	R310	V104	
	R661	E578	N507	Q415	E189	I105	
	W664	L580	THR	V422	A192	Q110	
	V665	S581	GLU	Q425	V193	E111	
	W666	F582	GLY	F426	E198	N112	
	L667	K583	VAL	D427	E202	K113	
	W676	E585	ARG	A428	E216	G114	
			MET	P429	V217	T115	
			L588	F430	L206	P118	
			S589	V431	V211	V122	
			Q590	E434	I212	L125	
			A591	W435	F213	G128	
			Y594	N436	E216	G131	
			L599	S437	N218	A132	
			E600	D438	D219	R133	
			S603	L439	I220	I134	
			L604	I440	R221	V135	
				Y441	R223	D139	
				I442	Y227	R140	
				A444	F230	S141	
				K445	L235	R143	
				K446	K362	L144	
				V451	K363	S145	
				V452	K366	I146	
				V455	W370	Q147	
				D456	E377	S148	
				H459	R282	P150	
				I460	S259	I153	
				G461	V261	V154	
				K462	N267	V154	
				L463	A268	R158	
				L464	K269		
				I464	R270		
				W465			
				T466			
				R467			
				K467			
				T548			

## 4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	101.90Å 72.32Å 135.03Å 90.00° 105.90° 90.00°	Depositor
Resolution (Å)	10.00 – 2.50 72.32 – 2.20	Depositor EDS
% Data completeness (in resolution range)	78.5 (10.00-2.50) 62.6 (72.32-2.20)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.82 (at 2.20Å)	Xtriage
Refinement program	X-PLOR 3.1	Depositor
R, $R_{free}$	0.183 , 0.275 0.182 , (Not available)	Depositor DCC
$R_{free}$ test set	No test flags present.	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	30.4	Xtriage
Anisotropy	0.239	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.28 , 95.3	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.47$ , $\langle L^2 \rangle = 0.29$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.94	EDS
Total number of atoms	11492	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	33.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.96% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: SR

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.56	0/5741	0.79	1/7792 (0.0%)
1	B	0.60	0/5782	0.81	3/7847 (0.0%)
All	All	0.58	0/11523	0.80	4/15639 (0.0%)

There are no bond length outliers.

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	354	LEU	CA-CB-CG	5.76	128.56	115.30
1	B	170	LEU	N-CA-C	-5.73	95.53	111.00
1	B	588	LEU	CA-CB-CG	5.51	127.97	115.30
1	B	26	LEU	CA-CB-CG	5.05	126.91	115.30

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	5610	0	5467	166	0
1	B	5650	0	5495	189	0
2	A	1	0	0	0	0
2	B	1	0	0	0	0
3	A	115	0	0	1	0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	B	115	0	0	4	0
All	All	11492	0	10962	353	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 16.

The worst 5 of 353 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:331:PRO:HG2	1:A:379:TRP:HB3	1.46	0.97
1:B:544:HIS:HA	1:B:579:PRO:HB3	1.52	0.90
1:B:44:PHE:O	1:B:45:LEU:HB2	1.73	0.87
1:A:709:MET:HB3	1:A:717:VAL:HB	1.58	0.85
1:B:591:ALA:HA	1:B:594:TYR:CE2	2.15	0.82

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	694/731 (95%)	631 (91%)	52 (8%)	11 (2%)	9	17
1	B	699/731 (96%)	613 (88%)	71 (10%)	15 (2%)	7	11
All	All	1393/1462 (95%)	1244 (89%)	123 (9%)	26 (2%)	8	13

5 of 26 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	55	GLU
1	A	281	ASN
1	B	45	LEU

*Continued on next page...*



*Continued from previous page...*

Mol	Chain	Res	Type
1	B	53	PHE
1	B	55	GLU

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	616/644 (96%)	559 (91%)	57 (9%)	9	17
1	B	619/644 (96%)	539 (87%)	80 (13%)	4	8
All	All	1235/1288 (96%)	1098 (89%)	137 (11%)	6	11

5 of 137 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	47	VAL
1	B	193	VAL
1	B	617	ASP
1	B	59	THR
1	B	140	ARG

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 18 such sidechains are listed below:

Mol	Chain	Res	Type
1	A	726	GLN
1	B	46	ASN
1	B	468	GLN
1	A	686	ASN
1	A	716	HIS

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 5.6 Ligand geometry [i](#)

Of 2 ligands modelled in this entry, 2 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	700/731 (95%)	-0.73	2 (0%) 94 94	8, 31, 66, 83	0
1	B	705/731 (96%)	-0.71	1 (0%) 95 96	4, 28, 71, 85	0
All	All	1405/1462 (96%)	-0.72	3 (0%) 95 95	4, 29, 68, 85	0

All (3) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	577	LEU	2.8
1	A	168	GLY	2.5
1	A	167	TYR	2.2

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

### 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
2	SR	B	732	1/1	0.99	0.14	45,45,45,45	0
2	SR	A	732	1/1	0.99	0.17	51,51,51,51	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.