



wwPDB X-ray Structure Validation Summary Report ⓘ

Aug 21, 2020 – 02:00 PM BST

PDB ID : 6QS1
Title : Crystal structure of human Angiotensin-1 converting enzyme N-domain in complex with BPPb
Authors : Cozier, G.E.; Acharya, K.R.
Deposited on : 2019-02-20
Resolution : 1.80 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.13.1
buster-report : 1.1.7 (2018)
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.13.1


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.80 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.






Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	5950 (1.80-1.80)
Clashscore	141614	6793 (1.80-1.80)
Ramachandran outliers	138981	6697 (1.80-1.80)
Sidechain outliers	138945	6696 (1.80-1.80)
RSRZ outliers	127900	5850 (1.80-1.80)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	629	<div> <div>2%</div> <div> <div></div> <div>88%</div> <div>9%</div> <div></div> </div> </div>
1	B	629	<div> <div>7%</div> <div> <div></div> <div>84%</div> <div>11%</div> <div></div> </div> </div>
2	E	11	<div> <div>55%</div> <div> <div></div> <div>64%</div> <div>27%</div> <div>9%</div> </div> </div>
2	F	11	<div> <div></div> <div> <div>36%</div> <div>9%</div> <div>55%</div> </div> </div>
3	C	2	<div> <div></div> <div> <div>50%</div> <div>50%</div> </div> </div>
4	D	4	<div> <div></div> <div>100%</div> </div>

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
5	G	3	 33% 33% 33%
5	H	3	 100%
6	I	2	 100%

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
9	PEG	B	709	-	-	X	-

2 Entry composition [i](#)

There are 16 unique types of molecules in this entry. The entry contains 21210 atoms, of which 10081 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Angiotensin-converting enzyme.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	608	Total	C	H	N	O	S	0	19	0
			9851	3230	4811	863	928	19			
1	B	603	Total	C	H	N	O	S	0	17	0
			9776	3203	4785	858	911	19			

There are 18 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	9	GLN	ASN	engineered mutation	UNP P12821
A	25	GLN	ASN	engineered mutation	UNP P12821
A	82	GLN	ASN	engineered mutation	UNP P12821
A	117	GLN	ASN	engineered mutation	UNP P12821
A	131	GLN	ASN	engineered mutation	UNP P12821
A	289	GLN	ASN	engineered mutation	UNP P12821
A	545	ARG	GLN	engineered mutation	UNP P12821
A	576	LEU	PRO	engineered mutation	UNP P12821
A	629	LEU	-	expression tag	UNP P12821
B	9	GLN	ASN	engineered mutation	UNP P12821
B	25	GLN	ASN	engineered mutation	UNP P12821
B	82	GLN	ASN	engineered mutation	UNP P12821
B	117	GLN	ASN	engineered mutation	UNP P12821
B	131	GLN	ASN	engineered mutation	UNP P12821
B	289	GLN	ASN	engineered mutation	UNP P12821
B	545	ARG	GLN	engineered mutation	UNP P12821
B	576	LEU	PRO	engineered mutation	UNP P12821
B	629	LEU	-	expression tag	UNP P12821

- Molecule 2 is a protein called Bradykinin potentiating peptide b.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	E	11	Total	C	H	N	O	0	0	0
			176	56	92	15	13			

Continued on next page...

Continued from previous page...


Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	F	5	Total	C	H	N	O	0	0	0
			84	27	45	6	6			

- Molecule 3 is an oligosaccharide called 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.




Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	C	2	Total	C	H	N	O	0	0	0
			55	16	27	2	10			

- Molecule 4 is an oligosaccharide called beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-[alpha-L-fucopyranose-(1-6)]2-acetamido-2-deoxy-beta-D-glucopyranose.



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
4	D	4	Total	C	H	N	O	0	0	0
			95	28	46	2	19			

- Molecule 5 is an oligosaccharide called 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-[alpha-L-fucopyranose-(1-6)]2-acetamido-2-deoxy-beta-D-glucopyranose.




Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
5	G	3	Total	C	H	N	O	0	0	0
			73	22	35	2	14			
5	H	3	Total	C	H	N	O	0	0	0
			74	22	36	2	14			

- Molecule 6 is an oligosaccharide called alpha-L-fucopyranose-(1-6)-2-acetamido-2-deoxy-beta-D-glucopyranose.



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
6	I	2	Total	C	H	N	O	0	0	0
			47	14	23	1	9			

- Molecule 7 is 3,6,9,12,15,18,21,24,27-NONAOXANONACOSANE-1,29-DIOL (three-letter code: XPE) (formula: $C_{20}H_{42}O_{11}$).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
7	A	1	Total	C	H	O	0	0
			73	20	42	11		

- Molecule 8 is 1,2-ETHANEDIOL (three-letter code: EDO) (formula: $C_2H_6O_2$).




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
8	A	1	Total	C	H	O	0	0
			10	2	6	2		
8	A	1	Total	C	H	O	0	0
			10	2	6	2		
8	A	1	Total	C	H	O	0	0
			10	2	6	2		
8	A	1	Total	C	H	O	0	0
			10	2	6	2		
8	B	1	Total	C	H	O	0	0
			10	2	6	2		
8	B	1	Total	C	H	O	0	0
			10	2	6	2		

- Molecule 9 is DI(HYDROXYETHYL)ETHER (three-letter code: PEG) (formula: C₄H₁₀O₃).




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
9	A	1	Total	C	H	O	0	0
			17	4	10	3		
9	A	1	Total	C	H	O	0	0
			17	4	10	3		
9	B	1	Total	C	H	O	0	0
			17	4	10	3		
9	B	1	Total	C	H	O	0	0
			17	4	10	3		

- Molecule 10 is BORIC ACID (three-letter code: BO3) (formula: BH_3O_3).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
10	A	1	Total B H O 7 1 3 3	0	0
10	A	1	Total B H O 7 1 3 3	0	0
10	A	1	Total B H O 7 1 3 3	0	0
10	A	1	Total B H O 7 1 3 3	0	0
10	B	1	Total B H O 7 1 3 3	0	0
10	B	1	Total B H O 7 1 3 3	0	0
10	B	1	Total B H O 7 1 3 3	0	0

- Molecule 11 is TRIETHYLENE GLYCOL (three-letter code: PGE) (formula: C₆H₁₄O₄).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
11	A	1	Total C H O 48 12 28 8	0	1

- Molecule 12 is ZINC ION (three-letter code: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
12	B	1	Total Zn 1 1	0	0
12	A	1	Total Zn 1 1	0	0


- Molecule 13 is CHLORIDE ION (three-letter code: CL) (formula: Cl).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
13	B	2	Total	Cl	0	0
			2	2		
13	A	2	Total	Cl	0	0
			2	2		

- Molecule 14 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
14	B	1	Total	Mg	0	0
			1	1		
14	A	1	Total	Mg	0	0
			1	1		

- Molecule 15 is 2-acetamido-2-deoxy-beta-D-glucopyranose (three-letter code: NAG) (formula: C₈H₁₅NO₆).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
15	B	1	Total	C	H	N	O	0	0
			28	8	14	1	5		

- Molecule 16 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
16	A	387	Total	O	0	3
			390	390		

Continued on next page...


Continued from previous page...

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
16	B	243	Total 245	O 245	0	2
16	E	6	Total 6	O 6	0	0
16	F	4	Total 4	O 4	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Angiotensin-converting enzyme




- Molecule 1: Angiotensin-converting enzyme



- Molecule 2: Bradykinin potentiating peptide b



- Molecule 2: Bradykinin potentiating peptide b

Chain F:  36% 9% 55%




- Molecule 3: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain C:  50% 50%




- Molecule 4: beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-[alpha-L-fucopyranose-(1-6)]2-acetamido-2-deoxy-beta-D-glucopyranose

Chain D:  100%



- Molecule 5: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-[alpha-L-fucopyranose-(1-6)]2-acetamido-2-deoxy-beta-D-glucopyranose

Chain G:  33% 33% 33%



- Molecule 5: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-[alpha-L-fucopyranose-(1-6)]2-acetamido-2-deoxy-beta-D-glucopyranose

Chain H:  100%



- Molecule 6: alpha-L-fucopyranose-(1-6)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain I:  100%



4 Data and refinement statistics

Property	Value	Source
Space group	P 1	Depositor
Cell constants a, b, c, α , β , γ	72.68Å 77.07Å 82.36Å 88.86° 64.59° 75.03°	Depositor
Resolution (Å)	74.05 – 1.80 74.05 – 1.80	Depositor EDS
% Data completeness (in resolution range)	98.6 (74.05-1.80) 98.7 (74.05-1.80)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.59 (at 1.80Å)	Xtriage
Refinement program	PHENIX (dev_3409: ???)	Depositor
R, R_{free}	0.202 , 0.232 0.201 , 0.232	Depositor DCC
R_{free} test set	1818 reflections (1.28%)	wwPDB-VP
Wilson B-factor (Å ²)	29.8	Xtriage
Anisotropy	0.037	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.41 , 48.3	EDS
L-test for twinning ²	$\langle L \rangle = 0.51$, $\langle L^2 \rangle = 0.34$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	21210	wwPDB-VP
Average B, all atoms (Å ²)	45.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.97% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: MG, BMA, NAG, CL, BO3, ZN, XPE, EDO, PGE, FUC, PEG, PCA

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.58	0/5271	0.68	0/7178
1	B	0.55	0/5220	0.65	0/7109
2	E	0.49	0/81	0.57	0/112
2	F	0.53	0/41	0.50	0/54
All	All	0.56	0/10613	0.66	0/14453

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	5040	4811	4741	42	0
1	B	4991	4785	4710	49	0
2	E	84	92	92	4	0
2	F	39	45	45	1	0
3	C	28	27	25	0	0
4	D	49	46	43	0	0
5	G	38	35	34	1	0
5	H	38	36	34	0	0
6	I	24	23	22	2	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
7	A	31	42	42	3	0
8	A	16	24	24	1	0
8	B	8	12	12	0	0
9	A	14	20	20	0	0
9	B	14	20	20	6	0
10	A	16	12	12	0	0
10	B	12	9	9	1	0
11	A	20	28	28	4	0
12	A	1	0	0	0	0
12	B	1	0	0	0	0
13	A	2	0	0	0	0
13	B	2	0	0	1	0
14	A	1	0	0	0	0
14	B	1	0	0	0	0
15	B	14	14	13	0	0
16	A	390	0	0	9	0
16	B	245	0	0	6	0
16	E	6	0	0	0	0
16	F	4	0	0	0	0
All	All	11129	10081	9926	98	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

The worst 5 of 98 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:564:ALA:HB1	9:B:709:PEG:H42	1.62	0.82
1:A:66:GLU:OE2	16:A:801:HOH:O	1.99	0.79
1:A:157:LEU:HD11	1:A:477:VAL:HG13	1.66	0.77
10:B:712:BO3:O2	16:B:801:HOH:O	2.00	0.77
1:B:377:VAL:HA	1:B:380:ARG:HG3	1.71	0.72

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	623/629 (99%)	611 (98%)	12 (2%)	0	100	100
1	B	616/629 (98%)	599 (97%)	17 (3%)	0	100	100
2	E	9/11 (82%)	8 (89%)	1 (11%)	0	100	100
2	F	3/11 (27%)	3 (100%)	0	0	100	100
All	All	1251/1280 (98%)	1221 (98%)	30 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	540/541 (100%)	536 (99%)	4 (1%)	84	81
1	B	534/541 (99%)	527 (99%)	7 (1%)	69	62
2	E	9/9 (100%)	9 (100%)	0	100	100
2	F	5/9 (56%)	5 (100%)	0	100	100
All	All	1088/1100 (99%)	1077 (99%)	11 (1%)	76	71

5 of 11 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	35	SER
1	B	129	LEU
1	B	368	TYR

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	388	HIS
1	B	326	ARG

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

1 non-standard protein/DNA/RNA residue is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	PCA	E	1	2	7,8,9	2.46	2 (28%)	9,10,12	2.04	5 (55%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	PCA	E	1	2	-	0/0/11/13	0/1/1/1

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	E	1	PCA	CD-N	4.85	1.47	1.34
2	E	1	PCA	CA-N	4.12	1.51	1.46

All (5) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	E	1	PCA	CA-N-CD	-3.02	103.24	113.58
2	E	1	PCA	CB-CA-N	2.67	110.97	103.30
2	E	1	PCA	CB-CA-C	-2.47	109.30	112.70
2	E	1	PCA	OE-CD-CG	-2.41	122.56	126.76
2	E	1	PCA	CG-CD-N	2.16	113.98	108.39

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	E	1	PCA	1	0

5.5 Carbohydrates ⓘ

14 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	NAG	C	1	1,3	14,14,15	0.41	0	17,19,21	0.52	0
3	NAG	C	2	3	14,14,15	0.20	0	17,19,21	0.74	1 (5%)
4	NAG	D	1	1,4	14,14,15	0.61	1 (7%)	17,19,21	0.61	0
4	NAG	D	2	4	14,14,15	0.40	0	17,19,21	0.75	1 (5%)
4	BMA	D	3	4	11,11,12	1.00	1 (9%)	15,15,17	0.77	0
4	FUC	D	4	4	10,10,11	0.95	1 (10%)	14,14,16	1.21	2 (14%)
5	NAG	G	1	1,5	14,14,15	0.53	0	17,19,21	0.86	1 (5%)
5	NAG	G	2	5	14,14,15	0.45	0	17,19,21	0.46	0
5	FUC	G	3	5	10,10,11	1.07	1 (10%)	14,14,16	1.36	2 (14%)
5	NAG	H	1	1,5	14,14,15	0.42	0	17,19,21	0.53	0
5	NAG	H	2	5	14,14,15	0.46	0	17,19,21	0.43	0
5	FUC	H	3	5	10,10,11	0.74	0	14,14,16	0.96	0
6	NAG	I	1	1,6	14,14,15	0.78	1 (7%)	17,19,21	0.86	1 (5%)

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
6	FUC	I	2	6	10,10,11	0.85	0	14,14,16	1.83	3 (21%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	NAG	C	1	1,3	-	0/6/23/26	0/1/1/1
3	NAG	C	2	3	-	0/6/23/26	0/1/1/1
4	NAG	D	1	1,4	-	0/6/23/26	0/1/1/1
4	NAG	D	2	4	-	0/6/23/26	0/1/1/1
4	BMA	D	3	4	-	0/2/19/22	0/1/1/1
4	FUC	D	4	4	-	-	0/1/1/1
5	NAG	G	1	1,5	-	3/6/23/26	0/1/1/1
5	NAG	G	2	5	-	2/6/23/26	0/1/1/1
5	FUC	G	3	5	-	-	0/1/1/1
5	NAG	H	1	1,5	-	2/6/23/26	0/1/1/1
5	NAG	H	2	5	-	2/6/23/26	0/1/1/1
5	FUC	H	3	5	-	-	0/1/1/1
6	NAG	I	1	1,6	-	2/6/23/26	0/1/1/1
6	FUC	I	2	6	-	-	0/1/1/1

All (5) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	D	4	FUC	O5-C1	-2.49	1.39	1.43
5	G	3	FUC	C1-C2	2.34	1.57	1.52
4	D	1	NAG	O5-C1	-2.04	1.40	1.43
4	D	3	BMA	C2-C3	2.01	1.55	1.52
6	I	1	NAG	C1-C2	2.00	1.55	1.52

The worst 5 of 11 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
6	I	2	FUC	C1-C2-C3	4.29	114.94	109.67
6	I	2	FUC	O5-C1-C2	3.46	116.11	110.77
5	G	3	FUC	O2-C2-C1	3.22	115.75	109.15
6	I	1	NAG	C1-O5-C5	3.02	116.29	112.19
6	I	2	FUC	C1-O5-C5	2.77	119.05	112.78

There are no chirality outliers.

5 of 11 torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
6	I	1	NAG	O5-C5-C6-O6
6	I	1	NAG	C4-C5-C6-O6
5	G	1	NAG	O5-C5-C6-O6
5	G	1	NAG	C4-C5-C6-O6
5	H	1	NAG	O5-C5-C6-O6


There are no ring outliers.

3 monomers are involved in 3 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
6	I	2	FUC	1	0
5	G	3	FUC	1	0
6	I	1	NAG	2	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.





5.6 Ligand geometry

Of 29 ligands modelled in this entry, 8 are monoatomic - leaving 21 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
9	PEG	B	709	-	6,6,6	0.52	0	5,5,5	0.43	0
10	BO3	A	716	-	3,3,3	0.18	0	3,3,3	0.51	0
8	EDO	A	711	-	3,3,3	0.55	0	2,2,2	0.24	0
10	BO3	B	710	-	3,3,3	0.26	0	3,3,3	0.48	0
9	PEG	B	708	-	6,6,6	0.53	0	5,5,5	0.41	0
7	XPE	A	710	-	30,30,30	0.59	0	29,29,29	0.67	0
11	PGE	A	720[A]	-	9,9,9	0.36	0	8,8,8	0.47	0
8	EDO	A	719	-	3,3,3	0.49	0	2,2,2	0.66	0
11	PGE	A	720[B]	-	9,9,9	0.30	0	8,8,8	0.66	0
15	NAG	B	701	1	14,14,15	0.53	0	17,19,21	0.55	0
9	PEG	A	713	-	6,6,6	0.51	0	5,5,5	0.44	0
10	BO3	A	721	-	3,3,3	0.20	0	3,3,3	0.37	0
8	EDO	B	707	-	3,3,3	0.56	0	2,2,2	0.10	0
9	PEG	A	717	-	6,6,6	0.43	0	5,5,5	0.48	0
10	BO3	A	715	-	3,3,3	0.41	0	3,3,3	0.64	0
10	BO3	B	711	-	3,3,3	0.21	0	3,3,3	0.16	0
8	EDO	A	712	-	3,3,3	0.44	0	2,2,2	0.47	0
10	BO3	B	712	-	3,3,3	0.44	0	3,3,3	0.26	0
8	EDO	A	718	-	3,3,3	0.54	0	2,2,2	0.46	0
10	BO3	A	714	-	3,3,3	0.66	0	3,3,3	0.70	0
8	EDO	B	713	-	3,3,3	0.47	0	2,2,2	0.41	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
9	PEG	B	709	-	-	4/4/4/4	-
8	EDO	A	711	-	-	0/1/1/1	-
7	XPE	A	710	-	-	15/28/28/28	-

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
9	PEG	B	708	-	-	2/4/4/4	-
11	PGE	A	720[A]	-	-	3/7/7/7	-
8	EDO	A	719	-	-	0/1/1/1	-
11	PGE	A	720[B]	-	-	6/7/7/7	-
15	NAG	B	701	1	-	3/6/23/26	0/1/1/1
9	PEG	A	713	-	-	2/4/4/4	-
9	PEG	A	717	-	-	3/4/4/4	-
8	EDO	A	718	-	-	1/1/1/1	-
8	EDO	A	712	-	-	1/1/1/1	-
8	EDO	B	707	-	-	0/1/1/1	-
8	EDO	B	713	-	-	1/1/1/1	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

5 of 41 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
11	A	720[A]	PGE	O2-C3-C4-O3
7	A	710	XPE	O7-C8-C9-O10
9	B	709	PEG	O1-C1-C2-O2
11	A	720[B]	PGE	O3-C5-C6-O4
9	B	708	PEG	O1-C1-C2-O2


There are no ring outliers.

6 monomers are involved in 15 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
9	B	709	PEG	4	0
9	B	708	PEG	2	0
7	A	710	XPE	3	0
11	A	720[A]	PGE	4	0
10	B	712	BO3	1	0
8	A	718	EDO	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is

within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	608/629 (96%)	0.23	10 (1%) 72 68	22, 35, 54, 78	0
1	B	603/629 (95%)	0.52	41 (6%) 17 13	25, 43, 66, 88	0
2	E	10/11 (90%)	2.04	6 (60%) 0 0	27, 44, 55, 55	10 (100%)
2	F	5/11 (45%)	0.56	0 100 100	30, 34, 44, 47	5 (100%)
All	All	1226/1280 (95%)	0.39	57 (4%) 32 26	22, 38, 63, 88	15 (1%)

The worst 5 of 57 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	E	3	LEU	7.0
1	A	325	GLY	6.8
1	B	105	LEU	5.7
1	B	1	LEU	5.0
1	B	101[A]	ALA	4.6

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	PCA	E	1	8/9	0.82	0.26	45,55,60,65	14

6.3 Carbohydrates [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum,




median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
5	NAG	G	2	14/15	0.60	0.27	60,73,93,103	27
4	BMA	D	3	11/12	0.69	0.17	65,76,88,92	0
3	NAG	C	2	14/15	0.70	0.24	61,81,98,103	0
5	FUC	G	3	10/11	0.73	0.26	54,65,82,91	0
6	FUC	I	2	10/11	0.77	0.29	59,71,86,94	20
3	NAG	C	1	14/15	0.80	0.15	49,61,74,78	0
5	NAG	H	2	14/15	0.82	0.29	62,71,86,86	28
5	FUC	H	3	10/11	0.83	0.25	64,71,81,83	20
6	NAG	I	1	14/15	0.87	0.14	49,60,74,86	0
5	NAG	H	1	14/15	0.87	0.20	57,67,81,81	0
4	FUC	D	4	10/11	0.89	0.15	42,50,60,63	20
4	NAG	D	2	14/15	0.89	0.18	46,58,75,75	27
5	NAG	G	1	14/15	0.89	0.12	43,55,67,68	0
4	NAG	D	1	14/15	0.94	0.11	38,46,58,58	0

The following is a graphical depiction of the model fit to experimental electron density for oligosaccharide. Each fit is shown from different orientation to approximate a three-dimensional view.




Electron density around Chain G:


$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



Electron density around Chain H:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)





6.4 Ligands ⓘ

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.


Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
15	NAG	B	701	14/15	0.52	0.23	66,78,93,94	28
8	EDO	A	718	4/4	0.68	0.21	52,63,68,77	0
10	BO3	A	721	4/4	0.72	0.28	45,54,65,73	0
8	EDO	A	711	4/4	0.77	0.12	51,61,64,66	0
10	BO3	B	711	4/4	0.77	0.19	46,55,66,66	7
8	EDO	B	713	4/4	0.77	0.29	66,80,84,86	0
8	EDO	B	707	4/4	0.83	0.10	54,65,70,75	0
8	EDO	A	712	4/4	0.84	0.10	52,63,66,70	0
7	XPE	A	710	31/31	0.84	0.14	39,55,67,74	0
10	BO3	A	715	4/4	0.85	0.12	30,33,39,40	7
11	PGE	A	720[B]	10/10	0.86	0.19	29,37,44,44	24
11	PGE	A	720[A]	10/10	0.86	0.19	28,37,44,45	24
9	PEG	A	717	7/7	0.86	0.18	46,56,61,66	17
10	BO3	B	710	4/4	0.88	0.17	36,42,49,52	7

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
9	PEG	B	709	7/7	0.89	0.14	44,53,59,59	17
10	BO3	A	716	4/4	0.90	0.19	40,42,50,53	7
10	BO3	B	712	4/4	0.90	0.11	50,55,66,67	0
9	PEG	B	708	7/7	0.92	0.24	31,39,46,47	17
13	CL	B	716	1/1	0.92	0.08	44,44,44,44	0
8	EDO	A	719	4/4	0.92	0.12	44,53,60,61	10
9	PEG	A	713	7/7	0.93	0.10	44,53,62,71	0
10	BO3	A	714	4/4	0.94	0.13	37,43,51,54	7
14	MG	A	725	1/1	0.94	0.11	38,38,38,38	1
14	MG	B	717	1/1	0.94	0.13	47,47,47,47	0
13	CL	B	715	1/1	0.98	0.13	32,32,32,32	0
12	ZN	B	714	1/1	0.99	0.11	29,29,29,29	1
13	CL	A	724	1/1	1.00	0.08	36,36,36,36	0
13	CL	A	723	1/1	1.00	0.14	25,25,25,25	0
12	ZN	A	722	1/1	1.00	0.13	30,30,30,30	1

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.



6.5 Other polymers [i](#)

There are no such residues in this entry.