



Full wwPDB X-ray Structure Validation Report ⓘ

Jan 30, 2021 – 09:08 PM EST

PDB ID : 3QU2
Title : Crystal structure of pyrophosphatase from bacteroides thetaiotaomicron, a closed cap conformation
Authors : Patskovsky, Y.; Huang, H.; Toro, R.; Gerlt, J.A.; Burley, S.K.; Dunaway-Mariano, D.; Almo, S.C.; New York SGX Research Center for Structural Genomics (NYSGXRC); Enzyme Function Initiative (EFI)
Deposited on : 2011-02-23
Resolution : 1.95 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix)	:	1.13
EDS	:	2.16
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac	:	5.8.0158
CCP4	:	7.0.044 (Gargrove)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.16


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.95 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	2580 (1.96-1.96)
Clashscore	141614	2705 (1.96-1.96)
Ramachandran outliers	138981	2678 (1.96-1.96)
Sidechain outliers	138945	2678 (1.96-1.96)
RSRZ outliers	127900	2539 (1.96-1.96)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	243	<div> <div>84%</div> <div>7%</div> <div>9%</div> </div>
1	B	243	<div> <div>87%</div> <div>9%</div> </div>
1	C	243	<div> <div>88%</div> <div>9%</div> </div>
1	D	243	<div> <div>84%</div> <div>7%</div> <div>9%</div> </div>

2 Entry composition

There are 6 unique types of molecules in this entry. The entry contains 7744 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called INORGANIC PYROPHOSPHATASE.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	221	Total	C	N	O	S	0	9	0
			1761	1132	284	333	12			
1	B	221	Total	C	N	O	S	0	10	0
			1768	1133	284	339	12			
1	C	221	Total	C	N	O	S	0	4	0
			1742	1117	285	328	12			
1	D	221	Total	C	N	O	S	0	8	0
			1758	1130	285	329	14			

There are 76 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-18	GLY	-	expression tag	UNP Q8A5V9
A	-17	SER	-	expression tag	UNP Q8A5V9
A	-16	SER	-	expression tag	UNP Q8A5V9
A	-15	HIS	-	expression tag	UNP Q8A5V9
A	-14	HIS	-	expression tag	UNP Q8A5V9
A	-13	HIS	-	expression tag	UNP Q8A5V9
A	-12	HIS	-	expression tag	UNP Q8A5V9
A	-11	HIS	-	expression tag	UNP Q8A5V9
A	-10	HIS	-	expression tag	UNP Q8A5V9
A	-9	SER	-	expression tag	UNP Q8A5V9
A	-8	SER	-	expression tag	UNP Q8A5V9
A	-7	GLY	-	expression tag	UNP Q8A5V9
A	-6	LEU	-	expression tag	UNP Q8A5V9
A	-5	VAL	-	expression tag	UNP Q8A5V9
A	-4	PRO	-	expression tag	UNP Q8A5V9
A	-3	ARG	-	expression tag	UNP Q8A5V9
A	-2	GLY	-	expression tag	UNP Q8A5V9
A	-1	SER	-	expression tag	UNP Q8A5V9
A	0	HIS	-	expression tag	UNP Q8A5V9
B	-18	GLY	-	expression tag	UNP Q8A5V9
B	-17	SER	-	expression tag	UNP Q8A5V9

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
B	-16	SER	-	expression tag	UNP Q8A5V9
B	-15	HIS	-	expression tag	UNP Q8A5V9
B	-14	HIS	-	expression tag	UNP Q8A5V9
B	-13	HIS	-	expression tag	UNP Q8A5V9
B	-12	HIS	-	expression tag	UNP Q8A5V9
B	-11	HIS	-	expression tag	UNP Q8A5V9
B	-10	HIS	-	expression tag	UNP Q8A5V9
B	-9	SER	-	expression tag	UNP Q8A5V9
B	-8	SER	-	expression tag	UNP Q8A5V9
B	-7	GLY	-	expression tag	UNP Q8A5V9
B	-6	LEU	-	expression tag	UNP Q8A5V9
B	-5	VAL	-	expression tag	UNP Q8A5V9
B	-4	PRO	-	expression tag	UNP Q8A5V9
B	-3	ARG	-	expression tag	UNP Q8A5V9
B	-2	GLY	-	expression tag	UNP Q8A5V9
B	-1	SER	-	expression tag	UNP Q8A5V9
B	0	HIS	-	expression tag	UNP Q8A5V9
C	-18	GLY	-	expression tag	UNP Q8A5V9
C	-17	SER	-	expression tag	UNP Q8A5V9
C	-16	SER	-	expression tag	UNP Q8A5V9
C	-15	HIS	-	expression tag	UNP Q8A5V9
C	-14	HIS	-	expression tag	UNP Q8A5V9
C	-13	HIS	-	expression tag	UNP Q8A5V9
C	-12	HIS	-	expression tag	UNP Q8A5V9
C	-11	HIS	-	expression tag	UNP Q8A5V9
C	-10	HIS	-	expression tag	UNP Q8A5V9
C	-9	SER	-	expression tag	UNP Q8A5V9
C	-8	SER	-	expression tag	UNP Q8A5V9
C	-7	GLY	-	expression tag	UNP Q8A5V9
C	-6	LEU	-	expression tag	UNP Q8A5V9
C	-5	VAL	-	expression tag	UNP Q8A5V9
C	-4	PRO	-	expression tag	UNP Q8A5V9
C	-3	ARG	-	expression tag	UNP Q8A5V9
C	-2	GLY	-	expression tag	UNP Q8A5V9
C	-1	SER	-	expression tag	UNP Q8A5V9
C	0	HIS	-	expression tag	UNP Q8A5V9
D	-18	GLY	-	expression tag	UNP Q8A5V9
D	-17	SER	-	expression tag	UNP Q8A5V9
D	-16	SER	-	expression tag	UNP Q8A5V9
D	-15	HIS	-	expression tag	UNP Q8A5V9
D	-14	HIS	-	expression tag	UNP Q8A5V9
D	-13	HIS	-	expression tag	UNP Q8A5V9

Continued on next page...


Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
D	-12	HIS	-	expression tag	UNP Q8A5V9
D	-11	HIS	-	expression tag	UNP Q8A5V9
D	-10	HIS	-	expression tag	UNP Q8A5V9
D	-9	SER	-	expression tag	UNP Q8A5V9
D	-8	SER	-	expression tag	UNP Q8A5V9
D	-7	GLY	-	expression tag	UNP Q8A5V9
D	-6	LEU	-	expression tag	UNP Q8A5V9
D	-5	VAL	-	expression tag	UNP Q8A5V9
D	-4	PRO	-	expression tag	UNP Q8A5V9
D	-3	ARG	-	expression tag	UNP Q8A5V9
D	-2	GLY	-	expression tag	UNP Q8A5V9
D	-1	SER	-	expression tag	UNP Q8A5V9
D	0	HIS	-	expression tag	UNP Q8A5V9

- Molecule 2 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).


Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	B	1	Total Mg 1 1	0	0
2	A	1	Total Mg 1 1	0	0
2	D	1	Total Mg 1 1	0	0
2	C	1	Total Mg 1 1	0	0

- Molecule 3 is GLYCEROL (three-letter code: GOL) (formula: C₃H₈O₃).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	C	O	0	0
			6	3	3		
3	B	1	Total	C	O	0	0
			6	3	3		
3	B	1	Total	C	O	0	0
			6	3	3		
3	C	1	Total	C	O	0	0
			6	3	3		
3	C	1	Total	C	O	0	0
			6	3	3		
3	D	1	Total	C	O	0	0
			6	3	3		
3	D	1	Total	C	O	0	0
			6	3	3		
3	D	1	Total	C	O	0	0
			6	3	3		

- Molecule 4 is CITRIC ACID (three-letter code: CIT) (formula: $C_6H_8O_7$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	C	O	0	0
			13	6	7		

- Molecule 5 is CHLORIDE ION (three-letter code: CL) (formula: Cl).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	B	1	Total	Cl	0	0
			1	1		


- Molecule 6 is water.


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	167	Total 167	O 167	0	0
6	B	152	Total 152	O 152	0	0
6	C	161	Total 161	O 161	0	0
6	D	169	Total 169	O 169	0	0

3 Residue-property plots [i](#)


These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: INORGANIC PYROPHOSPHATASE

Chain A: 




• Molecule 1: INORGANIC PYROPHOSPHATASE

Chain B: 




• Molecule 1: INORGANIC PYROPHOSPHATASE

Chain C: 



• Molecule 1: INORGANIC PYROPHOSPHATASE

Chain D: 



4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, α , β , γ	137.60Å 71.76Å 114.53Å 90.00° 105.80° 90.00°	Depositor
Resolution (Å)	20.00 – 1.95 44.46 – 1.94	Depositor EDS
% Data completeness (in resolution range)	100.0 (20.00-1.95) 99.8 (44.46-1.94)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	0.12	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.28 (at 1.94Å)	Xtriage
Refinement program	REFMAC 5.3.0034	Depositor
R, R_{free}	0.232 , 0.275 0.234 , 0.275	Depositor DCC
R_{free} test set	2378 reflections (3.00%)	wwPDB-VP
Wilson B-factor (Å ²)	24.3	Xtriage
Anisotropy	0.319	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.36 , 57.4	EDS
L-test for twinning ²	$\langle L \rangle = 0.52$, $\langle L^2 \rangle = 0.35$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	7744	wwPDB-VP
Average B, all atoms (Å ²)	20.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 84.01 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 1.8669e-07. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: GOL, MG, CL, CIT

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.49	0/1827	0.58	0/2471
1	B	0.51	0/1837	0.61	0/2482
1	C	0.50	0/1793	0.60	0/2421
1	D	0.52	0/1822	0.63	0/2462
All	All	0.50	0/7279	0.61	0/9836

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1761	0	1771	7	0
1	B	1768	0	1768	4	0
1	C	1742	0	1744	4	0
1	D	1758	0	1766	13	0
2	A	1	0	0	0	0
2	B	1	0	0	0	0
2	C	1	0	0	0	0
2	D	1	0	0	0	0
3	A	6	0	8	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	B	12	0	16	0	0
3	C	12	0	16	0	0
3	D	18	0	24	2	0
4	A	13	0	5	1	0
5	B	1	0	0	0	0
6	A	167	0	0	0	0
6	B	152	0	0	0	0
6	C	161	0	0	0	0
6	D	169	0	0	3	0
All	All	7744	0	7118	29	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (29) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:D:140:ALA:HA	1:D:143:VAL:HG22	1.71	0.69
1:D:77[B]:HIS:CD2	6:D:685:HOH:O	2.50	0.64
1:D:77[B]:HIS:CE1	6:D:685:HOH:O	2.50	0.64
1:A:102[A]:VAL:HG13	1:A:107:LEU:HB2	1.88	0.56
1:D:117:GLN:HB3	1:D:120:LEU:HD13	1.88	0.55
1:D:191:ASN:HD22	3:D:227:GOL:H12	1.71	0.54
1:D:116:GLY:HA3	1:D:140:ALA:HB3	1.90	0.53
1:D:77[B]:HIS:CG	6:D:685:HOH:O	2.62	0.52
1:C:30:VAL:HG21	1:C:79:LYS:HA	1.92	0.50
4:A:227:CIT:H42	4:A:227:CIT:O2	2.11	0.49
1:A:30:VAL:HG21	1:A:79:LYS:HA	1.93	0.49
1:B:140:ALA:HA	1:B:143:VAL:HG22	1.95	0.48
1:D:30:VAL:HG21	1:D:79:LYS:HA	1.97	0.47
1:B:5:LEU:HD13	1:B:168[A]:VAL:HG23	1.95	0.47
1:D:98:LEU:HD13	1:D:216:CYS:HA	1.97	0.47
1:C:102:VAL:HG13	1:C:107:LEU:HB2	1.97	0.46
1:A:111:VAL:HB	1:A:137:MET:HG2	1.98	0.45
1:D:26:ALA:HA	1:D:82:LEU:HD23	1.99	0.45
1:D:171:GLU:O	1:D:189:ALA:HA	2.17	0.45
1:D:134:LYS:H	3:D:228:GOL:H32	1.82	0.44
1:A:5:LEU:HD11	1:A:168[B]:VAL:HG22	1.99	0.43
1:A:178:GLU:O	1:A:182:LYS:HG2	2.18	0.43
1:D:116:GLY:O	1:D:141:PHE:HE1	2.02	0.43
1:A:20:MET:HB3	1:A:44:TYR:CE2	2.54	0.42

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:118:LEU:O	1:B:122[A]:GLU:HG2	2.19	0.42
1:A:169:VAL:HB	1:A:187[B]:THR:HG22	2.01	0.42
1:B:102:VAL:HG13	1:B:107:LEU:HB2	2.03	0.41
1:C:98:LEU:HD13	1:C:216:CYS:HA	2.02	0.41
1:C:149:ASN:O	1:C:152:PRO:HD2	2.21	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	228/243 (94%)	225 (99%)	3 (1%)	0	100	100
1	B	229/243 (94%)	227 (99%)	2 (1%)	0	100	100
1	C	223/243 (92%)	218 (98%)	5 (2%)	0	100	100
1	D	227/243 (93%)	224 (99%)	3 (1%)	0	100	100
All	All	907/972 (93%)	894 (99%)	13 (1%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	194/204 (95%)	189 (97%)	5 (3%)	46	36

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	B	195/204 (96%)	192 (98%)	3 (2%)	65	60
1	C	189/204 (93%)	189 (100%)	0	100	100
1	D	193/204 (95%)	192 (100%)	1 (0%)	88	88
All	All	771/816 (94%)	762 (99%)	9 (1%)	78	68

All (9) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	66	GLU
1	A	182	LYS
1	A	190[A]	VAL
1	A	190[B]	VAL
1	A	224	LEU
1	B	9	LEU
1	B	151[A]	GLU
1	B	151[B]	GLU
1	D	9	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
1	D	133	HIS

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no monosaccharides in this entry.

5.6 Ligand geometry

Of 14 ligands modelled in this entry, 5 are monoatomic - leaving 9 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	GOL	B	226	-	5,5,5	0.22	0	5,5,5	0.84	0
3	GOL	A	226	-	5,5,5	0.22	0	5,5,5	0.66	0
3	GOL	C	226	-	5,5,5	0.34	0	5,5,5	0.41	0
3	GOL	D	227	-	5,5,5	0.37	0	5,5,5	0.48	0
3	GOL	D	228	-	5,5,5	0.43	0	5,5,5	0.26	0
3	GOL	B	227	-	5,5,5	0.34	0	5,5,5	0.16	0
4	CIT	A	227	-	3,12,12	2.50	1 (33%)	3,17,17	1.67	1 (33%)
3	GOL	C	227	-	5,5,5	0.37	0	5,5,5	0.47	0
3	GOL	D	226	-	5,5,5	0.19	0	5,5,5	0.45	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	GOL	B	226	-	-	2/4/4/4	-
3	GOL	A	226	-	-	0/4/4/4	-
3	GOL	C	226	-	-	0/4/4/4	-
3	GOL	D	227	-	-	2/4/4/4	-
3	GOL	D	228	-	-	2/4/4/4	-
3	GOL	B	227	-	-	0/4/4/4	-
4	CIT	A	227	-	-	4/6/16/16	-
3	GOL	C	227	-	-	0/4/4/4	-
3	GOL	D	226	-	-	2/4/4/4	-

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	A	227	CIT	C4-C3	-4.12	1.49	1.54

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	A	227	CIT	C3-C4-C5	-2.88	110.37	114.98

There are no chirality outliers.

All (12) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	D	227	GOL	O1-C1-C2-C3
3	D	228	GOL	O1-C1-C2-C3
4	A	227	CIT	C2-C3-C4-C5
4	A	227	CIT	O7-C3-C4-C5
4	A	227	CIT	C6-C3-C4-C5
3	D	226	GOL	O1-C1-C2-C3
3	D	226	GOL	O1-C1-C2-O2
3	B	226	GOL	O1-C1-C2-C3
3	D	227	GOL	O1-C1-C2-O2
3	D	228	GOL	O1-C1-C2-O2
4	A	227	CIT	C1-C2-C3-O7
3	B	226	GOL	O1-C1-C2-O2

There are no ring outliers.

3 monomers are involved in 3 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	D	227	GOL	1	0
3	D	228	GOL	1	0
4	A	227	CIT	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	221/243 (90%)	-0.13	0 100 100	12, 18, 31, 54	0
1	B	221/243 (90%)	-0.15	3 (1%) 75 82	10, 17, 37, 72	0
1	C	221/243 (90%)	-0.14	2 (0%) 84 89	11, 18, 30, 49	0
1	D	221/243 (90%)	-0.10	4 (1%) 68 76	9, 18, 38, 65	0
All	All	884/972 (90%)	-0.13	9 (1%) 82 87	9, 18, 33, 72	0

All (9) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	C	224	LEU	8.5
1	D	141	PHE	7.2
1	D	224	LEU	7.2
1	B	224	LEU	6.1
1	D	118	LEU	3.3
1	B	116	GLY	2.9
1	C	104[A]	SER	2.7
1	D	144	LYS	2.1
1	B	143	VAL	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
4	CIT	A	227	13/13	0.85	0.15	16,30,64,66	0
3	GOL	D	226	6/6	0.88	0.13	17,23,26,32	0
3	GOL	D	227	6/6	0.90	0.16	14,28,45,50	0
3	GOL	B	227	6/6	0.92	0.16	10,19,22,29	0
3	GOL	C	226	6/6	0.92	0.11	12,22,29,50	0
3	GOL	D	228	6/6	0.92	0.11	16,38,44,58	0
3	GOL	B	226	6/6	0.93	0.12	12,22,41,46	0
3	GOL	A	226	6/6	0.93	0.10	13,16,17,18	0
2	MG	C	225	1/1	0.94	0.12	16,16,16,16	0
3	GOL	C	227	6/6	0.94	0.12	7,24,43,48	0
2	MG	A	225	1/1	0.96	0.12	12,12,12,12	0
2	MG	B	225	1/1	0.98	0.12	13,13,13,13	0
5	CL	B	228	1/1	0.98	0.10	32,32,32,32	0
2	MG	D	225	1/1	0.99	0.14	13,13,13,13	0

6.5 Other polymers [i](#)

There are no such residues in this entry.