



Full wwPDB X-ray Structure Validation Report ⓘ

Oct 16, 2021 – 10:03 PM EDT

PDB ID : 1QVY
Title : Crystal structure of RhoGDI K(199,200)R double mutant
Authors : Czepas, J.; Devedjiev, Y.; Krowarsh, D.; Derewenda, U.; Derewenda, Z.S.
Deposited on : 2003-08-29
Resolution : 1.60 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix)	:	1.13
EDS	:	2.23.2
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac	:	5.8.0158
CCP4	:	7.0.044 (Gargrove)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.23.2


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.60 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	3398 (1.60-1.60)
Clashscore	141614	3665 (1.60-1.60)
Ramachandran outliers	138981	3564 (1.60-1.60)
Sidechain outliers	138945	3563 (1.60-1.60)
RSRZ outliers	127900	3321 (1.60-1.60)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	139	<div> <div style="width: 100%; height: 10px; background: linear-gradient(to right, red, orange, yellow, green, grey);"></div> <div style="display: flex; justify-content: space-between; margin-top: 5px;"> % 86% 12% .. </div> </div>
1	B	139	<div> <div style="width: 100%; height: 10px; background: linear-gradient(to right, red, orange, yellow, green, grey);"></div> <div style="display: flex; justify-content: space-between; margin-top: 5px;"> % 88% 12% . </div> </div>
1	C	139	<div> <div style="width: 100%; height: 10px; background: linear-gradient(to right, red, orange, yellow, green, grey);"></div> <div style="display: flex; justify-content: space-between; margin-top: 5px;"> % 88% 9% ... </div> </div>
1	D	139	<div> <div style="width: 100%; height: 10px; background: linear-gradient(to right, red, orange, yellow, green, grey);"></div> <div style="display: flex; justify-content: space-between; margin-top: 5px;"> % 88% 9% .. </div> </div>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard

residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	SO4	A	1006	-	-	X	X

2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 5291 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Rho GDP-dissociation inhibitor 1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	138	Total	C	N	O	S	0	0	0
			1118	713	189	212	4			
1	B	138	Total	C	N	O	S	0	0	0
			1118	713	189	212	4			
1	C	138	Total	C	N	O	S	0	0	0
			1118	713	189	212	4			
1	D	138	Total	C	N	O	S	0	0	0
			1118	713	189	212	4			

There are 12 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	66	MET	-	cloning artifact	UNP P52565
A	199	ARG	LYS	engineered mutation	UNP P52565
A	200	ARG	LYS	engineered mutation	UNP P52565
B	66	MET	-	cloning artifact	UNP P52565
B	199	ARG	LYS	engineered mutation	UNP P52565
B	200	ARG	LYS	engineered mutation	UNP P52565
C	66	MET	-	cloning artifact	UNP P52565
C	199	ARG	LYS	engineered mutation	UNP P52565
C	200	ARG	LYS	engineered mutation	UNP P52565
D	66	MET	-	cloning artifact	UNP P52565
D	199	ARG	LYS	engineered mutation	UNP P52565
D	200	ARG	LYS	engineered mutation	UNP P52565

- Molecule 2 is SULFATE ION (three-letter code: SO4) (formula: O₄S).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	O	S	0	0
			5	4	1		
2	A	1	Total	O	S	0	0
			5	4	1		
2	A	1	Total	O	S	0	0
			5	4	1		
2	B	1	Total	O	S	0	0
			5	4	1		
2	B	1	Total	O	S	0	0
			5	4	1		
2	C	1	Total	O	S	0	0
			5	4	1		
2	D	1	Total	O	S	0	0
			5	4	1		
2	D	1	Total	O	S	0	0
			5	4	1		


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	172	Total	O	1	0
			172	172		
3	B	220	Total	O	2	0
			220	220		
3	C	182	Total	O	0	0
			182	182		
3	D	205	Total	O	0	0
			205	205		



3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Rho GDP-dissociation inhibitor 1




- Molecule 1: Rho GDP-dissociation inhibitor 1



- Molecule 1: Rho GDP-dissociation inhibitor 1



- Molecule 1: Rho GDP-dissociation inhibitor 1



4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, α , β , γ	148.66Å 58.24Å 75.05Å 90.00° 92.49° 90.00°	Depositor
Resolution (Å)	27.02 – 1.60 27.14 – 1.60	Depositor EDS
% Data completeness (in resolution range)	100.0 (27.02-1.60) 99.6 (27.14-1.60)	Depositor EDS
R_{merge}	0.05	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.30 (at 1.60Å)	Xtriage
Refinement program	REFMAC 5.0	Depositor
R, R_{free}	0.172 , 0.210 0.182 , 0.220	Depositor DCC
R_{free} test set	1676 reflections (1.99%)	wwPDB-VP
Wilson B-factor (Å ²)	16.4	Xtriage
Anisotropy	0.282	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.40 , 51.9	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.34$	Xtriage
Estimated twinning fraction	0.004 for -h,-k,l	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	5291	wwPDB-VP
Average B, all atoms (Å ²)	21.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 42.18 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 2.1170e-04. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

¹ Intensities estimated from amplitudes.

² Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: SO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.77	0/1143	0.98	5/1543 (0.3%)
1	B	0.72	0/1143	0.98	7/1543 (0.5%)
1	C	0.76	0/1143	1.00	5/1543 (0.3%)
1	D	0.74	0/1143	0.94	4/1543 (0.3%)
All	All	0.75	0/4572	0.98	21/6172 (0.3%)

There are no bond length outliers.

All (21) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	C	172	ARG	NE-CZ-NH2	-8.11	116.24	120.30
1	B	196	LEU	CA-CB-CG	7.99	133.66	115.30
1	B	201	ASP	CB-CG-OD2	7.79	125.31	118.30
1	A	172	ARG	NE-CZ-NH2	-7.62	116.49	120.30
1	A	120	ARG	NE-CZ-NH2	-7.24	116.68	120.30
1	D	204	ASP	CB-CG-OD1	-6.85	112.13	118.30
1	C	140	ASP	CB-CG-OD2	6.68	124.31	118.30
1	C	134	ARG	NE-CZ-NH2	-6.51	117.05	120.30
1	C	77	LEU	CA-CB-CG	6.50	130.24	115.30
1	C	201	ASP	CB-CG-OD2	6.29	123.96	118.30
1	A	134	ARG	NE-CZ-NH2	-6.25	117.17	120.30
1	D	204	ASP	CB-CG-OD2	5.72	123.45	118.30
1	D	185	ASP	CB-CG-OD2	5.53	123.28	118.30
1	B	185	ASP	CB-CG-OD2	5.53	123.28	118.30
1	D	120	ARG	NE-CZ-NH1	5.52	123.06	120.30
1	A	188	ASP	CB-CG-OD2	5.35	123.11	118.30
1	B	134	ARG	NE-CZ-NH2	-5.23	117.69	120.30
1	B	172	ARG	NE-CZ-NH2	-5.22	117.69	120.30
1	B	196	LEU	N-CA-CB	-5.13	100.14	110.40
1	B	120	ARG	NE-CZ-NH2	-5.08	117.76	120.30

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	201	ASP	CB-CG-OD2	5.02	122.82	118.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1118	0	1112	15	0
1	B	1118	0	1112	7	0
1	C	1118	0	1112	13	0
1	D	1118	0	1112	15	0
2	A	15	0	0	3	0
2	B	10	0	0	1	0
2	C	5	0	0	0	0
2	D	10	0	0	0	0
3	A	172	0	0	6	0
3	B	220	0	0	4	2
3	C	182	0	0	7	2
3	D	205	0	0	7	0
All	All	5291	0	4448	50	2

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

All (50) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:87:GLU:CD	3:C:1075:HOH:O	2.26	0.74
1:A:204:ASP:OD2	3:A:1148:HOH:O	2.07	0.73
1:A:203:LYS:O	1:A:204:ASP:HB2	1.92	0.68
1:B:67:VAL:N	3:B:1178:HOH:O	2.28	0.67
1:A:155:GLU:OE1	3:A:1119:HOH:O	2.14	0.65
1:C:204:ASP:HB3	3:C:1053:HOH:O	1.96	0.64
1:D:203:LYS:O	1:D:204:ASP:HB2	1.97	0.64

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:D:77:LEU:HD11	1:D:110:TYR:CD2	2.36	0.61
1:C:203:LYS:O	1:C:204:ASP:HB2	2.01	0.60
1:C:134:ARG:NE	3:C:1132:HOH:O	2.35	0.59
1:D:77:LEU:HD13	1:D:112:ILE:CD1	2.35	0.57
1:B:203:LYS:O	1:B:204:ASP:HB2	2.05	0.57
1:D:201:ASP:OD2	3:D:1166:HOH:O	2.17	0.57
1:C:77:LEU:HD12	1:C:112:ILE:CD1	2.35	0.56
1:C:204:ASP:OD2	3:C:1080:HOH:O	2.17	0.56
1:D:164:GLU:OE1	3:D:1203:HOH:O	2.17	0.56
1:D:77:LEU:CD2	1:D:196:LEU:HD11	2.37	0.55
1:B:204:ASP:HB3	3:B:1030:HOH:O	2.06	0.54
1:D:117:ARG:HD2	3:D:1142:HOH:O	2.07	0.54
1:A:204:ASP:HB3	3:A:1070:HOH:O	2.07	0.54
1:A:77:LEU:CD1	1:A:196:LEU:HD11	2.42	0.50
1:C:77:LEU:HD23	1:C:86:LEU:HD13	1.94	0.49
1:C:87:GLU:CG	3:C:1075:HOH:O	2.59	0.49
1:C:77:LEU:HD12	1:C:112:ILE:HD13	1.95	0.49
1:C:87:GLU:HB3	3:C:1075:HOH:O	2.14	0.48
1:A:200:ARG:HD2	3:A:1140:HOH:O	2.13	0.48
1:C:130:GLN:HG2	1:C:177:ILE:CG2	2.45	0.47
1:A:123:VAL:CG1	1:A:126:MET:SD	3.04	0.46
1:A:69:ASN:HB3	2:A:1006:SO4:O2	2.16	0.45
1:D:117:ARG:NH1	3:D:1181:HOH:O	2.49	0.45
1:D:77:LEU:HD22	1:D:196:LEU:HD11	1.98	0.45
1:D:204:ASP:HB3	3:D:1027:HOH:O	2.16	0.45
1:C:106:GLU:HB2	1:C:167:LYS:HA	1.99	0.44
1:B:204:ASP:OD2	3:B:1158:HOH:O	2.20	0.44
1:D:194:TRP:CZ3	1:D:196:LEU:HD13	2.53	0.44
1:A:69:ASN:CB	2:A:1006:SO4:O2	2.67	0.43
1:A:68:PRO:HA	2:A:1006:SO4:O4	2.19	0.42
1:D:194:TRP:HZ3	1:D:196:LEU:HD13	1.85	0.42
1:A:123:VAL:HG12	1:A:126:MET:SD	2.60	0.42
1:B:131:HIS:CD2	1:B:141:LYS:HG2	2.55	0.42
1:A:169:MET:HE2	3:A:1129:HOH:O	2.20	0.41
1:A:184:ASP:HB2	3:A:1125:HOH:O	2.20	0.41
1:C:87:GLU:CB	3:C:1075:HOH:O	2.67	0.41
1:D:95:GLU:OE2	3:D:1209:HOH:O	2.22	0.41
1:A:123:VAL:HG11	1:A:126:MET:SD	2.61	0.41
1:B:99:LYS:HD3	3:B:1227:HOH:O	2.20	0.41
1:D:178:LYS:NZ	3:D:1211:HOH:O	2.50	0.41
1:D:77:LEU:CD1	1:D:112:ILE:HD13	2.49	0.41

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:200:ARG:NH1	2:B:1002:SO4:O1	2.53	0.41
1:A:106:GLU:HB2	1:A:167:LYS:HA	2.02	0.41

All (2) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:B:1109:HOH:O	3:C:1148:HOH:O[2_557]	2.04	0.16
3:B:1126:HOH:O	3:C:1125:HOH:O[2_557]	2.16	0.04

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	136/139 (98%)	134 (98%)	2 (2%)	0	100	100
1	B	136/139 (98%)	133 (98%)	3 (2%)	0	100	100
1	C	136/139 (98%)	134 (98%)	2 (2%)	0	100	100
1	D	136/139 (98%)	133 (98%)	3 (2%)	0	100	100
All	All	544/556 (98%)	534 (98%)	10 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	124/125 (99%)	122 (98%)	2 (2%)	62	41
1	B	124/125 (99%)	121 (98%)	3 (2%)	49	24
1	C	124/125 (99%)	120 (97%)	4 (3%)	39	15
1	D	124/125 (99%)	120 (97%)	4 (3%)	39	15
All	All	496/500 (99%)	483 (97%)	13 (3%)	46	21

All (13) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	169	MET
1	A	196	LEU
1	B	104	LEU
1	B	138	LYS
1	B	170	LEU
1	C	130	GLN
1	C	134	ARG
1	C	159	LEU
1	C	196	LEU
1	D	104	LEU
1	D	117	ARG
1	D	134	ARG
1	D	196	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
1	C	195	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

8 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	SO4	A	1001	-	4,4,4	0.28	0	6,6,6	0.29	0
2	SO4	A	1006	-	4,4,4	0.22	0	6,6,6	0.32	0
2	SO4	C	1003	-	4,4,4	0.30	0	6,6,6	0.45	0
2	SO4	D	1008	-	4,4,4	0.22	0	6,6,6	0.52	0
2	SO4	D	1004	-	4,4,4	0.47	0	6,6,6	0.47	0
2	SO4	B	1007	-	4,4,4	0.19	0	6,6,6	0.27	0
2	SO4	A	1005	-	4,4,4	0.28	0	6,6,6	0.49	0
2	SO4	B	1002	-	4,4,4	0.32	0	6,6,6	0.39	0

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

2 monomers are involved in 4 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	1006	SO4	3	0
2	B	1002	SO4	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	138/139 (99%)	-0.04	1 (0%) 87 87	12, 18, 29, 36	0
1	B	138/139 (99%)	-0.18	1 (0%) 87 87	13, 17, 28, 33	0
1	C	138/139 (99%)	-0.10	1 (0%) 87 87	13, 19, 28, 34	0
1	D	138/139 (99%)	-0.14	1 (0%) 87 87	12, 18, 29, 34	0
All	All	552/556 (99%)	-0.11	4 (0%) 87 87	12, 18, 29, 36	0

All (4) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	D	67	VAL	4.4
1	A	67	VAL	3.6
1	C	184	ASP	2.0
1	B	93	ASP	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
2	SO4	A	1006	5/5	0.78	0.58	64,64,65,65	0
2	SO4	A	1005	5/5	0.86	0.40	55,56,56,56	0
2	SO4	D	1008	5/5	0.95	0.24	37,37,38,39	0
2	SO4	B	1002	5/5	0.97	0.18	31,33,35,35	0
2	SO4	D	1004	5/5	0.97	0.15	27,28,30,32	0
2	SO4	A	1001	5/5	0.97	0.15	31,33,34,35	0
2	SO4	B	1007	5/5	0.98	0.16	42,43,44,45	0
2	SO4	C	1003	5/5	0.98	0.17	31,33,34,36	0

6.5 Other polymers [i](#)

There are no such residues in this entry.