



Full wwPDB X-ray Structure Validation Report ⓘ

May 29, 2020 – 03:54 pm BST

PDB ID : 2QVB
Title : Crystal Structure of Haloalkane Dehalogenase Rv2579 from Mycobacterium tuberculosis
Authors : Mazumdar, P.A.; Hulecki, J.; Cherney, M.M.; Garen, C.R.; James, M.N.G.; TB Structural Genomics Consortium (TBSGC)
Deposited on : 2007-08-08
Resolution : 1.19 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix)	:	1.13
EDS	:	2.11
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac	:	5.8.0158
CCP4	:	7.0.044 (Gargrove)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.11


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.19 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	1223 (1.22-1.18)
Clashscore	141614	1286 (1.22-1.18)
Ramachandran outliers	138981	1240 (1.22-1.18)
Sidechain outliers	138945	1239 (1.22-1.18)
RSRZ outliers	127900	1200 (1.22-1.18)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	297	<div> <div>3%</div> <div> <div></div> <div>86%</div> <div>14%</div> </div> <div></div> </div>
1	B	297	<div> <div>3%</div> <div> <div></div> <div>85%</div> <div>13%</div> </div> <div></div> </div>

2 Entry composition [i](#)

There are 4 unique types of molecules in this entry. The entry contains 5421 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Haloalkane dehalogenase 3.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	297	Total	C	N	O	S	3	10	1
			2365	1512	417	425	11			
1	B	297	Total	C	N	O	S	5	8	0
			2369	1513	418	427	11			

- Molecule 2 is CHLORIDE ION (three-letter code: CL) (formula: Cl).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	B	1	Total	Cl	0	0
			1	1		
2	A	1	Total	Cl	0	0
			1	1		

- Molecule 3 is 1,2-ETHANEDIOL (three-letter code: EDO) (formula: C₂H₆O₂).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total 8	C 4	O 4	0	1
3	B	1	Total 8	C 4	O 4	0	1


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	338	Total 338	O 338	0	0
4	B	331	Total 331	O 331	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Haloalkane dehalogenase 3



• Molecule 1: Haloalkane dehalogenase 3



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	60.70 Å 65.70 Å 129.70 Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	36.11 – 1.19 36.12 – 1.19	Depositor EDS
% Data completeness (in resolution range)	(Not available) (36.11-1.19) 81.0 (36.12-1.19)	Depositor EDS
R_{merge}	0.05	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.03 (at 1.19 Å)	Xtriage
Refinement program	REFMAC	Depositor
R, R_{free}	0.120 , 0.160 0.121 , 0.159	Depositor DCC
R_{free} test set	6695 reflections (4.99%)	wwPDB-VP
Wilson B-factor (Å ²)	8.3	Xtriage
Anisotropy	0.828	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.43 , 52.0	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.97	EDS
Total number of atoms	5421	wwPDB-VP
Average B, all atoms (Å ²)	13.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 58.48 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 2.0329e-05. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

¹ Intensities estimated from amplitudes.

² Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: EDO, CL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	1.28	14/2474 (0.6%)	1.22	18/3362 (0.5%)
1	B	1.30	19/2468 (0.8%)	1.28	19/3354 (0.6%)
All	All	1.29	33/4942 (0.7%)	1.25	37/6716 (0.6%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1

All (33) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	220	GLU	CD-OE2	12.31	1.39	1.25
1	B	253	ARG	NE-CZ	9.43	1.45	1.33
1	B	220	GLU	CD-OE2	8.14	1.34	1.25
1	A	87	ARG	CZ-NH1	7.29	1.42	1.33
1	A	79	ASP	CB-CG	7.21	1.66	1.51
1	A	151	VAL	CB-CG1	-7.12	1.38	1.52
1	A	233	GLU	CG-CD	6.88	1.62	1.51
1	B	253	ARG	CZ-NH2	6.75	1.41	1.33
1	A	79	ASP	CG-OD1	6.41	1.40	1.25
1	B	151	VAL	CB-CG1	-6.40	1.39	1.52
1	B	54[A]	GLU	CD-OE2	6.39	1.32	1.25
1	B	54[B]	GLU	CD-OE2	6.39	1.32	1.25
1	B	295	ARG	CZ-NH2	6.37	1.41	1.33
1	B	79	ASP	CB-CG	6.36	1.65	1.51
1	A	266	GLU	CD-OE1	6.33	1.32	1.25
1	B	144	TRP	CB-CG	-5.99	1.39	1.50

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	217	GLU	CB-CG	-5.99	1.40	1.52
1	B	87	ARG	CZ-NH1	5.74	1.40	1.33
1	A	217	GLU	CB-CG	-5.71	1.41	1.52
1	B	193	ARG	CZ-NH2	-5.57	1.25	1.33
1	A	11	GLN	C-O	-5.53	1.12	1.23
1	A	29	LYS	CD-CE	-5.52	1.37	1.51
1	B	237[A]	MET	CG-SD	-5.46	1.67	1.81
1	B	237[B]	MET	CG-SD	-5.46	1.67	1.81
1	B	157	SER	CB-OG	5.40	1.49	1.42
1	B	29	LYS	CD-CE	5.38	1.64	1.51
1	B	257	TYR	CB-CG	-5.35	1.43	1.51
1	A	87	ARG	CD-NE	-5.32	1.37	1.46
1	B	192	ARG	NE-CZ	-5.31	1.26	1.33
1	A	157	SER	CB-OG	5.30	1.49	1.42
1	A	49	ILE	CB-CG1	5.23	1.68	1.54
1	B	234	GLU	CD-OE2	-5.18	1.20	1.25
1	A	166	GLU	CD-OE1	5.06	1.31	1.25

All (37) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	79	ASP	CB-CG-OD1	11.93	129.04	118.30
1	B	87	ARG	NE-CZ-NH2	-11.51	114.55	120.30
1	B	237[A]	MET	CG-SD-CE	-11.37	82.01	100.20
1	B	237[B]	MET	CG-SD-CE	-11.37	82.01	100.20
1	A	79	ASP	CB-CG-OD1	10.89	128.10	118.30
1	A	295	ARG	NE-CZ-NH1	-10.02	115.29	120.30
1	B	193	ARG	NE-CZ-NH2	-9.77	115.42	120.30
1	A	87	ARG	NE-CZ-NH2	-8.91	115.85	120.30
1	A	87	ARG	NE-CZ-NH1	8.86	124.73	120.30
1	B	253	ARG	NE-CZ-NH1	8.33	124.47	120.30
1	A	156	ARG	NE-CZ-NH2	-8.24	116.18	120.30
1	B	253	ARG	NE-CZ-NH2	7.89	124.25	120.30
1	B	293	ARG	NE-CZ-NH2	-7.67	116.47	120.30
1	B	293	ARG	NE-CZ-NH1	7.54	124.07	120.30
1	B	253	ARG	NH1-CZ-NH2	-7.45	111.20	119.40
1	A	27	GLU	OE1-CD-OE2	-7.42	114.39	123.30
1	B	123	ARG	NE-CZ-NH2	-7.19	116.71	120.30
1	B	123	ARG	NE-CZ-NH1	6.89	123.75	120.30
1	B	109	ASP	CB-CG-OD2	6.73	124.36	118.30
1	A	58	ARG	NE-CZ-NH2	-6.45	117.08	120.30
1	A	253	ARG	NE-CZ-NH1	6.26	123.43	120.30

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	202	ARG	NE-CZ-NH2	-6.21	117.19	120.30
1	A	29	LYS	CB-CA-C	-6.10	98.20	110.40
1	B	47	ARG	NE-CZ-NH1	6.08	123.34	120.30
1	A	292	ARG	NE-CZ-NH1	-6.07	117.27	120.30
1	A	71	ASP	CB-CG-OD1	6.01	123.71	118.30
1	B	255	ARG	NE-CZ-NH1	5.84	123.22	120.30
1	A	91	PHE	CB-CG-CD2	-5.78	116.75	120.80
1	A	255	ARG	NE-CZ-NH1	5.78	123.19	120.30
1	B	87	ARG	NE-CZ-NH1	5.69	123.15	120.30
1	A	215	ASP	CB-CG-OD1	5.67	123.40	118.30
1	A	109	ASP	CB-CG-OD1	-5.55	113.31	118.30
1	B	203	ARG	NE-CZ-NH2	-5.49	117.56	120.30
1	B	217	GLU	OE1-CD-OE2	-5.32	116.92	123.30
1	A	151	VAL	CA-CB-CG1	5.29	118.83	110.90
1	A	193	ARG	NE-CZ-NH2	-5.19	117.70	120.30
1	B	259	ARG	NE-CZ-NH1	-5.16	117.72	120.30

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	28[B]	GLY	Mainchain

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2365	0	2264	19	0
1	B	2369	0	2265	21	0
2	A	1	0	0	0	0
2	B	1	0	0	0	0
3	A	8	0	12	0	0
3	B	8	0	12	0	0
4	A	338	0	0	5	0
4	B	331	0	0	3	0
All	All	5421	0	4553	40	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (40) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:120:ASN:HD21	1:B:123:ARG:HH11	1.24	0.84
1:A:243:ASN:HD21	1:A:255:ARG:HE	1.20	0.84
1:A:120:ASN:HD21	1:A:123:ARG:HH11	1.24	0.83
1:B:243:ASN:HD21	1:B:255:ARG:HE	1.29	0.80
1:B:37:HIS:HE1	1:B:64:LEU:H	1.35	0.73
1:B:253:ARG:NH1	4:B:600:HOH:O	2.24	0.69
1:A:37:HIS:HE1	1:A:64:LEU:H	1.40	0.69
1:A:237[B]:MET:SD	4:A:592:HOH:O	2.53	0.65
1:B:37:HIS:CE1	1:B:64:LEU:H	2.16	0.63
1:A:120:ASN:ND2	1:A:123:ARG:HH11	1.95	0.62
1:A:52:HIS:HE1	4:A:514:HOH:O	1.84	0.61
1:A:6[B]:VAL:HG22	1:A:187:GLU:HG3	1.83	0.60
1:B:120:ASN:ND2	1:B:123:ARG:HH11	1.99	0.59
1:A:37:HIS:CE1	1:A:64:LEU:H	2.21	0.58
1:B:39:ASN:HD21	1:B:205:THR:HA	1.69	0.57
1:B:127:GLN:HG2	1:B:294:LEU:HD13	1.87	0.56
1:A:243:ASN:HD21	1:A:255:ARG:NE	1.99	0.56
1:B:243:ASN:ND2	1:B:255:ARG:HE	1.99	0.56
1:A:37:HIS:HD2	1:A:38:GLY:O	1.89	0.55
1:B:37:HIS:HD2	1:B:38:GLY:O	1.91	0.54
1:A:243:ASN:ND2	1:A:255:ARG:HE	2.00	0.53
1:A:52:HIS:HD2	4:A:366:HOH:O	1.91	0.53
1:B:127:GLN:HG2	1:B:294:LEU:CD1	2.41	0.49
1:B:243:ASN:HD21	1:B:255:ARG:NE	2.02	0.49
1:A:108:HIS:HD2	4:A:316:HOH:O	1.97	0.47
1:A:6[B]:VAL:HG22	1:A:187:GLU:CG	2.45	0.47
1:B:126:VAL:HG12	1:B:237[A]:MET:HE3	1.98	0.46
1:B:293:ARG:HG3	4:B:469:HOH:O	2.14	0.46
1:B:108:HIS:HD2	4:B:313:HOH:O	1.99	0.46
1:B:237[A]:MET:HG3	1:B:238:PRO:N	2.31	0.46
1:A:20:LYS:HE3	4:A:372:HOH:O	2.15	0.46
1:B:126:VAL:HG12	1:B:237[A]:MET:CE	2.45	0.46
1:B:76:SER:HB3	1:B:200:GLU:HB3	2.00	0.43
1:A:83:TYR:H	1:A:211:ASN:ND2	2.16	0.43
1:A:108:HIS:HE1	1:A:273:HIS:O	2.03	0.42
1:A:6[B]:VAL:HG11	1:A:186:GLU:CG	2.50	0.42
1:B:83:TYR:H	1:B:211:ASN:ND2	2.18	0.41

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:237[A]:MET:HB3	1:A:237[A]:MET:HE2	1.79	0.41
1:B:132:MET:HB3	1:B:242:ILE:HB	2.02	0.41
1:B:108:HIS:HE1	1:B:273:HIS:O	2.04	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	304/297 (102%)	296 (97%)	8 (3%)	0	100	100
1	B	303/297 (102%)	294 (97%)	9 (3%)	0	100	100
All	All	607/594 (102%)	590 (97%)	17 (3%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	245/244 (100%)	243 (99%)	2 (1%)	81	55
1	B	242/244 (99%)	235 (97%)	7 (3%)	42	7
All	All	487/488 (100%)	478 (98%)	9 (2%)	60	21

All (9) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	170	PHE
1	A	253	ARG
1	B	40	PRO
1	B	170	PHE
1	B	224	LEU
1	B	228	TYR
1	B	237[A]	MET
1	B	237[B]	MET
1	B	238	PRO

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (17) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	37	HIS
1	A	52	HIS
1	A	108	HIS
1	A	120	ASN
1	A	197	ASN
1	A	211	ASN
1	A	243	ASN
1	A	289	GLN
1	B	36	GLN
1	B	37	HIS
1	B	39	ASN
1	B	108	HIS
1	B	120	ASN
1	B	197	ASN
1	B	211	ASN
1	B	243	ASN
1	B	289	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates

There are no carbohydrates in this entry.

5.6 Ligand geometry

Of 6 ligands modelled in this entry, 2 are monoatomic - leaving 4 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	EDO	A	1[B]	-	3,3,3	0.87	0	2,2,2	0.30	0
3	EDO	B	1[B]	-	3,3,3	0.62	0	2,2,2	0.39	0
3	EDO	B	1[A]	-	3,3,3	0.48	0	2,2,2	0.80	0
3	EDO	A	1[A]	-	3,3,3	0.66	0	2,2,2	0.55	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	EDO	A	1[B]	-	-	0/1/1/1	-
3	EDO	B	1[B]	-	-	1/1/1/1	-
3	EDO	B	1[A]	-	-	0/1/1/1	-
3	EDO	A	1[A]	-	-	0/1/1/1	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (1) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	B	1[B]	EDO	O1-C1-C2-O2

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	297/297 (100%)	0.15	8 (2%) 54 53	7, 10, 18, 31	1 (0%)
1	B	297/297 (100%)	0.21	8 (2%) 54 53	7, 11, 18, 34	1 (0%)
All	All	594/594 (100%)	0.18	16 (2%) 54 53	7, 11, 18, 34	2 (0%)

All (16) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	298	ALA	10.1
1	A	3	ALA	7.4
1	A	298	ALA	6.7
1	B	299	GLY	6.4
1	A	297	ALA	5.5
1	A	299	GLY	5.4
1	B	297	ALA	5.4
1	B	3	ALA	5.2
1	A	4	PHE	5.1
1	B	296	SER	5.0
1	B	294	LEU	4.2
1	A	296	SER	3.7
1	A	295	ARG	3.1
1	A	10	GLY	2.8
1	B	124	ASP	2.2
1	B	295	ARG	2.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
3	EDO	A	1[A]	4/4	0.86	0.20	26,26,26,28	4
3	EDO	A	1[B]	4/4	0.86	0.20	26,28,30,31	4
3	EDO	B	1[B]	4/4	0.91	0.12	25,27,28,30	4
3	EDO	B	1[A]	4/4	0.91	0.12	23,26,27,28	4
2	CL	A	2	1/1	1.00	0.05	11,11,11,11	1
2	CL	B	2	1/1	1.00	0.03	11,11,11,11	1

6.5 Other polymers [i](#)

There are no such residues in this entry.