



Full wwPDB X-ray Structure Validation Report ⓘ

May 13, 2020 – 08:18 pm BST

PDB ID : 3QVM
Title : The structure of olei00960, a hydrolase from Oleispira antarctica
Authors : Singer, A.U.; Kagan, O.; Kim, Y.; Edwards, A.M.; Joachimiak, A.; Savchenko, A.; Midwest Center for Structural Genomics (MCSG)
Deposited on : 2011-02-25
Resolution : 2.00 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.11
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.11


1 Overall quality at a glance ⓘ

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	8085 (2.00-2.00)
Clashscore	141614	9178 (2.00-2.00)
Ramachandran outliers	138981	9054 (2.00-2.00)
Sidechain outliers	138945	9053 (2.00-2.00)
RSRZ outliers	127900	7900 (2.00-2.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	282	<div> <div>2%</div> <div> <div></div> <div>82%</div> <div>15%</div> <div>••</div> </div> </div>
1	B	282	<div> <div>2%</div> <div> <div></div> <div>86%</div> <div>12%</div> <div>•</div> </div> </div>

2 Entry composition [i](#)

There are 6 unique types of molecules in this entry. The entry contains 4584 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Olei00960.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	273	Total	C	N	O	S	Se	7	2	0
			2116	1349	340	413	6	8			
1	B	280	Total	C	N	O	S	Se	8	4	0
			2197	1397	354	431	6	9			

- Molecule 2 is SULFATE ION (three-letter code: SO4) (formula: O₄S).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	O	S	0	0
			5	4	1		
2	A	1	Total	O	S	0	0
			5	4	1		
2	A	1	Total	O	S	0	0
			5	4	1		
2	B	1	Total	O	S	0	0
			5	4	1		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	B	1	Total	O	S	0	0
			5	4	1		

- Molecule 3 is SODIUM ION (three-letter code: NA) (formula: Na).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	B	1	Total	Na	0	0
			1	1		
3	A	1	Total	Na	0	0
			1	1		

- Molecule 4 is CHLORIDE ION (three-letter code: CL) (formula: Cl).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	B	1	Total	Cl	0	0
			1	1		

- Molecule 5 is CALCIUM ION (three-letter code: CA) (formula: Ca).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	B	1	Total	Ca	0	0
			1	1		


- Molecule 6 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	99	Total	O	0	2
			101	101		
6	B	141	Total	O	0	0
			141	141		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Olei00960



• Molecule 1: Olei00960



4 Data and refinement statistics

Property	Value
Space group	P 4
Cell constants a, b, c, α , β , γ	123.07Å 123.07Å 49.22Å 90.00° 90.00° 90.00°
Resolution (Å)	30.77 – 2.00 30.77 – 2.00
% Data completeness (in resolution range)	99.1 (30.77-2.00) 99.6 (30.77-2.00)
R_{merge}	0.12
R_{sym}	(Not available)
$\langle I/\sigma(I) \rangle$ ¹	4.63 (at 2.00Å)
Refinement program	PHENIX 1.6_289, REFMAC 5.2.0019 plus PHENIX (phenix.refine: 1.6
R, R_{free}	0.156 , 0.185 0.150 , 0.180
R_{free} test set	2002 reflections (3.98%)
Wilson B-factor (Å ²)	19.4
Anisotropy	0.530
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.35 , 23.7
L-test for twinning ²	$\langle L \rangle = 0.39$, $\langle L^2 \rangle = 0.22$
Estimated twinning fraction	0.338 for h,-k,-l
F_o, F_c correlation	0.95
Total number of atoms	4584
Average B, all atoms (Å ²)	19.0

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.56% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: NA, CA, SO4, CL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.34	0/2152	0.47	0/2909
1	B	0.36	0/2232	0.52	0/3011
All	All	0.35	0/4384	0.50	0/5920

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2116	0	2064	30	0
1	B	2197	0	2154	28	0
2	A	15	0	0	0	0
2	B	10	0	0	0	0
3	A	1	0	0	0	0
3	B	1	0	0	0	0
4	B	1	0	0	0	0
5	B	1	0	0	0	0
6	A	101	0	0	1	0
6	B	141	0	0	1	0
All	All	4584	0	4218	55	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

All (55) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:28:LYS:HB2	1:B:279:THR:HB	1.68	0.76
1:A:154:ILE:HG13	1:A:206:PHE:HE1	1.54	0.72
1:A:46:MSE:HE3	1:A:259:HIS:HA	1.72	0.71
1:A:208:ASP:OD1	1:A:210:ARG:HD3	1.90	0.70
1:A:46:MSE:CE	1:A:266:ILE:HD11	2.22	0.69
1:A:46:MSE:HE1	1:A:266:ILE:HD11	1.76	0.66
1:B:278:GLN:OE1	1:B:278:GLN:HA	1.96	0.65
1:A:154:ILE:HG13	1:A:206:PHE:CE1	2.30	0.65
1:B:193:ASP:O	1:B:196:VAL:HG22	2.02	0.59
1:A:154:ILE:CG1	1:A:206:PHE:HE1	2.15	0.59
1:B:28:LYS:CB	1:B:279:THR:HB	2.33	0.58
1:B:154:ILE:HG23	1:B:198:LYS:HE2	1.87	0.56
1:B:154:ILE:HG13	1:B:206:PHE:CZ	2.42	0.55
1:B:28:LYS:HB2	1:B:279:THR:CB	2.37	0.54
1:B:225:SER:HB3	1:B:228:ASP:HB2	1.89	0.54
1:B:278:GLN:O	1:B:279:THR:OG1	2.20	0.53
1:B:74:THR:O	1:B:78:SER:HB3	2.10	0.52
1:B:267:THR:HB	1:B:268:PRO:HD3	1.92	0.52
1:A:237:GLN:O	1:A:241:GLU:HG3	2.10	0.52
1:A:30[A]:VAL:HG12	1:A:100:SER:HB2	1.93	0.51
1:B:213:LEU:HD21	1:B:239:MSE:HE3	1.92	0.51
1:A:194:PRO:O	1:A:198:LYS:HG3	2.11	0.51
1:A:63:GLY:HA2	1:A:69:LEU:HD12	1.93	0.51
1:A:75:LYS:NZ	1:B:277:ASN:HD22	2.10	0.51
1:A:83:TYR:O	1:A:86:ASP:HB2	2.11	0.50
1:A:75:LYS:NZ	1:B:277:ASN:ND2	2.59	0.49
1:B:28:LYS:HD3	1:B:279:THR:OG1	2.12	0.49
1:A:27:GLU:HG3	6:A:383:HOH:O	2.12	0.49
1:B:109:ILE:HG12	6:B:310:HOH:O	2.14	0.48
1:B:28:LYS:HB2	1:B:279:THR:CG2	2.44	0.47
1:A:216:ILE:O	1:A:244:PRO:HD2	2.14	0.47
1:B:202:LYS:O	1:B:206:PHE:HB2	2.14	0.46
1:A:46:MSE:HE3	1:A:259:HIS:CA	2.41	0.46
1:A:63:GLY:HA2	1:A:69:LEU:CD1	2.47	0.45
1:B:85[B]:LYS:HD3	1:B:85[B]:LYS:HA	1.74	0.44
1:B:28:LYS:CG	1:B:279:THR:HB	2.47	0.44
1:A:46:MSE:HE3	1:A:259:HIS:CB	2.48	0.44

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:278:GLN:CA	1:B:278:GLN:OE1	2.65	0.44
1:A:124:ASP:HA	1:A:218:THR:HG22	2.00	0.44
1:B:277:ASN:O	1:B:278:GLN:HB2	2.17	0.43
1:A:32:LEU:CD2	1:A:102:ILE:HD12	2.49	0.43
1:B:0:GLY:HA3	1:B:5:ASP:OD2	2.18	0.43
1:A:225:SER:HB3	1:A:228:ASP:HB2	2.01	0.43
1:B:76:ARG:O	1:B:82:GLY:HA3	2.19	0.42
1:B:227:LYS:HB3	1:B:254:GLU:HG2	2.00	0.42
1:A:202:LYS:O	1:A:206:PHE:HB2	2.19	0.42
1:A:81:GLU:HG2	1:A:81:GLU:H	1.61	0.42
1:B:108:SER:CB	1:B:130:PRO:HB3	2.50	0.42
1:A:134:PHE:O	1:A:145:PHE:HB2	2.20	0.41
1:A:68:ASP:OD1	1:A:70:GLU:HB3	2.20	0.41
1:A:132:PRO:HD3	1:A:239:MSE:SE	2.70	0.41
1:A:267:THR:HB	1:A:268:PRO:HD3	2.03	0.41
1:A:47:LEU:HB2	1:A:48:PRO:HD3	2.03	0.41
1:A:75:LYS:HZ2	1:B:277:ASN:ND2	2.18	0.41
1:B:14:VAL:HG12	1:B:20:ILE:HD11	2.02	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	273/282 (97%)	264 (97%)	9 (3%)	0	100	100
1	B	282/282 (100%)	274 (97%)	7 (2%)	1 (0%)	34	30
All	All	555/564 (98%)	538 (97%)	16 (3%)	1 (0%)	47	44

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	278	GLN

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	236/234 (101%)	230 (98%)	6 (2%)	47	49
1	B	246/234 (105%)	237 (96%)	9 (4%)	34	32
All	All	482/468 (103%)	467 (97%)	15 (3%)	39	40

All (15) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	71	SER
1	A	120	ASP
1	A	124	ASP
1	A	200	PHE
1	A	237	GLN
1	A	245	ASN
1	B	1	MSE
1	B	4	GLN
1	B	27	GLU
1	B	107	SER
1	B	124	ASP
1	B	158	ASP
1	B	200	PHE
1	B	278	GLN
1	B	279	THR

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
1	B	277	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

Of 9 ligands modelled in this entry, 4 are monoatomic - leaving 5 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	SO4	B	282	-	4,4,4	0.14	0	6,6,6	0.24	0
2	SO4	A	282	-	4,4,4	0.18	0	6,6,6	0.17	0
2	SO4	B	283	-	4,4,4	0.25	0	6,6,6	0.29	0
2	SO4	A	283	-	4,4,4	0.21	0	6,6,6	0.17	0
2	SO4	A	284	-	4,4,4	0.22	0	6,6,6	0.26	0

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	265/282 (93%)	0.26	5 (1%) 66 65	14, 21, 32, 40	0
1	B	271/282 (96%)	0.12	5 (1%) 68 66	11, 16, 25, 34	0
All	All	536/564 (95%)	0.19	10 (1%) 66 65	11, 18, 29, 40	0

All (10) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	279	THR	9.2
1	A	8	CYS	7.7
1	A	9	TYR	6.5
1	A	7	ILE	4.1
1	B	148	ASP	3.2
1	B	278	GLN	3.2
1	A	6	VAL	3.1
1	B	277	ASN	2.2
1	B	0	GLY	2.2
1	A	5	ASP	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
4	CL	B	285	1/1	0.88	0.10	40,40,40,40	0
3	NA	A	285	1/1	0.90	0.09	24,24,24,24	0
2	SO4	A	284	5/5	0.90	0.31	35,38,42,48	0
2	SO4	A	282	5/5	0.92	0.20	37,39,41,44	0
2	SO4	B	282	5/5	0.93	0.26	29,29,34,34	0
2	SO4	B	283	5/5	0.93	0.30	21,25,30,31	0
3	NA	B	284	1/1	0.95	0.10	16,16,16,16	0
2	SO4	A	283	5/5	0.95	0.24	28,29,32,34	0
5	CA	B	286	1/1	0.97	0.22	32,32,32,32	0

6.5 Other polymers [i](#)

There are no such residues in this entry.