



# Full wwPDB X-ray Structure Validation Report ⓘ

May 29, 2020 – 10:11 pm BST

PDB ID : 3QZ7  
Title : T-3 ternary complex of Dpo4  
Authors : Pata, J.D.; Wu, Y.; Wilson, R.C.  
Deposited on : 2011-03-04  
Resolution : 2.00 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.11  
buster-report : 1.1.7 (2018)  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.11


# 1 Overall quality at a glance ⓘ

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	8085 (2.00-2.00)
Clashscore	141614	9178 (2.00-2.00)
Ramachandran outliers	138981	9054 (2.00-2.00)
Sidechain outliers	138945	9053 (2.00-2.00)
RSRZ outliers	127900	7900 (2.00-2.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	360	<div> <div>23%</div> <div> <div></div> <div>82%</div> <div>12%</div> <div>• 5%</div> </div> </div>
2	P	13	<div> <div>77%</div> <div>15%</div> <div>8%</div> </div>
3	T	19	<div> <div>53%</div> <div>21%</div> <div>5%</div> <div>21%</div> </div>

## 2 Entry composition

There are 6 unique types of molecules in this entry. The entry contains 6635 atoms, of which 3219 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called DNA polymerase IV.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	341	Total	C	H	N	O	S	0	0	0
			5634	1760	2891	472	504	7			

There are 8 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	353	GLY	-	EXPRESSION TAG	UNP Q97W02
A	354	GLY	-	EXPRESSION TAG	UNP Q97W02
A	355	HIS	-	EXPRESSION TAG	UNP Q97W02
A	356	HIS	-	EXPRESSION TAG	UNP Q97W02
A	357	HIS	-	EXPRESSION TAG	UNP Q97W02
A	358	HIS	-	EXPRESSION TAG	UNP Q97W02
A	359	HIS	-	EXPRESSION TAG	UNP Q97W02
A	360	HIS	-	EXPRESSION TAG	UNP Q97W02


- Molecule 2 is a DNA chain called 5'-D(\*GP\*GP\*CP\*AP\*CP\*TP\*GP\*AP\*TP\*CP\*GP\*GP\*G)-3'.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
2	P	13	Total	C	H	N	O	P	0	0	0
			415	127	147	53	76	12			

- Molecule 3 is a DNA chain called 5'-D(\*TP\*TP\*AP\*CP\*GP\*CP\*CP\*TP\*CP\*GP\*AP\*TP\*CP\*AP\*GP\*TP\*GP\*CP\*C)-3'.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
3	T	15	Total	C	H	N	O	P	0	0	0
			473	144	169	54	91	15			

- Molecule 4 is 2'-DEOXYCYTIDINE-5'-TRIPHOSPHATE (three-letter code: DCP) (formula: C<sub>9</sub>H<sub>16</sub>N<sub>3</sub>O<sub>13</sub>P<sub>3</sub>).



Mol	Chain	Residues	Atoms						ZeroOcc	AltConf
4	A	1	Total	C	H	N	O	P	0	0
			40	9	12	3	13	3		

- Molecule 5 is CALCIUM ION (three-letter code: CA) (formula: Ca).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	A	3	Total Ca 3 3	0	0

- Molecule 6 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
6	A	56	Total O 56 56	0	0
6	P	6	Total O 6 6	0	0
6	T	8	Total O 8 8	0	0



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 2	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	98.71Å 102.46Å 52.74Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	25.54 – 2.00 25.54 – 2.00	Depositor EDS
% Data completeness (in resolution range)	90.0 (25.54-2.00) 90.0 (25.54-2.00)	Depositor EDS
$R_{merge}$	0.05	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.71 (at 1.99Å)	Xtriage
Refinement program	PHENIX	Depositor
R, $R_{free}$	0.227 , 0.267 0.219 , 0.262	Depositor DCC
$R_{free}$ test set	1629 reflections (4.89%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	48.9	Xtriage
Anisotropy	0.152	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.45 , 53.9	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.47$ , $\langle L^2 \rangle = 0.30$	Xtriage
Estimated twinning fraction	0.014 for k,h,-l	Xtriage
$F_o, F_c$ correlation	0.95	EDS
Total number of atoms	6635	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	67.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.92% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality ⓘ

### 5.1 Standard geometry ⓘ

Bond lengths and bond angles in the following residue types are not validated in this section: CA, DCP

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z  > 5$	RMSZ	# $ Z  > 5$
1	A	0.37	0/2782	0.52	0/3736
2	P	0.69	0/301	1.43	3/464 (0.6%)
3	T	0.70	0/339	1.47	6/520 (1.2%)
All	All	0.45	0/3422	0.81	9/4720 (0.2%)

There are no bond length outliers.

All (9) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	T	16	DT	O4'-C4'-C3'	-8.59	100.84	106.00
3	T	16	DT	O4'-C1'-N1	7.09	112.96	108.00
3	T	9	DC	O4'-C1'-N1	7.08	112.96	108.00
2	P	6	DT	O4'-C1'-N1	-6.38	103.53	108.00
3	T	5	DG	O4'-C1'-N9	-6.04	103.78	108.00
2	P	12	DG	O4'-C1'-N9	5.61	111.93	108.00
2	P	12	DG	C1'-O4'-C4'	-5.23	104.87	110.10
3	T	10	DG	C5-C6-O6	-5.18	125.49	128.60
3	T	10	DG	N1-C6-O6	5.00	122.90	119.90

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2743	2891	2889	29	0
2	P	268	147	147	1	0
3	T	304	169	169	2	0
4	A	28	12	12	1	0
5	A	3	0	0	0	0
6	A	56	0	0	0	0
6	P	6	0	0	0	0
6	T	8	0	0	0	0
All	All	3416	3219	3217	32	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (32) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:1:MET:HB2	1:A:112:SER:OG	1.82	0.79
3:T:16:DT:H2"	3:T:17:DG:O5'	2.03	0.59
1:A:247:ARG:C	1:A:248:ILE:HD12	2.23	0.58
3:T:16:DT:H2'	3:T:17:DG:C8	2.40	0.57
1:A:122:TYR:CE2	1:A:126:LEU:HD11	2.39	0.57
1:A:133:LEU:O	1:A:137:LYS:HD3	2.05	0.56
1:A:186:ILE:HD11	1:A:225:LEU:HD21	1.86	0.56
1:A:108:TYR:C	1:A:109:LEU:HD12	2.26	0.56
1:A:289:VAL:HB	1:A:332:ARG:HB2	1.89	0.53
1:A:185:GLY:O	1:A:221:LYS:CE	2.59	0.50
1:A:251:MET:HG2	1:A:264:TYR:CD2	2.47	0.49
1:A:100:GLU:HB2	1:A:237:ILE:HG23	1.93	0.49
1:A:10:TYR:HA	4:A:361:DCP:PB	2.54	0.48
2:P:6:DT:H2"	2:P:7:DG:C8	2.50	0.47
1:A:260:GLU:O	1:A:263:PRO:HD2	2.15	0.47
1:A:67:ILE:HG22	1:A:68:LEU:HG	1.96	0.46
1:A:9:ASP:O	1:A:10:TYR:C	2.54	0.46
1:A:122:TYR:HE2	1:A:126:LEU:HD11	1.80	0.45
1:A:241:VAL:HG23	1:A:241:VAL:O	2.17	0.45
1:A:248:ILE:HD12	1:A:248:ILE:N	2.32	0.44
1:A:180:ILE:HD13	1:A:199:ILE:HG22	1.99	0.44
1:A:31:CYS:HB3	1:A:61:ILE:HD11	1.99	0.44
1:A:79:GLU:CD	1:A:79:GLU:H	2.21	0.43
1:A:248:ILE:HA	1:A:334:GLY:HA3	2.01	0.42
1:A:109:LEU:HD12	1:A:109:LEU:N	2.35	0.42
1:A:160:PRO:O	1:A:161:ASN:C	2.58	0.42

*Continued on next page...*



*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:197:LEU:HD22	1:A:212:LYS:HZ3	1.84	0.42
1:A:129:LYS:HE3	1:A:129:LYS:HB3	1.89	0.42
1:A:185:GLY:O	1:A:221:LYS:HE2	2.19	0.41
1:A:300:ARG:HD3	1:A:302:PHE:CZ	2.56	0.40
1:A:31:CYS:HB3	1:A:61:ILE:CD1	2.51	0.40
1:A:180:ILE:HD13	1:A:194:LEU:HD13	2.03	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	339/360 (94%)	325 (96%)	13 (4%)	1 (0%)	41 37

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	10	TYR

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	300/315 (95%)	292 (97%)	8 (3%)	44 46

All (8) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	1	MET
1	A	10	TYR
1	A	127	GLU
1	A	180	ILE
1	A	234	ASN
1	A	252	LYS
1	A	292	ASP
1	A	294	ASP

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 5.6 Ligand geometry [i](#)

Of 4 ligands modelled in this entry, 3 are monoatomic - leaving 1 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
4	DCP	A	361	5	23,29,29	2.23	7 (30%)	30,45,45	2.16	6 (20%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	DCP	A	361	5	-	0/19/34/34	0/2/2/2

All (7) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	A	361	DCP	C6-N1	5.75	1.42	1.35
4	A	361	DCP	C4-N3	4.23	1.42	1.35
4	A	361	DCP	C6-C5	3.95	1.46	1.38
4	A	361	DCP	C4-N4	3.17	1.44	1.35
4	A	361	DCP	C2-N3	3.05	1.44	1.38
4	A	361	DCP	C3'-C4'	-2.71	1.45	1.53
4	A	361	DCP	C5-C4	2.48	1.47	1.41

All (6) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	A	361	DCP	O5'-PA-O1A	-7.32	80.47	109.07
4	A	361	DCP	O2A-PA-O5'	-4.12	88.63	107.75
4	A	361	DCP	PB-O3B-PG	-3.59	120.50	132.83
4	A	361	DCP	O5'-C5'-C4'	3.42	120.76	108.99
4	A	361	DCP	O3G-PG-O3B	3.37	115.93	104.64
4	A	361	DCP	C2-N3-C4	3.25	119.64	116.34

There are no chirality outliers.

There are no torsion outliers.


There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	A	361	DCP	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be

highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



## 5.7 Other polymers [i](#)

There are no such residues in this entry.









## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	341/360 (94%)	1.42	81 (23%)  	36, 59, 83, 91	0
2	P	13/13 (100%)	0.41	0  	54, 60, 72, 73	0
3	T	15/19 (78%)	0.17	0  	51, 65, 69, 77	0
All	All	369/392 (94%)	1.33	81 (21%)  	36, 60, 82, 91	0

All (81) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	116	ARG	6.8
1	A	335	VAL	6.5
1	A	37	PHE	5.8
1	A	334	GLY	5.6
1	A	287	VAL	5.1
1	A	248	ILE	5.0
1	A	333	ILE	4.7
1	A	48	TYR	4.7
1	A	209	GLU	4.7
1	A	5	PHE	4.7
1	A	107	ALA	4.6
1	A	274	TYR	4.6
1	A	286	VAL	4.5
1	A	337	PHE	4.2
1	A	4	LEU	4.1
1	A	108	TYR	4.1
1	A	216	MET	4.1
1	A	113	ASP	3.9
1	A	6	VAL	3.8
1	A	241	VAL	3.8
1	A	144	ILE	3.8
1	A	115	VAL	3.7
1	A	249	VAL	3.6

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	A	288	ALA	3.6
1	A	284	ILE	3.5
1	A	151	ALA	3.5
1	A	238	ARG	3.4
1	A	143	GLY	3.4
1	A	239	THR	3.4
1	A	178	LEU	3.4
1	A	149	VAL	3.3
1	A	142	VAL	3.2
1	A	289	VAL	3.2
1	A	187	GLY	3.2
1	A	169	GLU	3.1
1	A	197	LEU	3.1
1	A	210	PHE	3.1
1	A	109	LEU	3.0
1	A	150	PHE	3.0
1	A	306	ILE	3.0
1	A	3	VAL	3.0
1	A	192	GLU	3.0
1	A	154	ALA	2.9
1	A	246	GLY	2.9
1	A	212	LYS	2.9
1	A	278	LYS	2.9
1	A	117	ASP	2.8
1	A	265	LEU	2.8
1	A	253	ARG	2.8
1	A	97	GLU	2.8
1	A	8	PHE	2.8
1	A	189	ILE	2.7
1	A	233	TYR	2.7
1	A	188	ASN	2.7
1	A	38	GLU	2.7
1	A	224	TYR	2.7
1	A	193	LYS	2.6
1	A	196	LYS	2.6
1	A	176	ARG	2.6
1	A	252	LYS	2.6
1	A	11	PHE	2.5
1	A	153	ILE	2.5
1	A	315	SER	2.5
1	A	311	ALA	2.5
1	A	13	ALA	2.5

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	A	326	ASP	2.4
1	A	341	ILE	2.4
1	A	62	VAL	2.4
1	A	179	ASP	2.3
1	A	285	HIS	2.3
1	A	61	ILE	2.3
1	A	329	LYS	2.2
1	A	336	ARG	2.2
1	A	119	ARG	2.2
1	A	203	VAL	2.2
1	A	277	ASP	2.1
1	A	215	GLY	2.1
1	A	245	ILE	2.1
1	A	222	ALA	2.1
1	A	36	ARG	2.0
1	A	173	ARG	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.


## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
5	CA	A	364	1/1	0.81	0.08	94,94,94,94	0
5	CA	A	363	1/1	0.93	0.17	71,71,71,71	0
4	DCP	A	361	28/28	0.98	0.15	34,43,52,56	0
5	CA	A	362	1/1	0.99	0.16	38,38,38,38	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers

as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.



## 6.5 Other polymers ⓘ

There are no such residues in this entry.