



# Full wwPDB X-ray Structure Validation Report ⓘ

Aug 9, 2020 – 08:32 AM BST

PDB ID : 6QZZ  
Title : full length OphA V404E in complex with SAH  
Authors : Song, H.; Naismith, J.H.  
Deposited on : 2019-03-12  
Resolution : 1.85 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.13.1  
buster-report : 1.1.7 (2018)  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.13.1


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

## *X-RAY DIFFRACTION*

The reported resolution of this entry is 1.85 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	2469 (1.86-1.86)
Clashscore	141614	2625 (1.86-1.86)
Ramachandran outliers	138981	2592 (1.86-1.86)
Sidechain outliers	138945	2592 (1.86-1.86)
RSRZ outliers	127900	2436 (1.86-1.86)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	416	
2	B	416	

## 2 Entry composition [i](#)

There are 4 unique types of molecules in this entry. The entry contains 6521 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Peptide N-methyltransferase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	393	Total	C	N	O	S	0	4	0
			3081	1963	527	574	17			

- Molecule 2 is a protein called Peptide N-methyltransferase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	392	Total	C	N	O	S	0	3	0
			3064	1951	521	575	17			

- Molecule 3 is S-ADENOSYL-L-HOMOCYSTEINE (three-letter code: SAH) (formula:  $C_{14}H_{20}N_6O_5S$ ).




- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	159	Total 159	O 159	0	0
4	B	165	Total 165	O 165	0	0


### 3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

#### • Molecule 1: Peptide N-methyltransferase



#### • Molecule 2: Peptide N-methyltransferase



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 2	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	164.00 Å 91.72 Å 85.22 Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	80.05 – 1.85 80.05 – 1.85	Depositor EDS
% Data completeness (in resolution range)	99.9 (80.05-1.85) 99.9 (80.05-1.85)	Depositor EDS
$R_{merge}$	0.04	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.67 (at 1.84 Å)	Xtriage
Refinement program	REFMAC 5.8.0238	Depositor
R, $R_{free}$	0.206 , 0.225 0.215 , 0.234	Depositor DCC
$R_{free}$ test set	5568 reflections (5.05%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	35.8	Xtriage
Anisotropy	0.081	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.38 , 48.5	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.96	EDS
Total number of atoms	6521	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	50.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 85.93 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 1.0908e-07. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality

### 5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: EME, MVA, SAH

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.68	1/3145 (0.0%)	0.76	1/4276 (0.0%)
2	B	0.68	1/3108 (0.0%)	0.76	1/4225 (0.0%)
All	All	0.68	2/6253 (0.0%)	0.76	2/8501 (0.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1
2	B	0	2
All	All	0	3

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	201	GLU	C-O	8.93	1.40	1.23
2	B	201	GLU	C-O	8.65	1.39	1.23

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed( $^{\circ}$ )	Ideal( $^{\circ}$ )
2	B	145	PRO	N-CA-CB	-5.91	96.09	102.60
1	A	145	PRO	N-CA-CB	-5.50	96.54	102.60

There are no chirality outliers.

All (3) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	266	LEU	Peptide
2	B	186	PHE	Peptide
2	B	266	LEU	Peptide

## 5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3081	0	3053	42	0
2	B	3064	0	3037	41	0
3	A	26	0	19	0	0
3	B	26	0	19	0	0
4	A	159	0	0	0	0
4	B	165	0	0	0	0
All	All	6521	0	6128	65	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (65) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:44:ILE:HG13	2:B:98[B]:TYR:CE1	2.14	0.82
1:A:104:PHE:CZ	1:A:171[B]:PHE:HD2	2.01	0.79
2:B:44:ILE:HG13	2:B:98[B]:TYR:CZ	2.23	0.72
2:B:402:ILE:O	2:B:402:ILE:HG23	1.92	0.70
1:A:402:ILE:HG13	2:B:106:ASN:OD1	1.92	0.69
1:A:104:PHE:CZ	1:A:171[B]:PHE:CD2	2.82	0.68
2:B:404:EME:H3	2:B:405:GLY:N	2.11	0.66
1:A:406:VAL:O	1:A:410:ILE:HG12	1.95	0.66
1:A:202:TYR:O	1:A:251:LYS:NZ	2.29	0.65
1:A:104:PHE:CD2	2:B:143:SER:HB3	2.31	0.65
1:A:143:SER:HB3	2:B:104:PHE:CD2	2.34	0.63
1:A:131:GLU:HB2	1:A:171[B]:PHE:HZ	1.62	0.63
1:A:176:VAL:HG23	1:A:190:LYS:HB2	1.82	0.61
1:A:104:PHE:CE1	1:A:171[B]:PHE:CD2	2.88	0.60
1:A:401:MVA:HN2	1:A:401:MVA:HG22	1.85	0.58

*Continued on next page...*



*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:272:VAL:HG13	2:B:409:VAL:HG13	1.87	0.55
1:A:44:ILE:CG1	2:B:402:ILE:HG21	2.37	0.55
2:B:372:ARG:HD2	2:B:400:TRP:CH2	2.43	0.54
2:B:267:LEU:O	2:B:269:ALA:N	2.41	0.53
1:A:402:ILE:CG1	2:B:106:ASN:OD1	2.55	0.53
1:A:267:LEU:O	1:A:269:ALA:N	2.43	0.52
2:B:81:GLU:HA	2:B:84:VAL:HG22	1.91	0.51
1:A:376:LYS:HB2	1:A:378:MET:HE3	1.93	0.51
2:B:265:GLU:HA	2:B:267:LEU:HG	1.92	0.51
1:A:316:GLN:HB3	1:A:378:MET:HE2	1.92	0.50
1:A:399:PRO:CG	2:B:44:ILE:HD13	2.41	0.50
2:B:162:VAL:HG21	2:B:201:GLU:HB3	1.94	0.50
1:A:265:GLU:HA	1:A:267:LEU:HG	1.93	0.50
2:B:199:GLU:CD	2:B:232:ARG:HH22	2.14	0.50
1:A:162:VAL:HG21	1:A:201:GLU:HB3	1.94	0.50
1:A:272:VAL:HA	1:A:410:ILE:HD11	1.93	0.49
1:A:398:PHE:N	1:A:399:PRO:CD	2.77	0.48
1:A:27:LEU:HD23	2:B:27:LEU:HD23	1.96	0.47
2:B:404:EME:C7	2:B:405:GLY:N	2.76	0.47
1:A:104:PHE:CE2	2:B:143:SER:HB3	2.50	0.46
1:A:132:ASP:OD2	2:B:132:ASP:OD2	2.32	0.46
1:A:148:LEU:HG	2:B:148:LEU:HG	1.96	0.45
1:A:104:PHE:CD2	2:B:143:SER:CB	2.98	0.45
1:A:44:ILE:HD12	1:A:98:TYR:CZ	2.51	0.45
1:A:399:PRO:O	2:B:98[A]:TYR:OH	2.14	0.45
1:A:104:PHE:CE2	2:B:143:SER:CB	2.99	0.45
2:B:401:MVA:HG22	2:B:401:MVA:HN2	1.99	0.45
1:A:400:TRP:HA	1:A:401:MVA:HN1	1.80	0.44
2:B:44:ILE:HG21	2:B:215:MET:CE	2.47	0.44
1:A:44:ILE:CG2	2:B:400:TRP:CE3	3.01	0.44
1:A:316:GLN:HB3	1:A:378:MET:CE	2.49	0.43
2:B:327:VAL:HG11	2:B:363:LEU:HD11	2.01	0.43
1:A:376:LYS:HB2	1:A:378:MET:CE	2.49	0.43
1:A:398:PHE:N	1:A:399:PRO:HD3	2.33	0.43
2:B:400:TRP:HA	2:B:401:MVA:HN1	1.78	0.43
1:A:176:VAL:HG22	1:A:188:ASN:OD1	2.19	0.42
1:A:143:SER:HB3	2:B:104:PHE:CE2	2.55	0.42
2:B:199:GLU:OE2	2:B:232:ARG:NH2	2.48	0.42
1:A:376:LYS:CB	1:A:378:MET:CE	2.98	0.42
1:A:143:SER:CB	2:B:104:PHE:CD2	3.02	0.42
2:B:13:THR:O	2:B:94:VAL:HA	2.21	0.41

*Continued on next page...*

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:44:ILE:HG12	2:B:402:ILE:HG21	2.03	0.41
1:A:327:VAL:HG11	1:A:363:LEU:HD11	2.03	0.41
2:B:73:LEU:HD12	2:B:73:LEU:HA	1.98	0.41
1:A:144[A]:ASN:HA	1:A:145:PRO:HA	1.88	0.41
1:A:158:ARG:NH2	2:B:409:VAL:HG23	2.36	0.40
2:B:144[A]:ASN:HA	2:B:145:PRO:HA	1.88	0.40
2:B:402:ILE:HA	2:B:403:MVA:HN1	1.89	0.40
2:B:196:ASP:OD1	2:B:232:ARG:NH1	2.54	0.40
2:B:404:EME:H3	2:B:405:GLY:H	1.86	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles ⓘ

### 5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	392/416 (94%)	373 (95%)	17 (4%)	2 (0%)	29	15
2	B	388/416 (93%)	369 (95%)	17 (4%)	2 (0%)	29	15
All	All	780/832 (94%)	742 (95%)	34 (4%)	4 (0%)	29	15

All (4) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	268	PRO
2	B	268	PRO
2	B	145	PRO
1	A	145	PRO

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	329/343 (96%)	329 (100%)	0	100	100
2	B	326/341 (96%)	325 (100%)	1 (0%)	92	91
All	All	655/684 (96%)	654 (100%)	1 (0%)	93	92

All (1) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
2	B	367	ASP

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

4 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
2	EME	B	404	2	5,9,10	0.56	0	4,10,12	1.35	0
1	MVA	A	401	1	6,7,8	0.59	0	7,8,10	0.91	0
2	MVA	B	403	2	6,7,8	0.60	0	7,8,10	1.09	1 (14%)
2	MVA	B	401	2	6,7,8	0.80	0	7,8,10	1.57	2 (28%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	EME	B	404	2	-	4/4/9/11	-
1	MVA	A	401	1	-	5/6/8/10	-
2	MVA	B	403	2	-	2/6/8/10	-
2	MVA	B	401	2	-	6/6/8/10	-

There are no bond length outliers.

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	401	MVA	CB-CA-C	-3.05	109.21	113.04
2	B	403	MVA	O-C-CA	-2.52	117.80	124.83
2	B	401	MVA	O-C-CA	-2.15	118.85	124.83

There are no chirality outliers.

All (17) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	B	404	EME	O-C-CA-CB
2	B	404	EME	N2-CA-CB-CG
2	B	404	EME	C-CA-CB-CG
2	B	404	EME	CA-CB-CG-CD
1	A	401	MVA	N-CA-CB-CG1
1	A	401	MVA	N-CA-CB-CG2
1	A	401	MVA	C-CA-CB-CG1
1	A	401	MVA	C-CA-CB-CG2
2	B	403	MVA	CB-CA-N-CN
2	B	403	MVA	O-C-CA-CB
2	B	401	MVA	N-CA-CB-CG1
2	B	401	MVA	N-CA-CB-CG2
2	B	401	MVA	C-CA-CB-CG1
2	B	401	MVA	C-CA-CB-CG2
1	A	401	MVA	CB-CA-N-CN
2	B	401	MVA	CB-CA-N-CN
2	B	401	MVA	O-C-CA-CB

There are no ring outliers.

4 monomers are involved in 8 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	B	404	EME	3	0
1	A	401	MVA	2	0
2	B	403	MVA	1	0
2	B	401	MVA	2	0

## 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 5.6 Ligand geometry [i](#)

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
3	SAH	B	501	-	21,28,28	0.63	0	20,40,40	0.84	1 (5%)
3	SAH	A	501	-	21,28,28	0.71	0	20,40,40	0.74	1 (5%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	SAH	B	501	-	-	0/7/31/31	0/3/3/3
3	SAH	A	501	-	-	0/7/31/31	0/3/3/3

There are no bond length outliers.

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	B	501	SAH	C5-C6-N6	2.30	123.85	120.35

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	501	SAH	C5-C6-N6	2.25	123.76	120.35


There are no chirality outliers.


There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data ⓘ

### 6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	392/416 (94%)	0.70	37 (9%) 8 8	26, 48, 84, 113	0
2	B	389/416 (93%)	0.58	28 (7%) 15 15	27, 45, 80, 126	0
All	All	781/832 (93%)	0.64	65 (8%) 11 11	26, 47, 82, 126	0

All (65) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	267	LEU	11.1
2	B	267	LEU	9.8
2	B	268	PRO	5.5
1	A	398	PHE	5.0
2	B	409	VAL	4.8
1	A	229	ALA	4.7
2	B	271	GLN	4.7
1	A	258	LEU	4.7
1	A	234	PRO	4.5
1	A	268	PRO	4.2
2	B	402	ILE	4.2
2	B	411	GLY	4.1
2	B	272	VAL	4.1
2	B	400	TRP	4.0
1	A	409	VAL	3.9
1	A	410	ILE	3.6
2	B	412	SER	3.5
1	A	406	VAL	3.5
1	A	400	TRP	3.4
2	B	270	GLY	3.4
1	A	269	ALA	3.4
2	B	234	PRO	3.3
1	A	238	LYS	3.3
1	A	204	ALA	3.2

*Continued on next page...*



*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
2	B	269	ALA	3.2
1	A	198	LEU	3.2
2	B	228	VAL	3.1
2	B	260	ILE	3.1
1	A	270	GLY	3.1
1	A	288	ASP	3.0
1	A	289	VAL	3.0
2	B	410	ILE	3.0
2	B	229	ALA	3.0
1	A	194	LEU	2.9
2	B	289	VAL	2.9
2	B	273	PRO	2.9
1	A	271	GLN	2.9
1	A	231	LEU	2.8
1	A	157	ILE	2.8
1	A	205	GLU	2.8
2	B	186	PHE	2.8
1	A	104	PHE	2.8
1	A	187	ASP	2.7
1	A	396	ASN	2.6
1	A	261	ILE	2.6
2	B	236	ILE	2.6
2	B	231	LEU	2.5
2	B	194	LEU	2.5
1	A	189	ASN	2.4
2	B	264	LEU	2.4
2	B	251	LYS	2.4
1	A	251	LYS	2.3
1	A	369	TRP	2.3
1	A	260	ILE	2.3
1	A	202	TYR	2.2
2	B	266	LEU	2.2
1	A	272	VAL	2.2
2	B	261	ILE	2.1
2	B	205	GLU	2.1
1	A	226	TYR	2.1
1	A	235	GLU	2.1
2	B	346	ARG	2.0
1	A	346	ARG	2.0
1	A	266	LEU	2.0
1	A	276	LYS	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
2	MVA	B	401	8/9	0.74	0.18	59,64,71,73	0
2	MVA	B	403	8/9	0.91	0.14	41,47,67,71	0
2	EME	B	404	10/11	0.91	0.25	40,54,109,128	0
1	MVA	A	401	8/9	0.98	0.11	35,41,51,58	0

## 6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 6.4 Ligands [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.


Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
3	SAH	A	501	26/26	0.92	0.12	27,33,43,46	0
3	SAH	B	501	26/26	0.97	0.12	28,32,36,37	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

**Electron density around SAH A 501:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)





## 6.5 Other polymers [i](#)

There are no such residues in this entry.