



Full wwPDB X-ray Structure Validation Report ⓘ

Apr 26, 2022 – 10:08 am BST

PDB ID : 7QZQ
Title : Crystal structure of the kelch domain of human KBTBD12
Authors : Manning, C.; Chen, Z.; Chen, X.; Bradshaw, W.J.; Bakshi, S.; Mckinley, G.;
Chalk, R.; Burgess-Brown, N.; von Delft, F.; Bullock, A.N.
Deposited on : 2022-01-31
Resolution : 1.88 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	1.8.4, CSD as541be (2020)
Xtriage (Phenix)	:	1.13
EDS	:	2.28
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac	:	5.8.0267
CCP4	:	7.1.010 (Gargrove)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.28


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.88 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	9470 (1.90-1.86)
Clashscore	141614	10282 (1.90-1.86)
Ramachandran outliers	138981	10152 (1.90-1.86)
Sidechain outliers	138945	10152 (1.90-1.86)
RSRZ outliers	127900	9303 (1.90-1.86)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	362	<div> <div style="width: 100%; height: 10px; background: linear-gradient(to right, red 0%, red 1%, green 1%, green 88%, yellow 88%, yellow 95%, grey 95%);"></div> <div style="display: flex; justify-content: space-between; width: 90%; margin: 0 auto;"> % 88% 7% 5% </div> </div>
1	B	362	<div> <div style="width: 100%; height: 10px; background: linear-gradient(to right, red 0%, red 6%, green 6%, green 86%, yellow 86%, yellow 94%, grey 94%);"></div> <div style="display: flex; justify-content: space-between; width: 90%; margin: 0 auto;"> 6% 86% 8% 6% </div> </div>

2 Entry composition [i](#)

There are 4 unique types of molecules in this entry. The entry contains 5761 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Kelch repeat and BTB domain-containing protein 12.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	344	Total	C	N	O	S	0	4	0
			2683	1700	458	508	17			
1	B	342	Total	C	N	O	S	0	1	0
			2627	1666	443	501	17			

There are 6 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	263	SER	-	expression tag	UNP Q3ZCT8
A	264	MET	-	expression tag	UNP Q3ZCT8
A	624	LEU	-	expression tag	UNP Q3ZCT8
B	263	SER	-	expression tag	UNP Q3ZCT8
B	264	MET	-	expression tag	UNP Q3ZCT8
B	624	LEU	-	expression tag	UNP Q3ZCT8

- Molecule 2 is 1,2-ETHANEDIOL (three-letter code: EDO) (formula: C₂H₆O₂).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	A	1	Total C O 4 2 2	0	0

- Molecule 3 is SODIUM ION (three-letter code: NA) (formula: Na).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	B	1	Total Na 1 1	0	0


- Molecule 4 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	243	Total O 243 243	0	0
4	B	203	Total O 203 203	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Kelch repeat and BTB domain-containing protein 12



- Molecule 1: Kelch repeat and BTB domain-containing protein 12



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	72.38Å 55.08Å 95.16Å 90.00° 108.67° 90.00°	Depositor
Resolution (Å)	45.08 – 1.88 45.08 – 1.88	Depositor EDS
% Data completeness (in resolution range)	98.2 (45.08-1.88) 98.2 (45.08-1.88)	Depositor EDS
R_{merge}	0.16	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.20 (at 1.88Å)	Xtriage
Refinement program	PHENIX 1.20.1_4487	Depositor
R, R_{free}	0.195 , 0.239 0.197 , 0.200	Depositor DCC
R_{free} test set	2859 reflections (5.04%)	wwPDB-VP
Wilson B-factor (Å ²)	32.0	Xtriage
Anisotropy	0.626	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	(Not available) , (Not available)	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.34$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	5761	wwPDB-VP
Average B, all atoms (Å ²)	40.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.49% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: EDO, NA

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.40	0/2759	0.61	0/3755
1	B	0.40	0/2690	0.61	0/3671
All	All	0.40	0/5449	0.61	0/7426

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2683	0	2587	15	0
1	B	2627	0	2479	16	0
2	A	4	0	6	0	0
3	B	1	0	0	0	0
4	A	243	0	0	0	0
4	B	203	0	0	3	1
All	All	5761	0	5072	31	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (31) close contacts within the same asymmetric unit are listed below, sorted by their clash

magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:512:LYS:NZ	4:B:801:HOH:O	2.20	0.74
1:B:590:VAL:HG23	1:B:592:MET:HG3	1.70	0.72
1:B:287:ILE:HD12	1:B:593:HIS:CE1	2.35	0.61
1:A:506:ASP:OD2	1:A:511:ARG:NH2	2.37	0.57
1:B:396:ILE:HD13	1:B:401:LEU:HD21	1.88	0.56
1:A:395:LYS:O	1:A:396:ILE:HD13	2.10	0.52
1:A:590:VAL:HG23	1:A:592:MET:HG2	1.92	0.52
1:A:359:HIS:O	1:A:364:GLN:HA	2.11	0.50
1:A:344:LYS:O	1:A:348:GLN:HG3	2.12	0.50
1:B:486:SER:HB2	1:B:497:LEU:HD13	1.94	0.49
1:A:392:GLY:O	1:A:403:THR:HG22	2.13	0.48
1:B:480:TYR:CE2	1:B:503:VAL:HG22	2.47	0.48
1:A:314:PRO:HG2	1:A:323:VAL:HG21	1.96	0.48
1:A:480:TYR:CE2	1:A:503:VAL:HG22	2.48	0.48
1:B:543:ASN:HB2	4:B:921:HOH:O	2.14	0.46
1:A:290:ARG:NH1	1:A:290:ARG:HG2	2.31	0.45
1:B:379:ALA:HB1	1:B:431:VAL:HG12	2.00	0.43
1:B:301:TYR:CD2	1:B:573:ILE:HD13	2.53	0.43
1:A:452:ASP:OD1	1:A:452:ASP:N	2.52	0.43
1:A:485:PHE:CD2	1:A:496:VAL:HG13	2.54	0.43
1:B:393:GLN:NE2	4:B:816:HOH:O	2.50	0.43
1:A:480:TYR:CZ	1:A:516:VAL:HG21	2.54	0.42
1:B:449:ASP:OD2	1:B:457:ARG:NH1	2.45	0.42
1:B:354:GLU:OE1	1:B:371:THR:HG22	2.20	0.41
1:B:462:LEU:HD23	1:B:475:ARG:HB2	2.03	0.41
1:A:423:PRO:HD2	1:A:463:LEU:HD13	2.02	0.41
1:A:460:ASN:O	1:A:478:MET:HB2	2.21	0.41
1:A:308:THR:O	1:A:612:PRO:HG3	2.21	0.41
1:B:438:LEU:HB2	1:B:465:TYR:HB3	2.03	0.41
1:B:478:MET:HG2	1:B:527:TRP:CZ2	2.57	0.40
1:B:449:ASP:HB2	1:B:457:ARG:HH22	1.86	0.40

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
4:B:824:HOH:O	4:B:972:HOH:O[2_455]	2.17	0.03

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	346/362 (96%)	336 (97%)	10 (3%)	0	100	100
1	B	341/362 (94%)	334 (98%)	7 (2%)	0	100	100
All	All	687/724 (95%)	670 (98%)	17 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	291/319 (91%)	291 (100%)	0	100	100
1	B	277/319 (87%)	275 (99%)	2 (1%)	84	83
All	All	568/638 (89%)	566 (100%)	2 (0%)	91	90

All (2) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	394	MET
1	B	418	ARG

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 2 ligands modelled in this entry, 1 is monoatomic - leaving 1 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	EDO	A	701	-	3,3,3	0.37	0	2,2,2	0.51	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	EDO	A	701	-	-	1/1/1/1	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (1) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	A	701	EDO	O1-C1-C2-O2

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	344/362 (95%)	-0.04	4 (1%) 79 80	27, 37, 58, 79	0
1	B	342/362 (94%)	0.34	23 (6%) 17 19	26, 37, 72, 119	0
All	All	686/724 (94%)	0.15	27 (3%) 39 41	26, 37, 65, 119	0

All (27) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	342	ALA	11.8
1	B	345	LEU	8.8
1	B	347	ARG	7.0
1	B	348	GLN	6.5
1	B	363	ASN	6.4
1	B	289	SER	5.5
1	B	291	HIS	5.1
1	B	344	LYS	4.8
1	A	289	SER	3.5
1	B	365	PHE	3.3
1	B	341	SER	3.2
1	B	346	SER	3.2
1	B	364	GLN	2.9
1	B	316	TYR	2.8
1	B	294	TYR	2.8
1	B	340	ALA	2.6
1	B	372	ALA	2.4
1	B	290	ARG	2.4
1	B	607	ARG	2.4
1	B	350	ASN	2.3
1	B	288	ARG	2.2
1	B	349	LYS	2.2
1	B	362	GLY	2.1
1	A	517	VAL	2.1

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	319	GLY	2.1
1	A	294	TYR	2.0
1	A	361	ARG	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
3	NA	B	701	1/1	0.93	0.14	54,54,54,54	0
2	EDO	A	701	4/4	0.94	0.08	34,36,36,43	0

6.5 Other polymers [i](#)

There are no such residues in this entry.