



# wwPDB X-ray Structure Validation Summary Report ⓘ

Aug 8, 2020 – 02:57 AM BST

PDB ID : 6R0X  
Title : The extracellular domain of G6b-B in complex with Fab fragment and DP12 heparin oligosaccharide.  
Authors : Ogg, D.J.; McMiken, H.J.; Howard, T.D.  
Deposited on : 2019-03-13  
Resolution : 3.13 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:


MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.13.1  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.13.1

# 1 Overall quality at a glance i

The following experimental techniques were used to determine the structure:  
*X-RAY DIFFRACTION*

The reported resolution of this entry is 3.13 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.






Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	1626 (3.18-3.10)
Clashscore	141614	1735 (3.18-3.10)
Ramachandran outliers	138981	1677 (3.18-3.10)
Sidechain outliers	138945	1677 (3.18-3.10)
RSRZ outliers	127900	1588 (3.18-3.10)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	240	<div><div>2%</div><div></div><div>69%</div><div>18%</div><div>•</div><div>12%</div></div>
1	C	240	<div><div>%</div><div></div><div>71%</div><div>16%</div><div>•</div><div>11%</div></div>
2	B	234	<div><div>%</div><div></div><div>69%</div><div>19%</div><div>•</div><div>9%</div></div>
2	D	234	<div><div>6%</div><div></div><div>69%</div><div>19%</div><div>•</div><div>10%</div></div>
3	E	116	<div><div>2%</div><div></div><div>53%</div><div>28%</div><div>•</div><div>15%</div></div>
3	F	116	<div><div>%</div><div></div><div>59%</div><div>17%</div><div>8%</div><div>16%</div></div>

Continued on next page...

*Continued from previous page...*

Mol	Chain	Length	Quality of chain
4	G	3	 33% 33% 33%
4	I	3	 100%
5	H	8	 100%

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
4	A2G	G	1	X	-	-	-
4	A2G	I	1	X	-	-	-
5	IDS	H	2	X	-	-	-
5	IDS	H	6	X	-	-	-

## 2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 7896 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called antibody fab fragment heavy chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	212	Total	C	N	O	S	0	0	0
			1575	996	255	319	5			
1	C	213	Total	C	N	O	S	0	0	0
			1576	999	252	319	6			

- Molecule 2 is a protein called antibody fab fragment light chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	214	Total	C	N	O	S	0	0	0
			1597	995	257	339	6			
2	D	211	Total	C	N	O	S	0	0	0
			1537	949	249	332	7			

- Molecule 3 is a protein called Megakaryocyte and platelet inhibitory receptor G6b.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	E	99	Total	C	N	O	S	0	0	0
			693	438	127	124	4			
3	F	98	Total	C	N	O	S	0	0	0
			688	438	124	123	3			

There are 10 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
E	32	ASP	ASN	engineered mutation	UNP O95866
E	67	ALA	SER	engineered mutation	UNP O95866
E	68	ALA	SER	engineered mutation	UNP O95866
E	69	ALA	SER	engineered mutation	UNP O95866
E	71	ALA	THR	engineered mutation	UNP O95866
F	32	ASP	ASN	engineered mutation	UNP O95866
F	67	ALA	SER	engineered mutation	UNP O95866
F	68	ALA	SER	engineered mutation	UNP O95866

*Continued on next page...*

Continued from previous page...


Chain	Residue	Modelled	Actual	Comment	Reference
F	69	ALA	SER	engineered mutation	UNP O95866
F	71	ALA	THR	engineered mutation	UNP O95866

- Molecule 4 is an oligosaccharide called beta-D-galactopyranose-(1-3)-[N-acetyl-alpha-neuraminic acid-(2-6)]2-acetamido-2-deoxy-alpha-D-galactopyranose.



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
4	G	3	Total	C	N	O	0	0	0
			45	25	2	18			
4	I	3	Total	C	N	O	0	0	0
			45	25	2	18			

- Molecule 5 is an oligosaccharide called 2-O-sulfo-alpha-L-idopyranuronic acid-(1-4)-2-deoxy-6-O-sulfo-2-(sulfoamino)-alpha-D-glucopyranose-(1-4)-2-O-sulfo-alpha-L-idopyranuronic acid-(1-4)-2-deoxy-6-O-sulfo-2-(sulfoamino)-alpha-D-glucopyranose-(1-4)-2-O-sulfo-alpha-L-idopyranuronic acid-(1-4)-2-deoxy-6-O-sulfo-2-(sulfoamino)-alpha-D-glucopyranose-(1-4)-2-O-sulfo-alpha-L-idopyranuronic acid-(1-4)-2-deoxy-6-O-sulfo-2-(sulfoamino)-alpha-D-glucopyranose.




Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
5	H	8	Total	C	N	O	S	0	0	0
			140	48	4	76	12			


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: antibody fab fragment heavy chain




- Molecule 1: antibody fab fragment heavy chain




- Molecule 2: antibody fab fragment light chain




- Molecule 2: antibody fab fragment light chain






- Molecule 3: Megakaryocyte and platelet inhibitory receptor G6b



- Molecule 3: Megakaryocyte and platelet inhibitory receptor G6b



- Molecule 4: beta-D-galactopyranose-(1-3)-[N-acetyl-alpha-neuraminic acid-(2-6)]2-acetamido-2-deoxy-alpha-D-galactopyranose



- Molecule 4: beta-D-galactopyranose-(1-3)-[N-acetyl-alpha-neuraminic acid-(2-6)]2-acetamido-2-deoxy-alpha-D-galactopyranose



- Molecule 5: 2-O-sulfo-alpha-L-idopyranuronic acid-(1-4)-2-deoxy-6-O-sulfo-2-(sulfoamino)-alpha-D-glucopyranose-(1-4)-2-O-sulfo-alpha-L-idopyranuronic acid-(1-4)-2-deoxy-6-O-sulfo-2-(sulfoamino)-alpha-D-glucopyranose-(1-4)-2-O-sulfo-alpha-L-idopyranuronic acid-(1-4)-2-deoxy-6-O-sulfo-2-(sulfoamino)-alpha-D-glucopyranose-(1-4)-2-O-sulfo-alpha-L-idopyranuronic acid-(1-4)-2-deoxy-6-O-sulfo-2-(sulfoamino)-alpha-D-glucopyranose

Chain H: 

100%

SGH1  
IDS2  
SGH3  
IDS4  
SGH5  
IDS6  
SGH7  
IDS8



## 4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	183.80Å 72.34Å 131.04Å 90.00° 124.52° 90.00°	Depositor
Resolution (Å)	63.10 – 3.13 63.10 – 3.13	Depositor EDS
% Data completeness (in resolution range)	97.1 (63.10-3.13) 97.2 (63.10-3.13)	Depositor EDS
$R_{merge}$	0.16	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.25 (at 3.13Å)	Xtriage
Refinement program	REFMAC 5.8.0238	Depositor
R, $R_{free}$	0.230 , 0.266 0.256 , 0.247	Depositor DCC
$R_{free}$ test set	1253 reflections (5.11%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	91.3	Xtriage
Anisotropy	0.415	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.30 , 103.3	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.44$ , $\langle L^2 \rangle = 0.26$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.91	EDS
Total number of atoms	7896	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	115.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.80% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: A2G, SIA, GAL, IDS, SGN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.42	0/1617	0.68	0/2219
1	C	0.44	0/1618	0.69	0/2222
2	B	0.47	0/1632	0.74	0/2229
2	D	0.42	0/1569	0.68	0/2146
3	E	0.64	0/710	0.88	0/966
3	F	0.64	0/705	0.89	1/960 (0.1%)
All	All	0.48	0/7851	0.73	1/10742 (0.0%)

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	F	93	ARG	N-CA-C	-5.18	97.01	111.00

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1575	0	1482	14	0
1	C	1576	0	1490	15	0
2	B	1597	0	1441	26	0
2	D	1537	0	1348	20	0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	E	693	0	643	21	0
3	F	688	0	644	18	0
4	G	45	0	37	4	0
4	I	45	0	37	1	0
5	H	140	0	48	0	0
All	All	7896	0	7170	111	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 7.

The worst 5 of 111 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:D:54:ASN:HD22	2:D:70:TYR:H	1.32	0.76
2:B:86:GLY:HA3	2:B:91:TYR:HA	1.68	0.74
1:A:22:GLN:HB3	1:A:44:SER:HB2	1.69	0.73
2:D:213:THR:OG1	2:D:228:SER:HA	1.92	0.69
2:D:168:TRP:HB3	2:D:214:CYS:HA	1.76	0.65

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	208/240 (87%)	198 (95%)	8 (4%)	2 (1%)	15	47
1	C	209/240 (87%)	193 (92%)	16 (8%)	0	100	100
2	B	212/234 (91%)	190 (90%)	20 (9%)	2 (1%)	17	50
2	D	207/234 (88%)	190 (92%)	15 (7%)	2 (1%)	15	47
3	E	93/116 (80%)	74 (80%)	17 (18%)	2 (2%)	6	27

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
3	F	92/116 (79%)	71 (77%)	18 (20%)	3 (3%)	4	19
All	All	1021/1180 (86%)	916 (90%)	94 (9%)	11 (1%)	14	45

5 of 11 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	84	GLN
2	D	27	THR
3	F	45	VAL
2	B	189	LYS
3	E	73	THR

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	174/206 (84%)	152 (87%)	22 (13%)	4	17
1	C	176/206 (85%)	158 (90%)	18 (10%)	7	25
2	B	177/209 (85%)	150 (85%)	27 (15%)	2	11
2	D	168/209 (80%)	147 (88%)	21 (12%)	4	18
3	E	65/91 (71%)	50 (77%)	15 (23%)	1	3
3	F	64/91 (70%)	53 (83%)	11 (17%)	2	8
All	All	824/1012 (81%)	710 (86%)	114 (14%)	3	15

5 of 114 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	C	42	THR
1	C	213	TRP
3	F	60	ARG
1	C	69	ARG
1	C	166	LEU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 14 such sidechains are listed below:

Mol	Chain	Res	Type
2	B	157	ASN
2	B	186	GLN
3	E	121	HIS
2	B	58	GLN
2	D	54	ASN

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates ⓘ

14 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
4	A2G	G	1	3,4	14,14,15	0.57	0	17,19,21	1.39	3 (17%)
4	GAL	G	2	4	11,11,12	0.56	0	15,15,17	0.87	0
4	SIA	G	3	4	17,20,21	0.30	0	21,28,31	1.56	4 (19%)
5	SGN	H	1	5	19,20,20	0.66	1 (5%)	24,31,31	2.07	4 (16%)
5	IDS	H	2	5	13,16,17	0.68	0	15,24,26	3.40	4 (26%)
5	SGN	H	3	5	18,19,20	0.39	0	22,29,31	2.17	4 (18%)
5	IDS	H	4	5	13,16,17	1.04	1 (7%)	15,24,26	3.00	6 (40%)
5	SGN	H	5	5	18,19,20	0.52	0	22,29,31	2.72	4 (18%)
5	IDS	H	6	5	13,16,17	0.59	0	15,24,26	2.92	5 (33%)
5	SGN	H	7	5	18,19,20	3.17	2 (11%)	22,29,31	2.26	5 (22%)

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
5	IDS	H	8	5	12,15,17	0.82	1 (8%)	12,22,26	2.69	4 (33%)
4	A2G	I	1	3,4	14,14,15	0.33	0	17,19,21	1.33	2 (11%)
4	GAL	I	2	4	11,11,12	0.37	0	15,15,17	0.71	0
4	SIA	I	3	4	17,20,21	0.26	0	21,28,31	1.24	3 (14%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	A2G	G	1	3,4	1/1/5/7	2/6/23/26	0/1/1/1
4	GAL	G	2	4	-	1/2/19/22	0/1/1/1
4	SIA	G	3	4	-	9/14/34/38	0/1/1/1
5	SGN	H	1	5	-	6/11/31/31	0/1/1/1
5	IDS	H	2	5	1/1/6/7	1/5/26/29	1/1/1/1
5	SGN	H	3	5	-	1/11/28/31	0/1/1/1
5	IDS	H	4	5	-	0/5/26/29	0/1/1/1
5	SGN	H	5	5	-	5/11/28/31	0/1/1/1
5	IDS	H	6	5	1/1/6/7	0/5/26/29	1/1/1/1
5	SGN	H	7	5	-	8/11/28/31	0/1/1/1
5	IDS	H	8	5	-	0/5/22/29	1/1/1/1
4	A2G	I	1	3,4	1/1/5/7	0/6/23/26	0/1/1/1
4	GAL	I	2	4	-	1/2/19/22	0/1/1/1
4	SIA	I	3	4	-	6/14/34/38	0/1/1/1

All (5) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
5	H	7	SGN	O1S-S1	12.99	1.56	1.42
5	H	4	IDS	O4-C4	3.14	1.50	1.43
5	H	7	SGN	O4-C4	2.66	1.49	1.43
5	H	1	SGN	O4-C4	2.39	1.48	1.43
5	H	8	IDS	C4-C5	2.30	1.56	1.52

The worst 5 of 48 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
5	H	5	SGN	O4-C4-C5	11.54	137.94	109.30
5	H	2	IDS	O4-C4-C5	9.32	127.85	110.05

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
5	H	1	SGN	O4-C4-C5	8.19	129.64	109.30
5	H	3	SGN	O4-C4-C3	8.11	129.10	110.35
5	H	7	SGN	O2S-S1-O1S	-7.52	102.39	120.16

All (4) chirality outliers are listed below:

Mol	Chain	Res	Type	Atom
4	G	1	A2G	C1
5	H	2	IDS	C4
4	I	1	A2G	C1
5	H	6	IDS	C4

5 of 40 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	I	3	SIA	C5-C6-C7-C8
4	I	3	SIA	C5-C6-C7-O7
4	I	3	SIA	O6-C6-C7-C8
4	I	3	SIA	O6-C6-C7-O7
5	H	1	SGN	C4-C5-C6-O6


All (3) ring outliers are listed below:

Mol	Chain	Res	Type	Atoms
5	H	2	IDS	C1-C2-C3-C4-C5-O5
5	H	8	IDS	C1-C2-C3-C4-C5-O5
5	H	6	IDS	C1-C2-C3-C4-C5-O5


2 monomers are involved in 5 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	G	1	A2G	4	0
4	I	2	GAL	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.







## 5.6 Ligand geometry [i](#)

There are no ligands in this entry.

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues ⓘ

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	212/240 (88%)	0.11	4 (1%) 66 48	63, 101, 132, 167	0
1	C	213/240 (88%)	0.08	3 (1%) 75 59	72, 112, 151, 181	0
2	B	214/234 (91%)	-0.04	2 (0%) 84 72	65, 111, 152, 200	0
2	D	211/234 (90%)	0.35	13 (6%) 20 9	63, 128, 279, 290	0
3	E	99/116 (85%)	0.07	2 (2%) 65 46	70, 100, 142, 252	0
3	F	98/116 (84%)	0.04	1 (1%) 82 70	75, 97, 135, 227	0
All	All	1047/1180 (88%)	0.11	25 (2%) 59 38	63, 107, 177, 290	0

The worst 5 of 25 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	D	201	LEU	10.0
3	F	43	ARG	4.2
2	D	234	CYS	4.1
2	D	228	SER	3.5
2	D	224	PRO	3.5

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)


There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.




Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
4	A2G	G	1	14/15	0.61	0.33	127,136,146,151	0
5	SGN	H	1	20/20	0.71	0.25	187,198,201,201	0
4	GAL	I	2	11/12	0.74	0.23	168,170,171,172	0
4	GAL	G	2	11/12	0.77	0.32	105,111,116,120	0
5	SGN	H	7	19/20	0.77	0.20	143,154,165,165	0
5	IDS	H	8	15/17	0.79	0.32	163,164,166,166	0
4	SIA	G	3	20/21	0.79	0.19	161,166,174,176	0
5	SGN	H	3	19/20	0.84	0.12	136,151,168,169	0
4	SIA	I	3	20/21	0.88	0.22	216,228,236,236	0
4	A2G	I	1	14/15	0.88	0.20	165,171,191,205	0
5	IDS	H	2	16/17	0.89	0.21	166,178,182,182	0
5	IDS	H	6	16/17	0.90	0.17	122,133,136,140	0
5	IDS	H	4	16/17	0.91	0.17	105,117,125,125	0
5	SGN	H	5	19/20	0.93	0.18	97,104,108,112	0

The following is a graphical depiction of the model fit to experimental electron density for oligosaccharide. Each fit is shown from different orientation to approximate a three-dimensional view.






**Electron density around Chain I:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)

**Electron density around Chain H:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)



## 6.4 Ligands [i](#)

There are no ligands in this entry.

## 6.5 Other polymers [i](#)

There are no such residues in this entry.