



# Full wwPDB X-ray Structure Validation Report ⓘ

May 17, 2020 – 08:27 pm BST

PDB ID : 4R1U  
Title : Crystal structure of Medicago truncatula cinnamoyl-CoA reductase  
Authors : Noel, J.P.; Bomati, E.K.; Louie, G.V.; Bowman, M.E.  
Deposited on : 2014-08-07  
Resolution : 2.18 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.11  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.11


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.18 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	6864 (2.20-2.16)
Clashscore	141614	7689 (2.20-2.16)
Ramachandran outliers	138981	7564 (2.20-2.16)
Sidechain outliers	138945	7564 (2.20-2.16)
RSRZ outliers	127900	6738 (2.20-2.16)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	349	<div> <div>7%</div> <div> <div></div> <div>81%</div> <div>6%</div> <div>12%</div> </div> </div>
1	B	349	<div> <div>5%</div> <div> <div></div> <div>79%</div> <div>9%</div> <div>11%</div> </div> </div>

## 2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 5139 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Cinnamoyl CoA reductase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	308	Total	C	N	O	S	0	0	0
			2406	1529	417	451	9			
1	B	309	Total	C	N	O	S	0	0	0
			2419	1539	418	453	9			

There are 34 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	-25	GLY	-	EXPRESSION TAG	UNP G7JEE5
A	-24	SER	-	EXPRESSION TAG	UNP G7JEE5
A	-23	HIS	-	EXPRESSION TAG	UNP G7JEE5
A	-22	GLY	-	EXPRESSION TAG	UNP G7JEE5
A	-21	GLY	-	EXPRESSION TAG	UNP G7JEE5
A	-20	SER	-	EXPRESSION TAG	UNP G7JEE5
A	-19	GLU	-	EXPRESSION TAG	UNP G7JEE5
A	-18	PHE	-	EXPRESSION TAG	UNP G7JEE5
A	-17	GLU	-	EXPRESSION TAG	UNP G7JEE5
A	-16	LEU	-	EXPRESSION TAG	UNP G7JEE5
A	-15	ARG	-	EXPRESSION TAG	UNP G7JEE5
A	-14	ARG	-	EXPRESSION TAG	UNP G7JEE5
A	-13	GLN	-	EXPRESSION TAG	UNP G7JEE5
A	-12	ALA	-	EXPRESSION TAG	UNP G7JEE5
A	-11	SER	-	EXPRESSION TAG	UNP G7JEE5
A	-10	CYS	-	EXPRESSION TAG	UNP G7JEE5
A	-9	LEU	-	EXPRESSION TAG	UNP G7JEE5
B	-25	GLY	-	EXPRESSION TAG	UNP G7JEE5
B	-24	SER	-	EXPRESSION TAG	UNP G7JEE5
B	-23	HIS	-	EXPRESSION TAG	UNP G7JEE5
B	-22	GLY	-	EXPRESSION TAG	UNP G7JEE5
B	-21	GLY	-	EXPRESSION TAG	UNP G7JEE5
B	-20	SER	-	EXPRESSION TAG	UNP G7JEE5
B	-19	GLU	-	EXPRESSION TAG	UNP G7JEE5
B	-18	PHE	-	EXPRESSION TAG	UNP G7JEE5

*Continued on next page...*

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
B	-17	GLU	-	EXPRESSION TAG	UNP G7JEE5
B	-16	LEU	-	EXPRESSION TAG	UNP G7JEE5
B	-15	ARG	-	EXPRESSION TAG	UNP G7JEE5
B	-14	ARG	-	EXPRESSION TAG	UNP G7JEE5
B	-13	GLN	-	EXPRESSION TAG	UNP G7JEE5
B	-12	ALA	-	EXPRESSION TAG	UNP G7JEE5
B	-11	SER	-	EXPRESSION TAG	UNP G7JEE5
B	-10	CYS	-	EXPRESSION TAG	UNP G7JEE5
B	-9	LEU	-	EXPRESSION TAG	UNP G7JEE5

- Molecule 2 is ACETATE ION (three-letter code: ACT) (formula:  $C_2H_3O_2$ ).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	A	1	Total C O 4 2 2	0	0
2	B	1	Total C O 4 2 2	0	0
2	B	1	Total C O 4 2 2	0	0


- Molecule 3 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	160	Total O 160 160	0	0
3	B	142	Total O 142 142	0	0


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

#### • Molecule 1: Cinnamoyl CoA reductase



#### • Molecule 1: Cinnamoyl CoA reductase



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	70.80 Å 96.82 Å 106.91 Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	19.98 – 2.18 19.98 – 2.18	Depositor EDS
% Data completeness (in resolution range)	98.3 (19.98-2.18) 98.3 (19.98-2.18)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	19.07 (at 2.19 Å)	Xtriage
Refinement program	PHENIX 1.8.1_1168	Depositor
R, $R_{free}$	0.198 , 0.242 0.199 , 0.243	Depositor DCC
$R_{free}$ test set	3836 reflections (10.00%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	36.2	Xtriage
Anisotropy	0.044	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.38 , 54.0	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.42$ , $\langle L^2 \rangle = 0.25$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.96	EDS
Total number of atoms	5139	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	37.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 46.29 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 1.1688e-04. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

<sup>1</sup> Intensities estimated from amplitudes.

<sup>2</sup> Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality

### 5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: ACT

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.41	0/2457	0.54	0/3344
1	B	0.41	0/2472	0.54	0/3366
All	All	0.41	0/4929	0.54	0/6710

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2406	0	2407	16	0
1	B	2419	0	2431	19	0
2	A	4	0	3	0	0
2	B	8	0	6	1	0
3	A	160	0	0	4	0
3	B	142	0	0	3	0
All	All	5139	0	4847	34	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (34) close contacts within the same asymmetric unit are listed below, sorted by their clash

magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:165:ASP:OD2	3:A:587:HOH:O	1.89	0.89
1:A:190:THR:O	3:A:617:HOH:O	2.02	0.78
1:A:186:LEU:HD22	1:A:191:ILE:HG13	1.67	0.76
1:B:309:GLU:OE2	3:B:601:HOH:O	2.05	0.74
1:B:203:ASN:O	3:B:586:HOH:O	2.06	0.73
1:A:313:LEU:HD12	1:A:314:PRO:HD2	1.72	0.71
1:A:191:ILE:HG22	1:A:192:ASN:H	1.56	0.70
1:B:1:ASP:N	3:B:630:HOH:O	2.31	0.63
1:B:126:PRO:HG3	1:B:144:HIS:ND1	2.18	0.58
1:B:1:ASP:OD1	1:B:1:ASP:N	2.42	0.52
1:A:126:PRO:HG3	1:A:144:HIS:CD2	2.46	0.51
1:A:130:ARG:HD3	1:A:277:LYS:O	2.11	0.50
1:A:188:GLN:H	1:A:188:GLN:NE2	2.11	0.49
1:A:51:GLU:HG2	1:A:52:ARG:HG3	1.94	0.49
1:B:16:TRP:CZ2	1:B:185:PRO:HG2	2.49	0.48
1:B:123:TYR:CE2	1:B:124:MET:HE2	2.48	0.48
1:B:243:GLU:HG3	1:B:294:THR:O	2.15	0.47
1:A:141:ASP:HB3	1:A:144:HIS:HB3	1.97	0.47
1:B:164:TRP:CE3	1:B:174:LEU:HD23	2.50	0.46
1:A:191:ILE:CG2	1:A:192:ASN:H	2.27	0.45
1:B:31:THR:HB	1:B:58:VAL:HB	1.99	0.45
1:A:132:VAL:O	3:A:611:HOH:O	2.20	0.45
1:B:125:ASP:OD2	1:B:141:ASP:HB2	2.16	0.45
1:B:193:ALA:HA	1:B:196:ILE:HB	1.99	0.44
1:B:189:PRO:HB3	1:B:312:HIS:CE1	2.52	0.44
1:A:109:LYS:HB2	1:B:2:GLN:NE2	2.33	0.43
1:A:191:ILE:HG22	1:A:192:ASN:N	2.29	0.43
1:A:313:LEU:HA	1:A:314:PRO:HD2	1.81	0.42
1:A:130:ARG:O	3:A:599:HOH:O	2.21	0.41
1:B:191:ILE:HD13	1:B:191:ILE:HA	1.60	0.41
1:B:141:ASP:HB3	1:B:144:HIS:HB3	2.01	0.41
1:B:243:GLU:OE1	1:B:296:VAL:HG23	2.20	0.40
1:B:248:ARG:NE	2:B:401:ACT:OXT	2.41	0.40
1:B:191:ILE:HG23	1:B:192:ASN:N	2.35	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	304/349 (87%)	292 (96%)	9 (3%)	3 (1%)	15	12
1	B	305/349 (87%)	294 (96%)	5 (2%)	6 (2%)	7	4
All	All	609/698 (87%)	586 (96%)	14 (2%)	9 (2%)	10	7

All (9) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	191	ILE
1	B	190	THR
1	B	191	ILE
1	B	192	ASN
1	B	272	LYS
1	A	189	PRO
1	A	180	VAL
1	B	180	VAL
1	B	80	ALA

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	266/304 (88%)	260 (98%)	6 (2%)	50	60
1	B	270/304 (89%)	263 (97%)	7 (3%)	46	55
All	All	536/608 (88%)	523 (98%)	13 (2%)	49	59

All (13) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	148	THR
1	A	149	LYS
1	A	188	GLN
1	A	243	GLU
1	A	261	GLU
1	A	285	LYS
1	B	1	ASP
1	B	33	ARG
1	B	64	GLN
1	B	188	GLN
1	B	191	ILE
1	B	215	SER
1	B	305	ARG

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	144	HIS
1	B	312	HIS

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 5.6 Ligand geometry [i](#)

3 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and

the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
2	ACT	B	401	-	1,3,3	1.63	0	0,3,3	0.00	-
2	ACT	A	401	-	1,3,3	1.60	0	0,3,3	0.00	-
2	ACT	B	402	-	1,3,3	1.36	0	0,3,3	0.00	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	B	401	ACT	1	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data ⓘ

### 6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	308/349 (88%)	0.23	24 (7%) 13 13	20, 34, 59, 76	0
1	B	309/349 (88%)	0.24	18 (5%) 23 24	17, 33, 59, 73	0
All	All	617/698 (88%)	0.23	42 (6%) 17 18	17, 34, 60, 76	0

All (42) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	315	ILE	4.4
1	B	193	ALA	4.2
1	B	38	PRO	4.0
1	B	316	PRO	3.9
1	B	317	PRO	3.8
1	A	88	ASP	3.8
1	A	149	LYS	3.7
1	A	189	PRO	3.6
1	B	90	MET	3.6
1	A	193	ALA	3.4
1	A	38	PRO	3.3
1	A	151	TRP	3.2
1	A	87	PRO	3.2
1	A	36	GLU	3.1
1	A	190	THR	3.1
1	A	191	ILE	3.1
1	B	271	GLU	3.0
1	A	50	ARG	2.9
1	A	132	VAL	2.9
1	A	314	PRO	2.9
1	B	33	ARG	2.8
1	A	127	ASN	2.8
1	A	51	GLU	2.8
1	A	33	ARG	2.8

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	A	131	ASP	2.8
1	B	188	GLN	2.6
1	B	189	PRO	2.6
1	B	81	SER	2.5
1	A	271	GLU	2.4
1	B	191	ILE	2.4
1	B	36	GLU	2.4
1	B	45	GLU	2.3
1	A	304	VAL	2.3
1	A	148	THR	2.2
1	B	198	ILE	2.2
1	A	146	LYS	2.2
1	A	89	GLU	2.2
1	B	151	TRP	2.2
1	A	147	ASN	2.2
1	B	39	LYS	2.1
1	B	47	GLU	2.1
1	A	198	ILE	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
2	ACT	B	402	4/4	0.87	0.20	42,43,44,49	0
2	ACT	B	401	4/4	0.97	0.26	37,38,40,42	0
2	ACT	A	401	4/4	0.98	0.13	42,43,44,44	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.