



# Full wwPDB X-ray Structure Validation Report ⓘ

Oct 17, 2021 – 12:34 AM EDT

PDB ID : 1R3C  
Title : THE STRUCTURE OF P38ALPHA C162S MUTANT  
Authors : Patel, S.B.; Cameron, P.M.; Frantz-Wattley, B.; O'Neill, E.; Becker, J.W.; Scapin, G.  
Deposited on : 2003-10-01  
Resolution : 2.00 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Xtriage (Phenix)	:	1.13
EDS	:	2.23.2
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac	:	5.8.0158
CCP4	:	7.0.044 (Gargrove)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.23.2


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	8085 (2.00-2.00)
Clashscore	141614	9178 (2.00-2.00)
Ramachandran outliers	138981	9054 (2.00-2.00)
Sidechain outliers	138945	9053 (2.00-2.00)
RSRZ outliers	127900	7900 (2.00-2.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	366	<div> <div>5%</div> <div> <div></div> <div>77%</div> <div>18%</div> <div>• 5%</div> </div> </div>

## 2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 3088 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Mitogen-activated protein kinase 14.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	349	Total	C	N	O	S	0	1	0
			2818	1803	481	522	12			

There are 7 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-5	GLY	-	cloning artifact	UNP Q16539
A	-4	SER	-	cloning artifact	UNP Q16539
A	-3	HIS	-	cloning artifact	UNP Q16539
A	-2	MET	-	cloning artifact	UNP Q16539
A	-1	LEU	-	cloning artifact	UNP Q16539
A	0	GLU	-	cloning artifact	UNP Q16539
A	162	SER	CYS	engineered mutation	UNP Q16539

- Molecule 2 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	2	Total	Mg	0	0
			2	2		


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	266	Total	O	0	2
			268	268		

### 3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Mitogen-activated protein kinase 14



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	45.16Å 84.80Å 124.91Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	17.00 – 2.00 36.60 – 1.79	Depositor EDS
% Data completeness (in resolution range)	97.9 (17.00-2.00) 90.9 (36.60-1.79)	Depositor EDS
$R_{merge}$	0.07	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.77 (at 1.79Å)	Xtriage
Refinement program	CNS	Depositor
R, $R_{free}$	0.200 , 0.224 0.202 , 0.227	Depositor DCC
$R_{free}$ test set	2091 reflections (5.01%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	24.6	Xtriage
Anisotropy	0.178	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.37 , 48.8	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.47$ , $\langle L^2 \rangle = 0.30$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.95	EDS
Total number of atoms	3088	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	29.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.99% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality

### 5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: MG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.47	0/2884	0.62	0/3918

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2818	0	2796	60	0
2	A	2	0	0	0	0
3	A	268	0	0	9	0
All	All	3088	0	2796	60	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 11.

All (60) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:201:ASN:OD1	1:A:203:THR:HG22	1.68	0.93

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:203:THR:HG21	1:A:293:SER:HB2	1.54	0.90
1:A:237:ARG:HD3	1:A:265:MET:HE1	1.55	0.87
1:A:245:GLU:HA	1:A:248:LYS:HE2	1.70	0.74
1:A:203:THR:CG2	1:A:293:SER:HB2	2.19	0.72
1:A:237:ARG:HD3	1:A:265:MET:CE	2.20	0.71
1:A:23:ARG:CZ	1:A:45:LYS:HD3	2.22	0.69
1:A:11:GLN:HG2	1:A:13:LEU:HD13	1.75	0.69
1:A:13:LEU:HD23	1:A:39:CYS:SG	2.39	0.62
1:A:201:ASN:OD1	1:A:203:THR:CG2	2.46	0.61
1:A:109:MET:HB3	1:A:157:ALA:HB1	1.84	0.59
1:A:57:ARG:HH11	1:A:57:ARG:HB3	1.68	0.59
1:A:253:GLU:HG3	1:A:256:ARG:HH21	1.68	0.58
1:A:57:ARG:HB3	1:A:60:GLN:HB3	1.85	0.58
1:A:189:ARG:HD2	3:A:1054:HOH:O	2.03	0.57
1:A:253:GLU:HG3	1:A:256:ARG:NH2	2.20	0.55
1:A:12:GLU:O	1:A:13:LEU:HD12	2.10	0.52
1:A:57:ARG:NH1	1:A:60:GLN:HB3	2.25	0.51
1:A:117:VAL:HB	1:A:122:LEU:HD11	1.93	0.51
1:A:160:GLU:HB2	3:A:1229:HOH:O	2.10	0.51
1:A:245:GLU:HA	1:A:248:LYS:CE	2.39	0.51
1:A:330[A]:ARG:HD3	3:A:1001[A]:HOH:O	2.09	0.51
1:A:113:LEU:O	1:A:117:VAL:HG22	2.11	0.51
1:A:184:ALA:HA	1:A:187:TRP:CE3	2.46	0.51
1:A:54:LYS:HE3	3:A:1150:HOH:O	2.11	0.50
1:A:237:ARG:HA	1:A:265:MET:CE	2.41	0.50
1:A:244:ALA:O	1:A:248:LYS:HG3	2.12	0.50
1:A:122:LEU:HD12	1:A:122:LEU:N	2.26	0.50
1:A:283:ASP:O	1:A:287:LYS:HG2	2.12	0.49
1:A:57:ARG:HH11	1:A:60:GLN:HB3	1.76	0.49
1:A:159:ASN:ND2	1:A:163:GLU:HB2	2.28	0.49
1:A:127:VAL:HG21	1:A:217:LEU:HD23	1.95	0.48
1:A:237:ARG:HA	1:A:265:MET:HE2	1.95	0.48
1:A:175:THR:HB	1:A:178:GLU:HG3	1.96	0.47
1:A:57:ARG:HH11	1:A:57:ARG:CB	2.27	0.47
1:A:61:SER:HA	1:A:334:ILE:HD11	1.96	0.46
1:A:330[B]:ARG:NH2	3:A:1153:HOH:O	2.48	0.46
1:A:118:LYS:O	1:A:120:GLN:HG2	2.15	0.46
1:A:256:ARG:HG2	1:A:260:GLN:HE21	1.81	0.45
1:A:140:TYR:CE1	1:A:320:ALA:HB2	2.53	0.44
1:A:159:ASN:HD21	1:A:163:GLU:HB2	1.83	0.44
1:A:109:MET:HA	1:A:109:MET:CE	2.48	0.43

*Continued on next page...*

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:158:VAL:CG1	1:A:162:SER:HA	2.48	0.43
1:A:71:GLU:HB2	1:A:171:LEU:HD13	1.99	0.43
1:A:350:PRO:HG3	3:A:1074:HOH:O	2.18	0.43
1:A:79:LYS:HE2	3:A:1103:HOH:O	2.17	0.43
1:A:250:ILE:HD13	1:A:259:ILE:CD1	2.49	0.43
1:A:189:ARG:HH11	1:A:189:ARG:HB3	1.83	0.43
1:A:179:MET:HG3	1:A:186:ARG:HG3	2.01	0.43
1:A:237:ARG:HG2	3:A:1187:HOH:O	2.19	0.43
1:A:10:ARG:NH2	1:A:17:ILE:HD13	2.34	0.42
1:A:203:THR:HG21	1:A:293:SER:CB	2.39	0.42
1:A:60:GLN:HE21	1:A:64:HIS:CD2	2.38	0.41
1:A:117:VAL:O	1:A:120:GLN:HB2	2.19	0.41
1:A:78:MET:HE2	1:A:146:ILE:HD13	2.01	0.41
1:A:55:LEU:CD1	1:A:68:THR:HG23	2.50	0.41
1:A:202:GLN:HG2	3:A:1149:HOH:O	2.21	0.41
1:A:23:ARG:NH2	1:A:45:LYS:HD3	2.36	0.41
1:A:181:GLY:HA2	1:A:226:THR:O	2.21	0.41
1:A:28:SER:O	1:A:30:VAL:HG13	2.22	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	348/366 (95%)	339 (97%)	7 (2%)	2 (1%)	25 19

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	115	ASN
1	A	15	LYS



### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	308/325 (95%)	302 (98%)	6 (2%)	57 61

All (6) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	4	GLU
1	A	57	ARG
1	A	189	ARG
1	A	237	ARG
1	A	330[A]	ARG
1	A	330[B]	ARG

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (5) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	14	ASN
1	A	60	GLN
1	A	114	ASN
1	A	115	ASN
1	A	260	GLN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 5.6 Ligand geometry

Of 2 ligands modelled in this entry, 2 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

## 5.7 Other polymers

There are no such residues in this entry.

## 5.8 Polymer linkage issues

There are no chain breaks in this entry.

## 6 Fit of model and data ⓘ

### 6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	349/366 (95%)	-0.00	18 (5%) 27 26	15, 26, 49, 57	0

All (18) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	110	GLY	5.3
1	A	172	ALA	5.3
1	A	34	ALA	5.1
1	A	35	TYR	4.6
1	A	176	ASP	3.9
1	A	199	HIS	3.7
1	A	175	THR	3.4
1	A	13	LEU	3.2
1	A	33	GLY	3.0
1	A	57	ARG	2.9
1	A	160	GLU	2.9
1	A	174	HIS	2.8
1	A	173	ARG	2.6
1	A	4	GLU	2.5
1	A	109	MET	2.5
1	A	177	ASP	2.2
1	A	119	CYS	2.1
1	A	14	ASN	2.1

### 6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

### 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
2	MG	A	2002	1/1	0.76	0.11	43,43,43,43	1
2	MG	A	2001	1/1	0.98	0.05	41,41,41,41	0

### 6.5 Other polymers [i](#)

There are no such residues in this entry.