



# Full wwPDB X-ray Structure Validation Report ⓘ

May 29, 2020 – 03:51 pm BST

PDB ID : 2R3B  
Title : CRYSTAL STRUCTURE OF a ribokinase-like superfamily protein (EF1790)  
FROM ENTEROCOCCUS FAECALIS V583 AT 1.80 Å RESOLUTION  
Authors : Joint Center for Structural Genomics (JCSG)  
Deposited on : 2007-08-29  
Resolution : 1.80 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.11  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.11


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 1.80 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	5950 (1.80-1.80)
Clashscore	141614	6793 (1.80-1.80)
Ramachandran outliers	138981	6697 (1.80-1.80)
Sidechain outliers	138945	6696 (1.80-1.80)
RSRZ outliers	127900	5850 (1.80-1.80)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	310	<div> <div>3%</div> <div> <div></div> <div>83%</div> <div>9%</div> <div>9%</div> </div> </div>
1	B	310	<div> <div>2%</div> <div> <div></div> <div>84%</div> <div>8%</div> <div>8%</div> </div> </div>

## 2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 5182 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called YjeF-related protein.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	283	Total	C	N	O	S	Se	0	17	0
			2258	1445	386	415	2	10			
1	B	284	Total	C	N	O	S	Se	0	17	0
			2260	1448	382	418	2	10			

There are 38 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-18	MSE	-	LEADER SEQUENCE	UNP Q833Y3
A	-17	GLY	-	LEADER SEQUENCE	UNP Q833Y3
A	-16	SER	-	LEADER SEQUENCE	UNP Q833Y3
A	-15	ASP	-	LEADER SEQUENCE	UNP Q833Y3
A	-14	LYS	-	LEADER SEQUENCE	UNP Q833Y3
A	-13	ILE	-	LEADER SEQUENCE	UNP Q833Y3
A	-12	HIS	-	LEADER SEQUENCE	UNP Q833Y3
A	-11	HIS	-	LEADER SEQUENCE	UNP Q833Y3
A	-10	HIS	-	LEADER SEQUENCE	UNP Q833Y3
A	-9	HIS	-	LEADER SEQUENCE	UNP Q833Y3
A	-8	HIS	-	LEADER SEQUENCE	UNP Q833Y3
A	-7	HIS	-	LEADER SEQUENCE	UNP Q833Y3
A	-6	GLU	-	LEADER SEQUENCE	UNP Q833Y3
A	-5	ASN	-	LEADER SEQUENCE	UNP Q833Y3
A	-4	LEU	-	LEADER SEQUENCE	UNP Q833Y3
A	-3	TYR	-	LEADER SEQUENCE	UNP Q833Y3
A	-2	PHE	-	LEADER SEQUENCE	UNP Q833Y3
A	-1	GLN	-	LEADER SEQUENCE	UNP Q833Y3
A	0	GLY	-	LEADER SEQUENCE	UNP Q833Y3
B	-18	MSE	-	LEADER SEQUENCE	UNP Q833Y3
B	-17	GLY	-	LEADER SEQUENCE	UNP Q833Y3
B	-16	SER	-	LEADER SEQUENCE	UNP Q833Y3
B	-15	ASP	-	LEADER SEQUENCE	UNP Q833Y3
B	-14	LYS	-	LEADER SEQUENCE	UNP Q833Y3
B	-13	ILE	-	LEADER SEQUENCE	UNP Q833Y3

*Continued on next page...*

*Continued from previous page...*

Chain	Residue	Modelled	Actual	Comment	Reference
B	-12	HIS	-	LEADER SEQUENCE	UNP Q833Y3
B	-11	HIS	-	LEADER SEQUENCE	UNP Q833Y3
B	-10	HIS	-	LEADER SEQUENCE	UNP Q833Y3
B	-9	HIS	-	LEADER SEQUENCE	UNP Q833Y3
B	-8	HIS	-	LEADER SEQUENCE	UNP Q833Y3
B	-7	HIS	-	LEADER SEQUENCE	UNP Q833Y3
B	-6	GLU	-	LEADER SEQUENCE	UNP Q833Y3
B	-5	ASN	-	LEADER SEQUENCE	UNP Q833Y3
B	-4	LEU	-	LEADER SEQUENCE	UNP Q833Y3
B	-3	TYR	-	LEADER SEQUENCE	UNP Q833Y3
B	-2	PHE	-	LEADER SEQUENCE	UNP Q833Y3
B	-1	GLN	-	LEADER SEQUENCE	UNP Q833Y3
B	0	GLY	-	LEADER SEQUENCE	UNP Q833Y3

- Molecule 2 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	B	1	Total Mg 1 1	0	0
2	A	1	Total Mg 1 1	0	0

- Molecule 3 is CHLORIDE ION (three-letter code: CL) (formula: Cl).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	B	2	Total Cl 2 2	0	0
3	A	1	Total Cl 1 1	0	0

- Molecule 4 is 1,2-ETHANEDIOL (three-letter code: EDO) (formula: C<sub>2</sub>H<sub>6</sub>O<sub>2</sub>).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	1	Total C O 4 2 2	0	0
4	A	1	Total C O 4 2 2	0	0
4	A	1	Total C O 4 2 2	0	0
4	B	1	Total C O 4 2 2	0	0
4	B	1	Total C O 4 2 2	0	0
4	B	1	Total C O 4 2 2	0	0
4	B	1	Total C O 4 2 2	0	0
4	B	1	Total C O 4 2 2	0	0
4	B	1	Total C O 4 2 2	0	0

- Molecule 5 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	A	323	Total O 324 324	0	1
5	B	298	Total O 299 299	0	1



## 4 Data and refinement statistics

Property	Value	Source
Space group	C 2 2 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	97.82Å 159.56Å 102.50Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	29.59 – 1.80 29.60 – 1.80	Depositor EDS
% Data completeness (in resolution range)	99.3 (29.59-1.80) 99.3 (29.60-1.80)	Depositor EDS
$R_{merge}$	0.10	Depositor
$R_{sym}$	0.10	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.69 (at 1.80Å)	Xtriage
Refinement program	REFMAC 5.2.0019, PHENIX	Depositor
R, $R_{free}$	0.150 , 0.183 0.162 , 0.191	Depositor DCC
$R_{free}$ test set	3709 reflections (5.03%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	16.6	Xtriage
Anisotropy	0.563	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.34 , 60.6	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	0.010 for 1/2*h-1/2*k,-3/2*h-1/2*k,-l 0.018 for 1/2*h+1/2*k,3/2*h-1/2*k,-l	Xtriage
$F_o, F_c$ correlation	0.96	EDS
Total number of atoms	5182	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	25.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.73% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality

### 5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: MG, EDO, CL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.77	4/2338 (0.2%)	0.71	0/3156
1	B	0.72	2/2351 (0.1%)	0.70	1/3176 (0.0%)
All	All	0.74	6/4689 (0.1%)	0.71	1/6332 (0.0%)

All (6) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	81[A]	GLU	CD-OE1	-10.44	1.14	1.25
1	A	81[B]	GLU	CD-OE1	-10.44	1.14	1.25
1	B	81[A]	GLU	CD-OE1	-7.57	1.17	1.25
1	B	81[B]	GLU	CD-OE1	-7.57	1.17	1.25
1	A	81[A]	GLU	CB-CG	5.10	1.61	1.52
1	A	81[B]	GLU	CB-CG	5.10	1.61	1.52

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	7	ASP	CB-CG-OD2	-5.00	113.80	118.30

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.



Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2258	0	2330	29	0
1	B	2260	0	2323	16	0
2	A	1	0	0	0	0
2	B	1	0	0	0	0
3	A	1	0	0	0	0
3	B	2	0	0	1	0
4	A	12	0	18	0	0
4	B	24	0	36	0	0
5	A	324	0	0	4	0
5	B	299	0	0	4	0
All	All	5182	0	4707	45	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (45) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:87[B]:THR:HG21	5:A:543:HOH:O	1.61	1.00
1:A:200:PHE:HE2	1:A:233[B]:ILE:HG22	1.59	0.67
1:A:175[B]:ARG:NH2	1:A:176:GLN:HE21	1.96	0.62
1:A:15:GLN:OE1	1:A:276:PRO:HA	2.00	0.62
1:A:2[B]:ARG:HD3	1:A:200:PHE:CE1	2.35	0.62
1:A:110[B]:GLN:HG2	1:A:114[B]:MSE:HE2	1.82	0.62
1:B:155[A]:MSE:SE	1:B:165:ILE:HD11	2.50	0.61
1:A:200:PHE:CE2	1:A:233[B]:ILE:HG22	2.35	0.61
1:B:30[B]:LEU:CD1	1:B:56:THR:HG23	2.31	0.61
1:A:200:PHE:HE2	1:A:233[B]:ILE:CG2	2.14	0.60
1:A:30[B]:LEU:CD1	1:A:56:THR:HG23	2.31	0.60
1:A:175[B]:ARG:NH2	1:A:176:GLN:NE2	2.50	0.59
1:B:118[A]:GLN:HG2	5:B:480:HOH:O	2.02	0.58
1:B:200:PHE:CE2	1:B:233[B]:ILE:HG22	2.38	0.58
1:B:1[B]:MSE:HE2	1:B:170:LEU:HD11	1.86	0.57
1:B:118[B]:GLN:NE2	5:B:368:HOH:O	2.36	0.57
1:B:45:THR:HG21	1:B:68[B]:LEU:HD11	1.90	0.53
1:A:83:THR:O	1:A:87[B]:THR:HG23	2.10	0.50
1:A:5:SER:O	1:A:8:ILE:HG22	2.12	0.50
1:A:88[B]:ASN:OD1	1:A:92:GLN:NE2	2.44	0.50
1:B:200:PHE:HE2	1:B:233[B]:ILE:HG22	1.78	0.48
1:B:147:LYS:HB2	5:B:478:HOH:O	2.12	0.48
1:B:5:SER:O	1:B:8:ILE:HG22	2.14	0.47
1:A:2[B]:ARG:NH1	5:A:548:HOH:O	2.48	0.47

*Continued on next page...*

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:154:GLN:HB3	1:A:165:ILE:HD12	1.96	0.46
1:B:1[B]:MSE:SE	1:B:201:GLN:HB2	2.65	0.45
1:A:200:PHE:CE2	1:A:233[B]:ILE:CG2	2.96	0.45
1:A:175[B]:ARG:CZ	1:A:176:GLN:HE21	2.30	0.44
1:A:30[A]:LEU:HD22	1:A:97:LEU:HD23	2.00	0.43
1:B:30[B]:LEU:HD11	1:B:56:THR:HG23	1.97	0.43
1:A:30[B]:LEU:CD1	1:A:56:THR:CG2	2.97	0.43
1:A:30[B]:LEU:HD11	1:A:56:THR:HG23	1.99	0.43
1:A:1[B]:MSE:SE	5:A:571:HOH:O	2.87	0.43
1:A:155[B]:MSE:HE1	5:A:556:HOH:O	2.18	0.42
1:A:59:ILE:HD13	1:A:86[A]:LEU:HG	2.00	0.42
1:B:13:ILE:HG22	1:B:274:ALA:HA	2.01	0.42
1:A:114[A]:MSE:HB3	1:A:114[A]:MSE:HE3	1.94	0.42
1:B:39:GLY:N	3:B:294:CL:CL	2.90	0.42
1:A:2[B]:ARG:HD3	1:A:200:PHE:CD1	2.55	0.41
1:B:15:GLN:CD	1:B:276:PRO:HA	2.42	0.41
1:B:36:GLN:NE2	5:B:336:HOH:O	2.50	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles

### 5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	298/310 (96%)	290 (97%)	8 (3%)	0	100	100
1	B	300/310 (97%)	294 (98%)	6 (2%)	0	100	100
All	All	598/620 (96%)	584 (98%)	14 (2%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	248 / 251 (99%)	247 (100%)	1 (0%)	91	89
1	B	249 / 251 (99%)	247 (99%)	2 (1%)	81	78
All	All	497 / 502 (99%)	494 (99%)	3 (1%)	86	84

All (3) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	166	GLU
1	B	86	LEU
1	B	166	GLU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	92	GLN
1	A	158	GLN

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

## 5.6 Ligand geometry

Of 14 ligands modelled in this entry, 5 are monoatomic - leaving 9 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
4	EDO	B	300	-	3,3,3	0.50	0	2,2,2	0.05	0
4	EDO	B	297	-	3,3,3	0.50	0	2,2,2	0.10	0
4	EDO	A	295	-	3,3,3	0.45	0	2,2,2	0.31	0
4	EDO	A	296	-	3,3,3	0.30	0	2,2,2	0.87	0
4	EDO	B	296	-	3,3,3	0.56	0	2,2,2	0.35	0
4	EDO	B	299	-	3,3,3	0.49	0	2,2,2	0.55	0
4	EDO	B	298	-	3,3,3	0.45	0	2,2,2	0.07	0
4	EDO	B	295	-	3,3,3	0.45	0	2,2,2	0.42	0
4	EDO	A	294	-	3,3,3	0.49	0	2,2,2	0.15	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	EDO	B	300	-	-	0/1/1/1	-
4	EDO	B	297	-	-	0/1/1/1	-
4	EDO	A	295	-	-	1/1/1/1	-
4	EDO	A	296	-	-	1/1/1/1	-
4	EDO	B	296	-	-	0/1/1/1	-
4	EDO	B	299	-	-	1/1/1/1	-
4	EDO	B	298	-	-	1/1/1/1	-
4	EDO	B	295	-	-	1/1/1/1	-
4	EDO	A	294	-	-	1/1/1/1	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (6) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	B	298	EDO	O1-C1-C2-O2
4	A	295	EDO	O1-C1-C2-O2
4	A	294	EDO	O1-C1-C2-O2
4	A	296	EDO	O1-C1-C2-O2
4	B	295	EDO	O1-C1-C2-O2
4	B	299	EDO	O1-C1-C2-O2

There are no ring outliers.

No monomer is involved in short contacts.

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2			OWAB(Å <sup>2</sup> )	Q<0.9
1	A	276/310 (89%)	-0.24	9 (3%)	46	40	12, 19, 41, 67	0
1	B	277/310 (89%)	-0.35	6 (2%)	62	57	12, 18, 40, 67	0
All	All	553/620 (89%)	-0.30	15 (2%)	54	49	12, 19, 41, 67	0

All (15) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	165	ILE	5.3
1	B	276	PRO	5.0
1	A	276	PRO	4.8
1	A	166	GLU	4.0
1	B	139	ASN	3.4
1	B	275	GLN	3.1
1	A	21	TYR	3.0
1	A	-6	GLU	2.7
1	B	166	GLU	2.7
1	B	-7	HIS	2.6
1	A	167	GLN	2.2
1	A	-2	PHE	2.1
1	B	167[A]	GLN	2.1
1	A	0[A]	GLY	2.1
1	A	275	GLN	2.1

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
4	EDO	B	297	4/4	0.76	0.33	67,68,68,69	0
4	EDO	B	295	4/4	0.86	0.19	51,51,52,57	0
3	CL	B	294	1/1	0.87	0.21	51,51,51,51	0
4	EDO	A	295	4/4	0.90	0.12	52,53,54,60	0
4	EDO	B	299	4/4	0.90	0.12	29,34,37,39	0
4	EDO	B	296	4/4	0.91	0.14	31,38,39,40	0
2	MG	A	292	1/1	0.91	0.17	51,51,51,51	0
3	CL	B	293	1/1	0.91	0.09	41,41,41,41	0
4	EDO	B	298	4/4	0.92	0.27	59,60,60,61	0
4	EDO	B	300	4/4	0.93	0.18	44,51,53,54	0
4	EDO	A	296	4/4	0.95	0.13	32,33,36,49	0
4	EDO	A	294	4/4	0.96	0.12	23,32,41,53	0
2	MG	B	292	1/1	0.98	0.17	40,40,40,40	0
3	CL	A	293	1/1	0.99	0.03	27,27,27,27	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.