



# Full wwPDB X-ray Structure Validation Report ⓘ

Jun 15, 2020 – 06:17 am BST

PDB ID : 4R30  
Title : Structure of human laforin dual specificity phosphatase domain  
Authors : Sankhala, R.S.; Koksai, A.C.; Cingolani, G.  
Deposited on : 2014-08-13  
Resolution : 2.30 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.11  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.11


# 1 Overall quality at a glance ⓘ

The following experimental techniques were used to determine the structure:

## *X-RAY DIFFRACTION*

The reported resolution of this entry is 2.30 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	5042 (2.30-2.30)
Clashscore	141614	5643 (2.30-2.30)
Ramachandran outliers	138981	5575 (2.30-2.30)
Sidechain outliers	138945	5575 (2.30-2.30)
RSRZ outliers	127900	4938 (2.30-2.30)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	184	<div> <div>5%</div> <div> <div></div> <div>90%</div> <div>6%</div> <div>••</div> </div> </div>
1	B	184	<div> <div>3%</div> <div> <div></div> <div>90%</div> <div>7%</div> <div>••</div> </div> </div>
1	C	184	<div> <div>4%</div> <div> <div></div> <div>86%</div> <div>11%</div> <div>•</div> </div> </div>
1	D	184	<div> <div>5%</div> <div> <div></div> <div>86%</div> <div>11%</div> <div>••</div> </div> </div>

## 2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 6051 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Laforin.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	179	Total	C	N	O	S	0	0	0
			1435	918	250	253	14			
1	B	179	Total	C	N	O	S	0	0	0
			1435	918	250	253	14			
1	C	179	Total	C	N	O	S	0	0	0
			1435	918	250	253	14			
1	D	179	Total	C	N	O	S	0	0	0
			1435	918	250	253	14			

There are 4 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	266	SER	CYS	ENGINEERED MUTATION	UNP O95278
B	266	SER	CYS	ENGINEERED MUTATION	UNP O95278
C	266	SER	CYS	ENGINEERED MUTATION	UNP O95278
D	266	SER	CYS	ENGINEERED MUTATION	UNP O95278

- Molecule 2 is SULFATE ION (three-letter code: SO4) (formula: O<sub>4</sub>S).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	O	S	0	0
			5	4	1		
2	B	1	Total	O	S	0	0
			5	4	1		
2	C	1	Total	O	S	0	0
			5	4	1		
2	D	1	Total	O	S	0	0
			5	4	1		

- Molecule 3 is BETA-MERCAPTOETHANOL (three-letter code: BME) (formula: C<sub>2</sub>H<sub>6</sub>OS).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	1	Total C O S 4 2 1 1	0	0
3	A	1	Total C O S 4 2 1 1	0	0
3	A	1	Total C O S 4 2 1 1	0	0
3	B	1	Total C O S 4 2 1 1	0	0
3	B	1	Total C O S 4 2 1 1	0	0
3	B	1	Total C O S 4 2 1 1	0	0
3	C	1	Total C O S 4 2 1 1	0	0
3	C	1	Total C O S 4 2 1 1	0	0
3	C	1	Total C O S 4 2 1 1	0	0
3	D	1	Total C O S 4 2 1 1	0	0
3	D	1	Total C O S 4 2 1 1	0	0
3	D	1	Total C O S 4 2 1 1	0	0



- Molecule 4 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	66	Total O 66 66	0	0
4	B	72	Total O 72 72	0	0
4	C	48	Total O 48 48	0	0
4	D	57	Total O 57 57	0	0



### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.



#### • Molecule 1: Laforin





#### • Molecule 1: Laforin



#### • Molecule 1: Laforin



#### • Molecule 1: Laforin



## 4 Data and refinement statistics

Property	Value	Source
Space group	I 41	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	123.97Å 123.97Å 160.77Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	14.88 – 2.30 14.87 – 2.30	Depositor EDS
% Data completeness (in resolution range)	96.9 (14.88-2.30) 71.7 (14.87-2.30)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	0.80 (at 2.29Å)	Xtriage
Refinement program	PHENIX (phenix.refine: 1.8.2_1309)	Depositor
R, $R_{free}$	0.188 , 0.219 0.200 , 0.229	Depositor DCC
$R_{free}$ test set	1689 reflections (3.80%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	34.6	Xtriage
Anisotropy	0.385	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.33 , 51.8	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.54$ , $\langle L^2 \rangle = 0.38$	Xtriage
Estimated twinning fraction	0.054 for -h,k,-l	Xtriage
$F_o, F_c$ correlation	0.96	EDS
Total number of atoms	6051	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	59.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 46.79 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 1.0890e-04. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: SO4, BME

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.42	0/1472	0.57	0/1996
1	B	0.41	0/1472	0.55	0/1996
1	C	0.40	0/1472	0.53	0/1996
1	D	0.41	0/1472	0.55	0/1996
All	All	0.41	0/5888	0.55	0/7984

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1435	0	1424	9	0
1	B	1435	0	1424	7	0
1	C	1435	0	1424	13	0
1	D	1435	0	1424	12	0
2	A	5	0	0	0	0
2	B	5	0	0	0	0
2	C	5	0	0	0	0
2	D	5	0	0	0	0
3	A	12	0	15	1	0

*Continued on next page...*



*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	B	12	0	15	2	0
3	C	12	0	14	3	0
3	D	12	0	15	5	0
4	A	66	0	0	1	0
4	B	72	0	0	1	0
4	C	48	0	0	1	0
4	D	57	0	0	2	0
All	All	6051	0	5755	41	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (41) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:250:CYS:SG	3:C:502:BME:O1	2.09	1.10
1:B:300:ARG:HB3	1:B:303:VAL:HB	1.64	0.79
1:A:274:THR:HG22	3:A:503:BME:H11	1.68	0.75
1:D:253:HIS:HD2	3:D:502:BME:H22	1.54	0.73
1:D:253:HIS:CD2	3:D:502:BME:H22	2.23	0.73
1:D:329:CYS:SG	3:D:504:BME:O1	2.52	0.67
1:C:300:ARG:HB3	1:C:303:VAL:HB	1.77	0.65
1:A:331:LEU:HB3	1:C:331:LEU:HB3	1.82	0.61
1:A:290:ARG:NH1	4:A:664:HOH:O	2.29	0.60
1:D:325:ARG:NH1	4:D:628:HOH:O	2.35	0.60
1:B:161:LEU:HD13	3:B:504:BME:H21	1.85	0.58
1:C:274:THR:HG22	3:C:503:BME:H11	1.86	0.57
1:D:157:TYR:OH	1:D:179:LYS:NZ	2.38	0.57
1:C:329:CYS:SG	3:C:504:BME:O1	2.57	0.55
1:D:171:ARG:NH2	4:D:610:HOH:O	2.22	0.54
1:D:275:ALA:HA	3:D:503:BME:H12	1.88	0.54
1:B:155:MET:HE1	1:B:180:LEU:HD13	1.91	0.52
1:D:241:ARG:NH1	1:D:304:TYR:O	2.37	0.50
1:A:300:ARG:HB3	1:A:303:VAL:HB	1.95	0.49
1:B:274:THR:HG23	1:B:296:LEU:HD13	1.95	0.49
1:A:155:MET:HE1	1:A:180:LEU:HD13	1.94	0.48
1:B:318:PHE:CD1	1:B:323:LYS:HE3	2.49	0.47
1:B:323:LYS:NZ	4:B:661:HOH:O	2.25	0.46
3:B:502:BME:H21	3:D:502:BME:O1	2.16	0.46
1:C:299:LYS:O	4:C:638:HOH:O	2.21	0.45
1:C:197:ASP:CG	1:C:267:ASN:HD21	2.20	0.45

*Continued on next page...*

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:299:LYS:H	1:A:299:LYS:HD2	1.83	0.44
1:A:287:TRP:CE2	1:A:291:LYS:HE2	2.53	0.43
1:C:155:MET:HE1	1:C:180:LEU:HD13	2.01	0.43
1:C:215:ASP:HA	1:C:218:ILE:HD12	2.00	0.43
1:D:230:TRP:CE2	1:D:232:PRO:HG3	2.55	0.42
1:C:283:TYR:CE1	1:C:314:GLN:HB2	2.55	0.42
1:C:291:LYS:NZ	1:C:331:LEU:HD21	2.34	0.42
1:C:331:LEU:HD23	1:C:331:LEU:O	2.20	0.41
1:A:274:THR:HG23	1:A:296:LEU:HD13	2.03	0.41
1:A:291:LYS:NZ	1:A:331:LEU:HD21	2.36	0.41
1:D:300:ARG:HB3	1:D:303:VAL:HB	2.02	0.41
1:D:163:ASN:ND2	1:D:256:LEU:HD22	2.37	0.40
1:C:195:GLU:HG3	1:C:214:PRO:CG	2.51	0.40
1:B:283:TYR:CE1	1:B:314:GLN:HB2	2.56	0.40
1:D:197:ASP:CG	1:D:267:ASN:HD21	2.24	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles

### 5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	177/184 (96%)	172 (97%)	5 (3%)	0	100	100
1	B	177/184 (96%)	172 (97%)	5 (3%)	0	100	100
1	C	177/184 (96%)	171 (97%)	6 (3%)	0	100	100
1	D	177/184 (96%)	171 (97%)	6 (3%)	0	100	100
All	All	708/736 (96%)	686 (97%)	22 (3%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	155/158 (98%)	151 (97%)	4 (3%)	46	63
1	B	155/158 (98%)	151 (97%)	4 (3%)	46	63
1	C	155/158 (98%)	153 (99%)	2 (1%)	69	82
1	D	155/158 (98%)	150 (97%)	5 (3%)	39	54
All	All	620/632 (98%)	605 (98%)	15 (2%)	49	66

All (15) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	180	LEU
1	A	329	CYS
1	A	330	SER
1	A	331	LEU
1	B	180	LEU
1	B	243	GLN
1	B	329	CYS
1	B	330	SER
1	C	296	LEU
1	C	330	SER
1	D	180	LEU
1	D	243	GLN
1	D	299	LYS
1	D	329	CYS
1	D	330	SER

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	243	GLN
1	C	243	GLN

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

## 5.6 Ligand geometry ⓘ

16 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
3	BME	D	502	1	3,3,3	1.28	0	1,2,2	2.06	1 (100%)
3	BME	C	504	1	3,3,3	0.90	0	1,2,2	2.10	1 (100%)
3	BME	A	504	1	3,3,3	0.65	0	1,2,2	1.49	0
3	BME	A	503	1	3,3,3	1.07	0	1,2,2	0.83	0
2	SO4	B	501	-	4,4,4	0.16	0	6,6,6	0.29	0
2	SO4	A	501	-	4,4,4	0.16	0	6,6,6	0.23	0
2	SO4	D	501	-	4,4,4	0.15	0	6,6,6	0.29	0
3	BME	A	502	1	3,3,3	2.06	1 (33%)	1,2,2	1.49	0
3	BME	D	504	1	3,3,3	0.70	0	1,2,2	0.28	0
3	BME	C	502	1	3,3,3	0.81	0	1,2,2	2.06	1 (100%)
3	BME	B	504	1	3,3,3	0.77	0	1,2,2	1.85	0
2	SO4	C	501	-	4,4,4	0.12	0	6,6,6	0.40	0
3	BME	C	503	1	3,3,3	0.94	0	1,2,2	1.24	0
3	BME	B	503	1	3,3,3	0.99	0	1,2,2	1.36	0
3	BME	D	503	1	3,3,3	0.84	0	1,2,2	1.47	0
3	BME	B	502	1	3,3,3	1.04	0	1,2,2	0.67	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	BME	D	502	1	-	1/1/1/1	-
3	BME	C	504	1	-	1/1/1/1	-
3	BME	A	504	1	-	1/1/1/1	-
3	BME	A	503	1	-	1/1/1/1	-
3	BME	A	502	1	-	0/1/1/1	-
3	BME	D	504	1	-	1/1/1/1	-
3	BME	C	502	1	-	1/1/1/1	-
3	BME	B	504	1	-	1/1/1/1	-
3	BME	C	503	1	-	1/1/1/1	-
3	BME	B	503	1	-	1/1/1/1	-
3	BME	D	503	1	-	1/1/1/1	-
3	BME	B	502	1	-	1/1/1/1	-

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	502	BME	C2-S2	-3.50	1.68	1.80

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	C	504	BME	O1-C1-C2	-2.10	102.55	110.83
3	D	502	BME	O1-C1-C2	-2.06	102.70	110.83
3	C	502	BME	O1-C1-C2	-2.06	102.72	110.83

There are no chirality outliers.

All (11) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	D	502	BME	O1-C1-C2-S2
3	B	502	BME	O1-C1-C2-S2
3	C	504	BME	O1-C1-C2-S2
3	A	504	BME	O1-C1-C2-S2
3	A	503	BME	O1-C1-C2-S2
3	D	504	BME	O1-C1-C2-S2
3	C	502	BME	O1-C1-C2-S2
3	B	504	BME	O1-C1-C2-S2
3	C	503	BME	O1-C1-C2-S2

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms
3	B	503	BME	O1-C1-C2-S2
3	D	503	BME	O1-C1-C2-S2

There are no ring outliers.

9 monomers are involved in 10 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	D	502	BME	3	0
3	C	504	BME	1	0
3	A	503	BME	1	0
3	D	504	BME	1	0
3	C	502	BME	1	0
3	B	504	BME	1	0
3	C	503	BME	1	0
3	D	503	BME	1	0
3	B	502	BME	1	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	179/184 (97%)	-0.16	10 (5%) 24 30	35, 56, 102, 125	0
1	B	179/184 (97%)	-0.21	6 (3%) 45 52	33, 53, 94, 122	0
1	C	179/184 (97%)	-0.04	8 (4%) 33 40	38, 59, 99, 126	0
1	D	179/184 (97%)	-0.26	9 (5%) 28 35	33, 52, 101, 133	0
All	All	716/736 (97%)	-0.17	33 (4%) 32 39	33, 55, 99, 133	0

All (33) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	C	300	ARG	4.6
1	D	301	PRO	4.2
1	C	330	SER	4.0
1	A	300	ARG	3.9
1	A	209	PRO	3.9
1	B	302	ALA	3.7
1	B	209	PRO	3.6
1	D	330	SER	3.6
1	D	196	TRP	3.6
1	A	330	SER	3.4
1	D	302	ALA	3.3
1	A	301	PRO	3.3
1	C	196	TRP	3.2
1	D	331	LEU	3.1
1	D	154	ALA	3.1
1	D	298	ALA	3.1
1	B	300	ARG	3.0
1	C	153	GLN	2.9
1	C	301	PRO	2.9
1	C	299	LYS	2.8
1	B	153	GLN	2.7

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	D	300	ARG	2.7
1	A	299	LYS	2.5
1	A	153	GLN	2.5
1	A	235	ASP	2.5
1	B	330	SER	2.4
1	C	298	ALA	2.3
1	D	235	ASP	2.3
1	A	211	PRO	2.2
1	A	298	ALA	2.2
1	C	235	ASP	2.1
1	A	331	LEU	2.0
1	B	210	GLU	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates ⓘ

There are no carbohydrates in this entry.

## 6.4 Ligands ⓘ

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
3	BME	B	504	4/4	0.91	0.18	41,44,45,117	0
3	BME	C	504	4/4	0.92	0.20	47,49,52,111	0
3	BME	A	504	4/4	0.94	0.14	49,51,56,138	0
3	BME	D	504	4/4	0.96	0.10	44,52,62,120	0
3	BME	A	502	4/4	0.97	0.12	12,40,51,58	0
3	BME	B	503	4/4	0.97	0.15	39,54,65,65	0
3	BME	D	502	4/4	0.98	0.14	34,58,63,72	0
3	BME	A	503	4/4	0.98	0.11	34,59,67,67	0
3	BME	C	502	4/4	0.98	0.15	16,32,38,259	0
2	SO4	B	501	5/5	0.98	0.09	49,55,60,62	0
3	BME	C	503	4/4	0.98	0.09	37,48,61,61	0

*Continued on next page...*



*Continued from previous page...*

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
2	SO4	A	501	5/5	0.98	0.10	47,55,57,62	0
3	BME	D	503	4/4	0.98	0.16	35,85,91,96	0
3	BME	B	502	4/4	0.98	0.14	31,49,63,64	0
2	SO4	D	501	5/5	0.99	0.09	52,55,56,63	0
2	SO4	C	501	5/5	0.99	0.07	53,55,60,61	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.