



Full wwPDB X-ray Structure Validation Report ⓘ

May 24, 2020 – 01:43 am BST

PDB ID : 3R62
Title : Structure of complement regulator Factor H mutant, T1184R.
Authors : Morgan, H.P.; Jiang, J.; Herbert, A.P.; Kavanagh, D.; Uhran, D.; Barlow, P.N.; Hannan, J.P.
Deposited on : 2011-03-21
Resolution : 1.52 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.11
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.11


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.52 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	4009 (1.54-1.50)
Clashscore	141614	4249 (1.54-1.50)
Ramachandran outliers	138981	4148 (1.54-1.50)
Sidechain outliers	138945	4146 (1.54-1.50)
RSRZ outliers	127900	3943 (1.54-1.50)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	129	<div> <div>4%</div> <div> <div></div> <div>76%</div> <div>16%</div> <div>5%</div> <div></div> </div> </div>
1	B	129	<div> <div>2%</div> <div> <div></div> <div>82%</div> <div>13%</div> <div></div> </div> </div>

2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 2580 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Complement factor H.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	125	Total	C	N	O	S	0	10	0
			1081	672	196	204	9			
1	B	127	Total	C	N	O	S	0	11	0
			1094	686	195	204	9			

There are 10 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	1103	GLU	-	EXPRESSION TAG	UNP P08603
A	1104	ALA	-	EXPRESSION TAG	UNP P08603
A	1105	GLU	-	EXPRESSION TAG	UNP P08603
A	1106	PHE	-	EXPRESSION TAG	UNP P08603
A	1184	ARG	THR	ENGINEERED MUTATION	UNP P08603
B	1103	GLU	-	EXPRESSION TAG	UNP P08603
B	1104	ALA	-	EXPRESSION TAG	UNP P08603
B	1105	GLU	-	EXPRESSION TAG	UNP P08603
B	1106	PHE	-	EXPRESSION TAG	UNP P08603
B	1184	ARG	THR	ENGINEERED MUTATION	UNP P08603

- Molecule 2 is GLYCEROL (three-letter code: GOL) (formula: C₃H₈O₃).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	C	O	0	0
			6	3	3		



- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	204	Total	O	0	0
			204	204		
3	B	195	Total	O	0	0
			195	195		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($\text{RSRZ} > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Complement factor H



• Molecule 1: Complement factor H



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	36.89Å 52.84Å 142.63Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	42.46 – 1.52 42.46 – 1.52	Depositor EDS
% Data completeness (in resolution range)	100.0 (42.46-1.52) 99.5 (42.46-1.52)	Depositor EDS
R_{merge}	0.11	Depositor
R_{sym}	0.12	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.04 (at 1.52Å)	Xtriage
Refinement program	REFMAC 5.5.0109	Depositor
R, R_{free}	0.178 , 0.208 0.176 , 0.206	Depositor DCC
R_{free} test set	2185 reflections (5.03%)	wwPDB-VP
Wilson B-factor (Å ²)	12.9	Xtriage
Anisotropy	0.066	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.34 , 42.5	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	2580	wwPDB-VP
Average B, all atoms (Å ²)	16.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 8.67% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: GOL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	1.40	5/1114 (0.4%)	1.29	6/1505 (0.4%)
1	B	1.30	2/1140 (0.2%)	1.21	5/1540 (0.3%)
All	All	1.35	7/2254 (0.3%)	1.25	11/3045 (0.4%)

All (7) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	1201	CYS	CB-SG	-9.91	1.65	1.82
1	A	1128	TYR	CG-CD1	5.73	1.46	1.39
1	A	1142	TYR	CE2-CZ	5.70	1.46	1.38
1	B	1120	ILE	CB-CG2	-5.54	1.35	1.52
1	B	1183	TRP	CB-CG	5.53	1.60	1.50
1	A	1190	TYR	CD2-CE2	5.42	1.47	1.39
1	A	1183	TRP	CE3-CZ3	5.18	1.47	1.38

All (11) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	1220	ASP	CB-CG-OD1	12.01	129.10	118.30
1	B	1135	GLU	OE1-CD-OE2	7.86	132.73	123.30
1	A	1153	ARG	NE-CZ-NH2	-7.69	116.46	120.30
1	A	1182	ARG	NE-CZ-NH1	-7.23	116.69	120.30
1	A	1119	ASP	CB-CG-OD2	-7.07	111.94	118.30
1	B	1202	LYS	CD-CE-NZ	-6.75	96.19	111.70
1	B	1214	LEU	CB-CG-CD1	-6.62	99.75	111.00
1	B	1177	TYR	CB-CG-CD2	-6.15	117.31	121.00
1	B	1120	ILE	CG1-CB-CG2	-5.61	99.07	111.40
1	A	1162	LYS	CD-CE-NZ	-5.52	99.00	111.70
1	A	1171	ARG	CG-CD-NE	5.47	123.28	111.80

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1081	0	1046	20	0
1	B	1094	0	1073	21	0
2	A	6	0	7	1	0
3	A	204	0	0	9	0
3	B	195	0	0	8	1
All	All	2580	0	2126	41	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 9.

All (41) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:1105:GLU:HG2	1:B:1154:ASN:HA	1.35	1.06
1:B:1183:TRP:HZ3	1:B:1198[B]:GLU:OE2	1.49	0.95
1:B:1203:ARG:HH11	1:B:1203:ARG:HB2	1.38	0.88
1:B:1135:GLU:OE1	3:B:497:HOH:O	2.00	0.78
1:A:1142:TYR:OH	2:A:1:GOL:H32	1.84	0.77
1:B:1183:TRP:CZ3	1:B:1198[B]:GLU:OE2	2.39	0.72
1:A:1107:GLY:O	3:A:515:HOH:O	2.10	0.70
1:A:1230[B]:LYS:HD3	3:A:498:HOH:O	1.94	0.67
1:B:1105:GLU:HG3	1:B:1106:PHE:N	2.08	0.67
1:A:1210:ARG:HD2	1:A:1210:ARG:O	1.96	0.65
1:B:1159[B]:GLU:HB3	3:B:310:HOH:O	1.97	0.63
1:B:1108:LYS:NZ	3:B:479:HOH:O	2.32	0.63
1:B:1105:GLU:HG3	1:B:1107:GLY:H	1.66	0.61
1:A:1206:ARG:HH11	1:A:1229:ALA:HB3	1.69	0.58
1:B:1183:TRP:HZ3	1:B:1198[B]:GLU:CD	2.07	0.56
1:B:1135:GLU:HG2	1:B:1149:ARG:HG2	1.89	0.55
1:A:1203:ARG:HH11	1:A:1203:ARG:HB3	1.71	0.55
1:B:1198[B]:GLU:HG2	1:B:1215:ARG:HG3	1.91	0.53

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:1230[B]:LYS:NZ	3:A:516:HOH:O	2.42	0.51
1:A:1202:LYS:HE3	1:A:1205:TYR:CE1	2.46	0.51
1:B:1215:ARG:HG2	3:B:97:HOH:O	2.12	0.50
1:A:1148:LYS:HD2	3:A:464:HOH:O	2.10	0.50
1:A:1203:ARG:HH11	1:A:1203:ARG:CB	2.24	0.50
1:A:1210:ARG:HD2	1:A:1210:ARG:C	2.32	0.50
1:A:1130:PRO:HB3	3:A:515:HOH:O	2.11	0.48
1:B:1222[B]:LYS:HD2	3:B:183:HOH:O	2.12	0.48
1:B:1191[B]:SER:HB2	3:B:28:HOH:O	2.14	0.48
1:A:1143:GLN:HB2	3:A:507:HOH:O	2.14	0.47
1:A:1135[B]:GLU:HG2	1:A:1149:ARG:HB2	1.96	0.47
1:A:1143:GLN:CG	3:A:507:HOH:O	2.64	0.46
1:A:1143:GLN:HG3	3:A:507:HOH:O	2.15	0.45
1:B:1203:ARG:HD3	3:B:229:HOH:O	2.16	0.44
1:B:1105:GLU:HG2	1:B:1154:ASN:CA	2.25	0.44
1:B:1108:LYS:HB3	1:B:1127[A]:VAL:HG22	2.00	0.43
1:B:1183:TRP:CZ3	1:B:1198[B]:GLU:CD	2.89	0.42
1:B:1222[B]:LYS:HE2	3:B:481:HOH:O	2.18	0.41
1:A:1143:GLN:CB	3:A:507:HOH:O	2.67	0.41
1:A:1135[B]:GLU:OE2	1:A:1149:ARG:HD2	2.20	0.41
1:A:1199:PHE:O	1:A:1214:LEU:HB3	2.21	0.40
1:A:1203:ARG:HB3	1:A:1203:ARG:NH1	2.34	0.40
1:B:1166:PRO:O	1:B:1221:GLY:HA2	2.22	0.40

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:B:84:HOH:O	3:B:462:HOH:O[3_544]	2.09	0.11

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	134/129 (104%)	129 (96%)	4 (3%)	1 (1%)	22	5
1	B	136/129 (105%)	134 (98%)	2 (2%)	0	100	100
All	All	270/258 (105%)	263 (97%)	6 (2%)	1 (0%)	34	13

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	1213	THR

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	123/115 (107%)	117 (95%)	6 (5%)	25	3
1	B	125/115 (109%)	123 (98%)	2 (2%)	62	35
All	All	248/230 (108%)	240 (97%)	8 (3%)	36	10

All (8) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	1130	PRO
1	A	1148	LYS
1	A	1203	ARG
1	A	1206	ARG
1	A	1210	ARG
1	A	1213	THR
1	B	1105	GLU
1	B	1203	ARG

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (3) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	1137	GLN
1	A	1187	GLN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	1165	HIS

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

5.6 Ligand geometry ⓘ

1 ligand is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	GOL	A	1	-	5,5,5	0.75	0	5,5,5	1.88	1 (20%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	GOL	A	1	-	-	2/4/4/4	-

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	1	GOL	C3-C2-C1	-3.97	96.25	111.70

There are no chirality outliers.

All (2) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	A	1	GOL	C1-C2-C3-O3
2	A	1	GOL	O2-C2-C3-O3

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	1	GOL	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	125/129 (96%)	0.03	5 (4%) 38 42	7, 11, 26, 44	0
1	B	127/129 (98%)	-0.01	3 (2%) 59 63	7, 12, 24, 38	0
All	All	252/258 (97%)	0.01	8 (3%) 47 52	7, 11, 25, 44	0

All (8) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	1106	PHE	3.5
1	B	1231	ARG	3.3
1	A	1213	THR	3.3
1	A	1212[A]	HIS	2.9
1	A	1203	ARG	2.6
1	B	1203	ARG	2.5
1	A	1210	ARG	2.5
1	A	1209	SER	2.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum,

median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
2	GOL	A	1	6/6	0.81	0.17	26,32,34,36	0

6.5 Other polymers [i](#)

There are no such residues in this entry.