



wwPDB X-ray Structure Validation Summary Report ⓘ

Oct 11, 2021 – 06:08 AM EDT

PDB ID : 2R7E
Title : Crystal Structure Analysis of Coagulation Factor VIII
Authors : Stoddard, B.L.; Shen, B.W.
Deposited on : 2007-09-07
Resolution : 3.70 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix)	:	1.13
EDS	:	2.23.2
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac	:	5.8.0158
CCP4	:	7.0.044 (Gargrove)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.23.2


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION





The reported resolution of this entry is 3.70 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	1049 (3.88-3.52)
Clashscore	141614	1027 (3.86-3.54)
Ramachandran outliers	138981	1069 (3.88-3.52)
Sidechain outliers	138945	1065 (3.88-3.52)
RSRZ outliers	127900	1578 (3.90-3.50)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	742	
2	B	770	
3	C	4	
4	D	2	
5	E	7	

2 Entry composition [i](#)

There are 7 unique types of molecules in this entry. The entry contains 10985 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Coagulation factor VIII.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	693	Total	C	N	O	S	0	0	0
			5590	3592	937	1035	26			

- Molecule 2 is a protein called Coagulation factor VIII.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	644	Total	C	N	O	S	0	0	0
			5229	3346	907	944	32			

There is a discrepancy between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	1838	SER	PHE	engineered mutation	UNP P00451

- Molecule 3 is an oligosaccharide called alpha-D-mannopyranose-(1-6)-beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.




Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	C	4	Total	C	N	O		0	0	0
			50	28	2	20				

- Molecule 4 is an oligosaccharide called 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
4	D	2	Total	C	N	O	0	0	0
			28	16	2	10			

- Molecule 5 is an oligosaccharide called alpha-D-mannopyranose-(1-2)-beta-D-mannopyranos e-(1-3)-[alpha-D-mannopyranose-(1-6)-alpha-D-mannopyranose-(1-6)]beta-D-mannopyranos e-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
5	E	7	Total	C	N	O	0	0	0
			83	46	2	35			

- Molecule 6 is CALCIUM ION (three-letter code: CA) (formula: Ca).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	3	Total	Ca	0	0
			3	3		


- Molecule 7 is COPPER (II) ION (three-letter code: CU) (formula: Cu).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
7	A	1	Total	Cu	0	0
			1	1		
7	B	1	Total	Cu	0	0
			1	1		


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Coagulation factor VIII




- Molecule 2: Coagulation factor VIII



L2297	D2298	P2299	P2300	L2301	L2302	T2303	R2304	Y2305	L2306	R2307	L2308	H2309	P2310	Q2311	S2312	V2313	H2314	H2315	Q2316	L2317	L2318	L2319	R2320	M2321	E2322	V2323	L2324	G2325	C2326	E2327	A2328	Q2329	D2330	L2331	Y2332
-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------

- Molecule 3: alpha-D-mannopyranose-(1-6)-beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain C:  25% 25% 50%

MAG1	MAG2	BMA3	MAN4
------	------	------	------

- Molecule 4: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain D:  100%

MAG1	MAG2
------	------

- Molecule 5: alpha-D-mannopyranose-(1-2)-beta-D-mannopyranose-(1-3)-[alpha-D-mannopyranose-(1-6)-alpha-D-mannopyranose-(1-6)]beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain E:  29% 71%

MAG1	MAG2	BMA3	BMA4	MAN5	MAN6	MAN7
------	------	------	------	------	------	------

4 Data and refinement statistics

Property	Value	Source
Space group	P 41 21 2	Depositor
Cell constants a, b, c, α , β , γ	134.57Å 134.57Å 359.50Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	57.37 – 3.70 57.36 – 3.60	Depositor EDS
% Data completeness (in resolution range)	85.0 (57.37-3.70) 81.8 (57.36-3.60)	Depositor EDS
R_{merge}	0.12	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	0.79 (at 3.57Å)	Xtriage
Refinement program	CNS 1.2	Depositor
R, R_{free}	0.279 , 0.347 0.293 , 0.357	Depositor DCC
R_{free} test set	1924 reflections (4.95%)	wwPDB-VP
Wilson B-factor (Å ²)	125.7	Xtriage
Anisotropy	0.320	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.25 , 131.9	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.89	EDS
Total number of atoms	10985	wwPDB-VP
Average B, all atoms (Å ²)	168.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.27% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: MAN, CA, NAG, CU, BMA

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z > 5$	RMSZ	# $ Z > 5$
1	A	0.56	7/5749 (0.1%)	0.83	9/7806 (0.1%)
2	B	0.44	0/5377	0.75	1/7280 (0.0%)
All	All	0.50	7/11126 (0.1%)	0.79	10/15086 (0.1%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	2

The worst 5 of 7 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	226	ARG	CZ-NH1	8.44	1.44	1.33
1	A	437	LYS	CD-CE	7.11	1.69	1.51
1	A	226	ARG	CZ-NH2	6.99	1.42	1.33
1	A	226	ARG	NE-CZ	5.36	1.40	1.33
1	A	437	LYS	CE-NZ	5.29	1.62	1.49

The worst 5 of 10 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	226	ARG	NE-CZ-NH2	-7.46	116.57	120.30
1	A	226	ARG	NE-CZ-NH1	6.08	123.34	120.30
1	A	229	PRO	N-CA-C	6.01	127.74	112.10
1	A	205	GLY	N-CA-C	-6.00	98.11	113.10
1	A	691	GLY	N-CA-C	-5.65	98.99	113.10

There are no chirality outliers.

All (2) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	385	TYR	Sidechain
1	A	656	TYR	Sidechain

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	5590	0	5429	1198	0
2	B	5229	0	5098	1107	0
3	C	50	0	43	3	0
4	D	28	0	25	0	0
5	E	83	0	70	7	0
6	A	3	0	0	0	0
7	A	1	0	0	0	0
7	B	1	0	0	0	0
All	All	10985	0	10665	2260	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 104.

The worst 5 of 2260 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:114:TYR:HE1	2:B:1997:ARG:HB2	1.11	1.13
2:B:2284:GLN:HA	2:B:2284:GLN:HE21	1.14	1.13
2:B:1901:ALA:HB1	2:B:1902:PRO:HD2	1.30	1.12
1:A:499:LYS:HG3	1:A:500:ASP:H	1.15	1.12
1:A:651:VAL:HG12	1:A:652:PHE:H	1.15	1.11

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	687/742 (93%)	312 (45%)	180 (26%)	195 (28%)	0	0
2	B	642/770 (83%)	316 (49%)	172 (27%)	154 (24%)	0	0
All	All	1329/1512 (88%)	628 (47%)	352 (26%)	349 (26%)	0	0

5 of 349 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	3	ARG
1	A	9	ALA
1	A	17	MET
1	A	19	SER
1	A	23	GLU

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	616/662 (93%)	505 (82%)	111 (18%)	1	11
2	B	572/688 (83%)	482 (84%)	90 (16%)	2	16
All	All	1188/1350 (88%)	987 (83%)	201 (17%)	2	13

5 of 201 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
2	B	1719	ARG

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
2	B	1905	ILE
2	B	2309	HIS
2	B	1737	GLU
2	B	1830	PHE

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 49 such sidechains are listed below:

Mol	Chain	Res	Type
2	B	1859	HIS
2	B	2054	HIS
2	B	1861	ASN
2	B	1915	ASN
2	B	2087	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

13 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	NAG	C	1	3,1	14,14,15	0.82	1 (7%)	17,19,21	0.87	0
3	NAG	C	2	3	14,14,15	0.71	0	17,19,21	0.76	1 (5%)
3	BMA	C	3	3	11,11,12	0.77	0	15,15,17	0.70	0
3	MAN	C	4	3	11,11,12	0.54	0	15,15,17	0.93	2 (13%)

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
4	NAG	D	1	4,2	14,14,15	0.68	0	17,19,21	0.52	0
4	NAG	D	2	4	14,14,15	0.72	0	17,19,21	0.52	0
5	NAG	E	1	5,2	14,14,15	0.71	0	17,19,21	1.23	3 (17%)
5	NAG	E	2	5	14,14,15	0.54	0	17,19,21	0.84	1 (5%)
5	BMA	E	3	5	11,11,12	1.53	4 (36%)	15,15,17	0.99	1 (6%)
5	BMA	E	4	5	11,11,12	1.05	1 (9%)	15,15,17	1.37	2 (13%)
5	MAN	E	5	5	11,11,12	0.66	0	15,15,17	0.91	1 (6%)
5	MAN	E	6	5	11,11,12	1.34	3 (27%)	15,15,17	0.76	0
5	MAN	E	7	5	11,11,12	1.79	1 (9%)	15,15,17	0.89	1 (6%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	NAG	C	1	3,1	-	5/6/23/26	0/1/1/1
3	NAG	C	2	3	-	4/6/23/26	0/1/1/1
3	BMA	C	3	3	-	1/2/19/22	0/1/1/1
3	MAN	C	4	3	-	1/2/19/22	0/1/1/1
4	NAG	D	1	4,2	-	4/6/23/26	0/1/1/1
4	NAG	D	2	4	-	2/6/23/26	0/1/1/1
5	NAG	E	1	5,2	-	5/6/23/26	0/1/1/1
5	NAG	E	2	5	-	4/6/23/26	0/1/1/1
5	BMA	E	3	5	-	2/2/19/22	0/1/1/1
5	BMA	E	4	5	-	2/2/19/22	0/1/1/1
5	MAN	E	5	5	-	1/2/19/22	0/1/1/1
5	MAN	E	6	5	-	0/2/19/22	0/1/1/1
5	MAN	E	7	5	-	2/2/19/22	0/1/1/1

The worst 5 of 10 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
5	E	7	MAN	C2-C3	4.53	1.59	1.52
5	E	4	BMA	C1-C2	2.63	1.58	1.52
5	E	6	MAN	C2-C3	2.47	1.56	1.52
5	E	6	MAN	C4-C5	2.35	1.58	1.53
3	C	1	NAG	C1-C2	2.34	1.55	1.52

The worst 5 of 12 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
5	E	4	BMA	C1-O5-C5	3.92	117.51	112.19
5	E	1	NAG	C2-N2-C7	-2.63	119.17	122.90
5	E	4	BMA	C1-C2-C3	2.47	112.70	109.67
5	E	5	MAN	C1-C2-C3	2.44	112.67	109.67
5	E	1	NAG	C3-C4-C5	2.32	114.38	110.24

There are no chirality outliers.

5 of 33 torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
3	C	1	NAG	C3-C2-N2-C7
3	C	1	NAG	C8-C7-N2-C2
3	C	1	NAG	O7-C7-N2-C2
3	C	2	NAG	C8-C7-N2-C2
3	C	2	NAG	O7-C7-N2-C2


There are no ring outliers.


7 monomers are involved in 10 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
5	E	5	MAN	3	0
5	E	3	BMA	3	0
3	C	2	NAG	2	0
5	E	1	NAG	2	0
5	E	4	BMA	3	0
5	E	2	NAG	1	0
3	C	1	NAG	3	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.







5.6 Ligand geometry [i](#)

Of 5 ligands modelled in this entry, 5 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	693/742 (93%)	-0.26	18 (2%) 56 43	75, 170, 206, 206	0
2	B	644/770 (83%)	-0.37	9 (1%) 75 64	89, 169, 206, 206	0
All	All	1337/1512 (88%)	-0.32	27 (2%) 65 53	75, 170, 206, 206	0

The worst 5 of 27 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	725	ASP	5.0
1	A	23	GLU	4.7
1	A	225	ALA	4.3
1	A	24	LEU	3.7
1	A	37	SER	3.7

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.


Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
3	BMA	C	3	11/12	0.45	0.32	148,148,148,148	11
3	MAN	C	4	11/12	0.68	0.34	115,115,115,115	11
3	NAG	C	2	14/15	0.79	0.23	122,122,122,122	14
5	BMA	E	3	11/12	0.82	0.12	203,203,203,203	0
4	NAG	D	2	14/15	0.84	0.23	180,180,180,180	0

Continued on next page...

Continued from previous page...




Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
5	BMA	E	4	11/12	0.84	0.20	145,145,145,145	11
3	NAG	C	1	14/15	0.86	0.18	205,205,205,205	0
4	NAG	D	1	14/15	0.87	0.19	205,205,205,205	0
5	NAG	E	2	14/15	0.89	0.25	202,205,205,205	0
5	MAN	E	6	11/12	0.89	0.17	187,187,187,187	0
5	MAN	E	5	11/12	0.90	0.18	104,104,104,104	11
5	MAN	E	7	11/12	0.92	0.14	190,190,190,190	0
5	NAG	E	1	14/15	0.95	0.15	150,153,155,156	0


The following is a graphical depiction of the model fit to experimental electron density for oligosaccharide. Each fit is shown from different orientation to approximate a three-dimensional view.



Electron density around Chain D:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)





6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
6	CA	A	2400	1/1	0.56	0.39	341,341,341,341	0
6	CA	A	2402	1/1	0.96	0.17	352,352,352,352	0
7	CU	A	2403	1/1	0.96	0.29	359,359,359,359	0
6	CA	A	2401	1/1	0.98	0.31	275,275,275,275	0
7	CU	B	2404	1/1	0.98	0.31	349,349,349,349	0

6.5 Other polymers [i](#)

There are no such residues in this entry.