



# wwPDB X-ray Structure Validation Summary Report ⓘ

May 22, 2020 – 02:27 am BST

PDB ID : 4R7Y  
Title : Crystal structure of an active MCM hexamer  
Authors : Miller, J.M.; Arachea, B.T.; Epling, L.B.; Enemark, E.J.  
Deposited on : 2014-08-28  
Resolution : 2.70 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix)	:	1.13
EDS	:	2.11
buster-report	:	1.1.7 (2018)
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac	:	5.8.0158
CCP4	:	7.0.044 (Gargrove)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.11


# 1 Overall quality at a glance ⓘ

The following experimental techniques were used to determine the structure:

## *X-RAY DIFFRACTION*

The reported resolution of this entry is 2.70 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	2808 (2.70-2.70)
Clashscore	141614	3122 (2.70-2.70)
Ramachandran outliers	138981	3069 (2.70-2.70)
Sidechain outliers	138945	3069 (2.70-2.70)
RSRZ outliers	127900	2737 (2.70-2.70)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	613	<div> <div>9%</div> <div>44%</div> <div>43%</div> <div>9%</div> <div>••</div> </div>
1	B	613	<div> <div>8%</div> <div>44%</div> <div>44%</div> <div>8%</div> <div>•</div> </div>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
5	CL	A	2006	-	-	X	-
5	CL	B	2006	-	-	X	-

## 2 Entry composition [i](#)

There are 5 unique types of molecules in this entry. The entry contains 9494 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Minichromosome maintenance protein MCM, Cell division control protein 21.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	591	Total	C	N	O	S	0	0	0
			4715	3001	822	878	14			
1	B	591	Total	C	N	O	S	0	0	0
			4715	3001	822	878	14			

There are 4 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	0	SER	-	EXPRESSION TAG	UNP Q9UXG1
A	1	LEU	-	EXPRESSION TAG	UNP Q9UXG1
B	0	SER	-	EXPRESSION TAG	UNP Q9UXG1
B	1	LEU	-	EXPRESSION TAG	UNP Q9UXG1

- Molecule 2 is ADENOSINE-5'-DIPHOSPHATE (three-letter code: ADP) (formula:  $C_{10}H_{15}N_5O_{10}P_2$ ).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total	C	N	O	P	0	0
			27	10	5	10	2		
2	B	1	Total	C	N	O	P	0	0
			27	10	5	10	2		

- Molecule 3 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	B	1	Total	Mg	0	0
			1	1		
3	A	1	Total	Mg	0	0
			1	1		

- Molecule 4 is ZINC ION (three-letter code: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	B	1	Total	Zn	0	0
			1	1		
4	A	1	Total	Zn	0	0
			1	1		


- Molecule 5 is CHLORIDE ION (three-letter code: CL) (formula: Cl).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	B	3	Total	Cl	0	0
			3	3		
5	A	3	Total	Cl	0	0
			3	3		


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Minichromosome maintenance protein MCM, Cell division control protein 21



- Molecule 1: Minichromosome maintenance protein MCM, Cell division control protein 21



GLU	ILE	PRO	PRO	ILE	P1919	L1852	H1773	R1319	S260	K173	I103	SER
GLY	PRO	PRO	PRO	ILE	I1920	R1853	E1774	G1320	S261	C174	I104	LEU
ILE	PRO	PRO	PRO	ILE	T1921	V1854	A1775	I1322	E263	G175	P105	GLU
PRO	PRO	PRO	PRO	ILE	R1855	R1855				K176	R106	ILE
PRO	PRO	PRO	PRO	ILE	R1856	R1856				P177	E108	PRO
ILE	PRO	PRO	PRO	ILE	G1857	T1780	L1325	L1325	K267	G178	L109	SER
P1919	I1920	T1921	R1923	R1923	F1858	I1781	L1326	V1327	I1261	G179	R110	LYS
I1920	T1921	R1923	R1923	R1923	S1782	I1783	V1327	G1328	E1264	F180	R111	Q7
R1923	R1923	R1923	R1923	R1923	I1783		D1329		E1265	R181	D9	I8
Q1924	Q1924	Q1924	Q1924	Q1924	T1789				Q1267	L182	Y10	R11
L1925	L1925	L1925	L1925	L1925	T1796				Q1267	I183	R113	R12
E1926	E1926	E1926	E1926	E1926	A1800				I1268	K186	F13	D12
A1927	A1927	A1927	A1927	A1927					I1269	T187	L120	F14
L1928	L1928	L1928	L1928	L1928	N1803				Q1336	K188	I121	R15
I1929	I1929	I1929	I1929	I1929	P1804				Q1336	Q193	T122	E16
R1930	R1930	R1930	R1930	R1930	K1805				L1271	R200	D124	N25
L1931	L1931	L1931	L1931	L1931					L1272	P201	V130	R26
S1932	S1932	S1932	S1932	S1932	R1808				K1273	E202	I131	Q27
E1933	E1933	E1933	E1933	E1933	F1809				K1274	E203	P132	Y30
A1934	A1934	A1934	A1934	A1934	M1810				K1276	P205	E135	I34
H1935	H1935	H1935	H1935	H1935	M1811				D1277	Q208	I137	R41
A1936	A1936	A1936	A1936	A1936	R1812				V1279	L209	Y138	I47
R1937	R1937	R1937	R1937	R1937	K1813				D1280	R211	T141	E48
M1938	M1938	M1938	M1938	M1938	M1814				I1281	E214	Y142	F49
R1939	R1939	R1939	R1939	R1939	N1815				I1282	L215	K143	Y61
L1940	L1940	L1940	L1940	L1940	I1816				D1284	L216	H144	E62
P1942	P1942	P1942	P1942	P1942	F1817					L217	I146	I63
I1943	I1943	I1943	I1943	I1943	R1818						P147	I64
V1944	V1944	V1944	V1944	V1944	Q1818						D148	N65
T1945	T1945	T1945	T1945	T1945	I1819						C149	K68
R1946	R1946	R1946	R1946	R1946	P1823						R190	I69
E1947	E1947	E1947	E1947	E1947	L1824						Q151	I70
D1948	D1948	D1948	D1948	D1948	L1825						E152	I71
A1949	A1949	A1949	A1949	A1949	L1826						F153	L71
R1950	R1950	R1950	R1950	R1950	S1827						E154	P72
E1951	E1951	E1951	E1951	E1951	R1828						V155	I73
A1952	A1952	A1952	A1952	A1952	F1829						P156	L74
I1953	I1953	I1953	I1953	I1953	D1830						E157	A77
K1954	K1954	K1954	K1954	K1954	F1833						E159	I82
L1955	L1955	L1955	L1955	L1955							E160	
I1963	I1963	I1963	I1963	I1963	I1836						M161	D86
A1964	A1964	A1964	A1964	A1964	D1837						P162	P87
M1965	M1965	M1965	M1965	M1965	E1838						E163	T88
ASP	ASP	ASP	ASP	ASP	P1839						E166	Y89
					R1900						M167	Q90
					M1901						P168	R91
					R1902						T169	V96
					K1903						C171	I100
					S1904						P172	
					VAL							
					LYS							
					LYS							
					THR							
					LYS							
					GLY							
					GLU							
					GLU							

## 4 Data and refinement statistics

Property	Value	Source
Space group	P 63	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	118.90Å 118.90Å 199.32Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	49.83 – 2.70 49.83 – 2.70	Depositor EDS
% Data completeness (in resolution range)	89.7 (49.83-2.70) 89.7 (49.83-2.70)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	0.11	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.24 (at 2.69Å)	Xtriage
Refinement program	CNS 1.3	Depositor
R, $R_{free}$	0.263 , 0.295 0.254 , 0.290	Depositor DCC
$R_{free}$ test set	1976 reflections (5.06%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	39.4	Xtriage
Anisotropy	0.248	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.32 , 44.7	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.46$ , $\langle L^2 \rangle = 0.29$	Xtriage
Estimated twinning fraction	0.044 for h,-h-k,-l	Xtriage
$F_o, F_c$ correlation	0.92	EDS
Total number of atoms	9494	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	61.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 49.42 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 7.4816e-05. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

<sup>1</sup> Intensities estimated from amplitudes.

<sup>2</sup> Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



## 5 Model quality

### 5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: ZN, MG, ADP, CL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.45	0/4793	0.69	0/6476
1	B	0.45	0/4793	0.70	0/6476
All	All	0.45	0/9586	0.70	0/12952

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4715	0	4876	420	0
1	B	4715	0	4876	393	0
2	A	27	0	12	3	0
2	B	27	0	12	1	0
3	A	1	0	0	0	0
3	B	1	0	0	0	0
4	A	1	0	0	0	0
4	B	1	0	0	0	0
5	A	3	0	0	2	0
5	B	3	0	0	2	0
All	All	9494	0	9776	798	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 41.

The worst 5 of 798 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:1346:PRO:HB3	1:B:1872:ARG:HD3	1.29	1.10
1:A:156:PRO:HB2	1:A:159:GLU:O	1.53	1.07
1:B:1261:ILE:HA	1:B:1873:LYS:HZ1	1.17	1.05
1:A:1878:ALA:HA	1:A:1882:ILE:HD11	1.37	1.05
1:B:1261:ILE:HA	1:B:1873:LYS:NZ	1.73	1.04

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	587/613 (96%)	519 (88%)	53 (9%)	15 (3%)	5	13
1	B	587/613 (96%)	524 (89%)	47 (8%)	16 (3%)	5	12
All	All	1174/1226 (96%)	1043 (89%)	100 (8%)	31 (3%)	5	13

5 of 31 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	155	TRP
1	A	247	ARG
1	A	1841	ASP
1	A	1857	GLY
1	B	155	TRP

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	517/537 (96%)	444 (86%)	73 (14%)	3	8
1	B	517/537 (96%)	461 (89%)	56 (11%)	6	15
All	All	1034/1074 (96%)	905 (88%)	129 (12%)	4	10

5 of 129 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	1892	GLU
1	A	1963	ILE
1	B	1879	ARG
1	A	1896	LYS
1	A	1940	LEU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 15 such sidechains are listed below:

Mol	Chain	Res	Type
1	A	1814	ASN
1	A	1962	GLN
1	B	1810	ASN
1	A	1779	GLN
1	B	1779	GLN

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 5.6 Ligand geometry [i](#)

Of 12 ligands modelled in this entry, 10 are monoatomic - leaving 2 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
2	ADP	B	2001	3	24,29,29	1.44	3 (12%)	29,45,45	1.80	4 (13%)
2	ADP	A	2001	3	24,29,29	1.32	3 (12%)	29,45,45	1.86	4 (13%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	ADP	B	2001	3	-	0/12/32/32	0/3/3/3
2	ADP	A	2001	3	-	0/12/32/32	0/3/3/3

The worst 5 of 6 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	2001	ADP	O4'-C1'	3.67	1.46	1.41
2	B	2001	ADP	PB-O1B	3.31	1.61	1.50
2	A	2001	ADP	O4'-C1'	3.18	1.45	1.41
2	A	2001	ADP	PB-O1B	3.13	1.60	1.50
2	B	2001	ADP	C8-N7	-2.80	1.29	1.34

The worst 5 of 8 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	2001	ADP	PA-O3A-PB	-6.59	110.20	132.83
2	B	2001	ADP	PA-O3A-PB	-6.43	110.77	132.83

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	2001	ADP	N3-C2-N1	-4.61	121.48	128.68
2	B	2001	ADP	N3-C2-N1	-4.13	122.22	128.68
2	B	2001	ADP	C3'-C2'-C1'	2.99	105.48	100.98

There are no chirality outliers.


There are no torsion outliers.

There are no ring outliers.

2 monomers are involved in 4 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	B	2001	ADP	1	0
2	A	2001	ADP	3	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



## 5.7 Other polymers ⓘ

There are no such residues in this entry.

## 5.8 Polymer linkage issues ⓘ

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	591/613 (96%)	0.54	56 (9%) <b>8</b> <b>6</b>	10, 63, 109, 125	0
1	B	591/613 (96%)	0.47	51 (8%) <b>10</b> <b>8</b>	9, 61, 110, 128	0
All	All	1182/1226 (96%)	0.51	107 (9%) <b>9</b> <b>7</b>	9, 62, 110, 128	0

The worst 5 of 107 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	248	GLY	8.0
1	B	249	SER	7.7
1	A	249	SER	7.6
1	B	1315	GLY	6.6
1	B	146	HIS	6.4

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.


### 6.4 Ligands [i](#)


In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.



Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
2	ADP	A	2001	27/27	0.80	0.27	113,114,115,115	0
2	ADP	B	2001	27/27	0.82	0.27	118,121,127,128	0
4	ZN	B	2003	1/1	0.88	0.05	169,169,169,169	0
5	CL	B	2006	1/1	0.92	0.09	39,39,39,39	0
4	ZN	A	2003	1/1	0.93	0.09	163,163,163,163	0
3	MG	B	2002	1/1	0.94	0.19	64,64,64,64	0
3	MG	A	2002	1/1	0.95	0.13	64,64,64,64	0
5	CL	B	2004	1/1	0.96	0.23	48,48,48,48	0
5	CL	A	2005	1/1	0.96	0.21	59,59,59,59	0
5	CL	B	2005	1/1	0.96	0.17	53,53,53,53	0
5	CL	A	2004	1/1	0.97	0.25	48,48,48,48	0
5	CL	A	2006	1/1	0.99	0.22	41,41,41,41	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.





## 6.5 Other polymers [i](#)

There are no such residues in this entry.