



Full wwPDB X-ray Structure Validation Report ⓘ

May 13, 2020 – 09:31 am BST

PDB ID : 5RAQ
Title : PanDDA analysis group deposition – Crystal Structure of JMJD1B in complex with FM001577a
Authors : Snee, M.; Nowak, R.; Johansson, C.; Burgess-Brown, N.A.; Arrowsmith, C.H.; Bountra, C.; Edwards, A.M.; Oppermann, U.
Deposited on : 2020-03-16
Resolution : 1.85 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.11
buster-report : 1.1.7 (2018)
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.11


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.85 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	2469 (1.86-1.86)
Clashscore	141614	2625 (1.86-1.86)
Ramachandran outliers	138981	2592 (1.86-1.86)
Sidechain outliers	138945	2592 (1.86-1.86)
RSRZ outliers	127900	2436 (1.86-1.86)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	372	 6% 81% 9% • 9%
1	B	372	 4% 84% 8% 8%

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:


Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	S5J	A	1801	-	-	-	X


ENTRY-COMPOSITION INFOmissingINFO

2 Residue-property plots [i](#)


These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


• Molecule 1: Lysine-specific demethylase 3B

Chain A: 



• Molecule 1: Lysine-specific demethylase 3B

Chain B: 



3 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	57.37Å 93.72Å 93.47Å 90.00° 108.11° 90.00°	Depositor
Resolution (Å)	64.56 – 1.85 64.48 – 1.85	Depositor EDS
% Data completeness (in resolution range)	98.6 (64.56-1.85) 98.7 (64.48-1.85)	Depositor EDS
R_{merge}	0.06	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.76 (at 1.86Å)	Xtriage
Refinement program	REFMAC 5.8.0238	Depositor
R, R_{free}	0.181 , 0.216 0.198 , 0.230	Depositor DCC
R_{free} test set	3908 reflections (4.94%)	wwPDB-VP
Wilson B-factor (Å ²)	29.7	Xtriage
Anisotropy	0.120	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.32 , 57.4	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtriage
Estimated twinning fraction	0.034 for h,-k,-h-l	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	6301	wwPDB-VP
Average B, all atoms (Å ²)	38.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.83% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

4 Model quality

4.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: MN, S5J, CL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.81	0/2835	0.86	0/3846
1	B	0.76	0/2872	0.82	0/3895
All	All	0.79	0/5707	0.84	0/7741

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

4.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2762	0	2646	29	0
1	B	2798	0	2685	26	0
2	A	17	0	0	0	0
2	B	17	0	0	0	0
3	A	4	0	0	0	0
3	B	1	0	0	0	0
4	A	1	0	0	0	0
4	B	1	0	0	0	0
5	A	369	0	0	15	0
5	B	331	0	0	6	0
All	All	6301	0	5331	55	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (55) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:1470:CYS:O	5:A:1901:HOH:O	1.60	1.19
1:B:1470:CYS:SG	5:B:2219:HOH:O	2.36	0.84
1:A:1406[A]:ARG:NH2	5:A:1904:HOH:O	2.10	0.84
1:B:1644:HIS:NE2	1:B:1649:ASP:OD2	2.21	0.73
1:B:1516:LYS:HE3	5:B:1928:HOH:O	1.88	0.73
1:A:1601:GLU:OE1	5:A:1903:HOH:O	2.05	0.73
1:B:1517[B]:ARG:HH11	1:B:1517[B]:ARG:CG	2.03	0.70
1:B:1602:ARG:O	1:B:1606:HIS:O	2.08	0.70
1:B:1517[B]:ARG:HH11	1:B:1517[B]:ARG:HG3	1.60	0.66
1:B:1517[B]:ARG:NH1	1:B:1517[B]:ARG:HG3	2.12	0.64
1:B:1518:ASP:HB3	5:B:2133:HOH:O	1.99	0.63
1:A:1711:LYS:CB	5:A:2088:HOH:O	2.47	0.62
1:A:1522:ASN:O	1:A:1526:ARG:NH1	2.33	0.61
1:A:1505:LEU:HD22	1:A:1700:VAL:HG11	1.82	0.61
1:B:1518:ASP:OD1	5:B:1901:HOH:O	2.16	0.58
1:B:1506:MET:CE	1:B:1538:PRO:HG3	2.34	0.57
1:A:1582:HIS:CE1	5:A:1934:HOH:O	2.57	0.57
1:A:1639:GLU:O	5:A:1905:HOH:O	2.18	0.57
1:B:1664[B]:GLU:HA	1:B:1664[B]:GLU:OE1	2.06	0.55
1:A:1406[A]:ARG:NH1	5:A:1915:HOH:O	2.39	0.55
1:A:1440:ASP:HB2	5:A:2038:HOH:O	2.06	0.54
1:B:1388:GLY:N	5:B:1904:HOH:O	2.29	0.54
1:B:1494:PHE:CZ	1:B:1498:MET:HE3	2.43	0.54
1:B:1505:LEU:HD22	1:B:1700:VAL:HG11	1.90	0.53
1:A:1506:MET:HE1	1:A:1538:PRO:HD3	1.91	0.52
1:A:1585:GLU:CD	1:A:1585:GLU:H	2.12	0.51
1:A:1631:LYS:HE2	1:A:1666:TYR:OH	2.10	0.51
1:B:1517[B]:ARG:CB	1:B:1517[B]:ARG:HH11	2.23	0.51
1:B:1494:PHE:CE2	1:B:1498:MET:HE3	2.47	0.50
1:B:1606:HIS:CD2	1:B:1606:HIS:H	2.29	0.50
1:A:1649:ASP:CB	5:A:2053:HOH:O	2.60	0.49
1:A:1505:LEU:HD22	1:A:1700:VAL:CG1	2.41	0.49
1:B:1517[B]:ARG:HB2	1:B:1517[B]:ARG:HH11	1.77	0.48
1:A:1384:TRP:HB3	1:A:1388:GLY:HA2	1.96	0.47
1:A:1582:HIS:O	1:A:1582:HIS:CG	2.68	0.46
1:A:1540:MET:HE2	1:A:1698:ILE:HG23	1.98	0.46
1:B:1505:LEU:HD23	1:B:1506:MET:HE2	1.97	0.46

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:1649:ASP:HB3	5:A:2053:HOH:O	2.16	0.46
1:B:1561:LEU:C	1:B:1561:LEU:HD12	2.37	0.45
1:B:1446:VAL:HG22	1:B:1453:ILE:HD13	1.98	0.45
1:B:1548:THR:O	1:B:1551:ASP:HB2	2.16	0.45
1:A:1407:GLU:CG	5:A:2204:HOH:O	2.64	0.44
1:B:1443:VAL:HG21	1:B:1485:LEU:HD11	1.98	0.44
1:A:1624:LYS:NZ	5:A:1902:HOH:O	1.84	0.44
1:A:1407:GLU:HG3	5:A:2204:HOH:O	2.17	0.44
1:A:1379[A]:MET:CE	5:A:2100:HOH:O	2.65	0.44
1:B:1606:HIS:HD2	1:B:1606:HIS:H	1.65	0.43
1:A:1582:HIS:HE1	5:A:1934:HOH:O	1.97	0.43
1:B:1379:MET:N	5:B:1913:HOH:O	2.50	0.43
1:A:1561:LEU:C	1:A:1561:LEU:HD12	2.40	0.42
1:A:1502:PHE:CE2	1:A:1506:MET:HE2	2.54	0.42
1:A:1385:LEU:HA	1:A:1385:LEU:HD23	1.88	0.41
1:A:1407:GLU:O	1:A:1410:LYS:HG2	2.19	0.41
1:B:1506:MET:HE3	1:B:1538:PRO:HG3	2.03	0.41
1:A:1506:MET:CE	1:A:1538:PRO:HD3	2.50	0.41

There are no symmetry-related clashes.

4.3 Torsion angles [i](#)

4.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	342/372 (92%)	333 (97%)	7 (2%)	2 (1%)	25	12
1	B	342/372 (92%)	335 (98%)	6 (2%)	1 (0%)	41	26
All	All	684/744 (92%)	668 (98%)	13 (2%)	3 (0%)	34	19

All (3) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	1581	ALA
1	B	1594	ASP
1	A	1594	ASP

4.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	298/331 (90%)	288 (97%)	10 (3%)	37	19
1	B	302/331 (91%)	296 (98%)	6 (2%)	55	40
All	All	600/662 (91%)	584 (97%)	16 (3%)	47	29

All (16) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	1406[A]	ARG
1	A	1406[B]	ARG
1	A	1577	ILE
1	A	1579	GLU
1	A	1583	ASP
1	A	1584	GLU
1	A	1585	GLU
1	A	1644	HIS
1	A	1661	ARG
1	A	1717	THR
1	B	1383	SER
1	B	1582	HIS
1	B	1587	LEU
1	B	1588	LYS
1	B	1603	ILE
1	B	1606	HIS

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (4) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	1640	ASN
1	A	1650	GLN
1	B	1606	HIS
1	B	1656	GLN

4.3.3 RNA [i](#)

There are no RNA molecules in this entry.

4.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

4.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

4.6 Ligand geometry [i](#)

Of 9 ligands modelled in this entry, 2 could not be matched to an existing wwPDB Chemical Component Dictionary definition at this stage and 7 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.



There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier.

The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



4.7 Other polymers [i](#)

There are no such residues in this entry.

4.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

5 Fit of model and data ⓘ

5.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2		OWAB(Å ²)	Q<0.9
1	A	340/372 (91%)	-0.01	24 (7%)	16 15	18, 31, 66, 110	12 (3%)
1	B	344/372 (92%)	-0.22	16 (4%)	31 30	21, 35, 67, 103	31 (9%)
All	All	684/744 (91%)	-0.12	40 (5%)	23 22	18, 33, 67, 110	43 (6%)

All (40) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	1384	TRP	13.3
1	A	1386	CYS	13.1
1	A	1385	LEU	12.4
1	A	1392	CYS	11.2
1	A	1379[A]	MET	10.2
1	A	1578	GLY	9.8
1	A	1577	ILE	9.7
1	A	1383	SER	9.2
1	A	1388	GLY	8.6
1	A	1394	HIS	8.5
1	A	1387	ASP	6.8
1	A	1581	ALA	6.2
1	B	1581	ALA	5.6
1	A	1588[A]	LYS	5.4
1	A	1564[A]	SER	5.3
1	A	1406[A]	ARG	4.4
1	B	1577	ILE	4.4
1	B	1517[A]	ARG	4.3
1	B	1582	HIS	3.8
1	B	1586	VAL	3.7
1	B	1583	ASP	3.6
1	B	1664[A]	GLU	3.4
1	A	1642	PRO	3.3
1	B	1606	HIS	3.1

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	1580	GLY	3.1
1	B	1387	ASP	3.0
1	A	1579	GLU	2.9
1	B	1642	PRO	2.9
1	A	1582	HIS	2.8
1	B	1603	ILE	2.7
1	A	1717	THR	2.7
1	B	1590	ILE	2.6
1	A	1378	SER	2.5
1	A	1639	GLU	2.5
1	A	1641	PRO	2.4
1	B	1587	LEU	2.2
1	A	1606	HIS	2.1
1	B	1639	GLU	2.1
1	B	1588	LYS	2.0
1	B	1589	THR	2.0

5.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.4 Ligands [i](#)

LIGAND-RSR INFOmissingINFO

5.5 Other polymers [i](#)

There are no such residues in this entry.