



Full wwPDB X-ray Structure Validation Report ⓘ

May 15, 2020 – 02:44 am BST

PDB ID : 3RBT
Title : Crystal structure of glutathione S-transferase Omega 3 from the silkworm Bombyx mori
Authors : Chen, B.-Y.; Ma, X.-X.; Tan, X.; Yang, J.-P.; Zhang, N.-N.; Li, W.-F.; Chen, Y.; Xia, Q.; Zhou, C.-Z.
Deposited on : 2011-03-29
Resolution : 2.20 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.11
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.11


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.20 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	4898 (2.20-2.20)
Clashscore	141614	5594 (2.20-2.20)
Ramachandran outliers	138981	5503 (2.20-2.20)
Sidechain outliers	138945	5504 (2.20-2.20)
RSRZ outliers	127900	4800 (2.20-2.20)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	246	<div> <div>3%</div> <div> <div></div> <div>81%</div> <div>11%</div> <div>7%</div> </div> </div>
1	B	246	<div> <div>2%</div> <div> <div></div> <div>82%</div> <div>11%</div> <div>6%</div> </div> </div>
1	C	246	<div> <div>9%</div> <div> <div></div> <div>86%</div> <div>7%</div> <div>7%</div> </div> </div>
1	D	246	<div> <div>4%</div> <div> <div></div> <div>85%</div> <div>12%</div> <div>..</div> </div> </div>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit crite-

ria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	GOL	A	241	-	-	X	-
2	GOL	B	241	-	-	X	-
2	GOL	C	241	-	-	X	-

2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 8238 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Glutathione transferase o1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	230	Total	C	N	O	S	0	0	0
			1946	1261	333	344	8			
1	B	232	Total	C	N	O	S	0	0	0
			1949	1257	336	348	8			
1	C	228	Total	C	N	O	S	0	0	0
			1922	1243	331	340	8			
1	D	242	Total	C	N	O	S	0	0	0
			2040	1320	352	360	8			

There are 28 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-5	HIS	-	EXPRESSION TAG	UNP Q1HPV9
A	-4	HIS	-	EXPRESSION TAG	UNP Q1HPV9
A	-3	HIS	-	EXPRESSION TAG	UNP Q1HPV9
A	-2	HIS	-	EXPRESSION TAG	UNP Q1HPV9
A	-1	HIS	-	EXPRESSION TAG	UNP Q1HPV9
A	0	HIS	-	EXPRESSION TAG	UNP Q1HPV9
A	1	GLY	-	EXPRESSION TAG	UNP Q1HPV9
B	-5	HIS	-	EXPRESSION TAG	UNP Q1HPV9
B	-4	HIS	-	EXPRESSION TAG	UNP Q1HPV9
B	-3	HIS	-	EXPRESSION TAG	UNP Q1HPV9
B	-2	HIS	-	EXPRESSION TAG	UNP Q1HPV9
B	-1	HIS	-	EXPRESSION TAG	UNP Q1HPV9
B	0	HIS	-	EXPRESSION TAG	UNP Q1HPV9
B	1	GLY	-	EXPRESSION TAG	UNP Q1HPV9
C	-5	HIS	-	EXPRESSION TAG	UNP Q1HPV9
C	-4	HIS	-	EXPRESSION TAG	UNP Q1HPV9
C	-3	HIS	-	EXPRESSION TAG	UNP Q1HPV9
C	-2	HIS	-	EXPRESSION TAG	UNP Q1HPV9
C	-1	HIS	-	EXPRESSION TAG	UNP Q1HPV9
C	0	HIS	-	EXPRESSION TAG	UNP Q1HPV9
C	1	GLY	-	EXPRESSION TAG	UNP Q1HPV9

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
D	-5	HIS	-	EXPRESSION TAG	UNP Q1HPV9
D	-4	HIS	-	EXPRESSION TAG	UNP Q1HPV9
D	-3	HIS	-	EXPRESSION TAG	UNP Q1HPV9
D	-2	HIS	-	EXPRESSION TAG	UNP Q1HPV9
D	-1	HIS	-	EXPRESSION TAG	UNP Q1HPV9
D	0	HIS	-	EXPRESSION TAG	UNP Q1HPV9
D	1	GLY	-	EXPRESSION TAG	UNP Q1HPV9

- Molecule 2 is GLYCEROL (three-letter code: GOL) (formula: $C_3H_8O_3$).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	A	1	Total C O 6 3 3	0	0
2	B	1	Total C O 6 3 3	0	0
2	C	1	Total C O 6 3 3	0	0
2	D	1	Total C O 6 3 3	0	0

- Molecule 3 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	96	Total O 96 96	0	0
3	B	98	Total O 98 98	0	0

Continued on next page...


Continued from previous page...

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	C	78	Total	O	0	0
			78	78		
3	D	85	Total	O	0	0
			85	85		



3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.



• Molecule 1: Glutathione transferase o1




• Molecule 1: Glutathione transferase o1



• Molecule 1: Glutathione transferase o1



• Molecule 1: Glutathione transferase o1





4 Data and refinement statistics

Property	Value	Source
Space group	C 2 2 21	Depositor
Cell constants a, b, c, α , β , γ	108.54Å 139.03Å 128.82Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	41.50 – 2.20 42.94 – 2.20	Depositor EDS
% Data completeness (in resolution range)	99.7 (41.50-2.20) 99.7 (42.94-2.20)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.25 (at 2.20Å)	Xtrriage
Refinement program	REFMAC 5.5.0072	Depositor
R, R_{free}	0.215 , 0.263 0.220 , 0.271	Depositor DCC
R_{free} test set	2508 reflections (5.06%)	wwPDB-VP
Wilson B-factor (Å ²)	29.3	Xtrriage
Anisotropy	0.187	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.38 , 47.1	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.32$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	8238	wwPDB-VP
Average B, all atoms (Å ²)	32.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 29.63 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 1.4990e-03. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: GOL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.43	0/2003	0.54	0/2712
1	B	0.49	1/2003 (0.0%)	0.56	1/2711 (0.0%)
1	C	0.42	0/1976	0.52	0/2674
1	D	0.42	0/2102	0.54	0/2847
All	All	0.44	1/8084 (0.0%)	0.54	1/10944 (0.0%)

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	122	GLU	CB-CG	-5.33	1.42	1.52

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	123	LEU	CA-CB-CG	5.38	127.68	115.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1946	0	1911	22	0
1	B	1949	0	1915	27	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	C	1922	0	1895	13	0
1	D	2040	0	1991	17	0
2	A	6	0	8	9	0
2	B	6	0	8	12	0
2	C	6	0	8	6	0
2	D	6	0	8	1	0
3	A	96	0	0	1	0
3	B	98	0	0	0	0
3	C	78	0	0	1	0
3	D	85	0	0	0	0
All	All	8238	0	7744	79	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (79) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:110:LYS:NZ	2:B:241:GOL:H12	1.53	1.24
1:A:95:GLU:HG3	2:A:241:GOL:H2	1.42	0.98
1:C:95:GLU:HG3	2:C:241:GOL:H2	1.48	0.94
1:B:110:LYS:HZ1	2:B:241:GOL:H12	1.16	0.92
1:C:110:LYS:NZ	2:C:241:GOL:H32	1.85	0.90
1:B:141:GLN:O	1:B:145:THR:HG22	1.74	0.86
1:B:110:LYS:HZ2	2:B:241:GOL:H12	1.42	0.85
1:C:110:LYS:HZ2	2:C:241:GOL:H32	1.40	0.83
1:C:123:LEU:HB2	1:C:145:THR:HG21	1.65	0.77
1:B:110:LYS:NZ	2:B:241:GOL:C1	2.45	0.74
1:A:103:HIS:CD2	2:A:241:GOL:H32	2.24	0.72
1:A:95:GLU:CG	2:A:241:GOL:H2	2.20	0.70
2:A:241:GOL:H31	3:A:258:HOH:O	1.94	0.67
1:C:141:GLN:O	1:C:145:THR:HG22	1.96	0.66
1:B:123:LEU:HB2	1:B:145:THR:HG21	1.79	0.65
1:C:103:HIS:CD2	2:C:241:GOL:H11	2.34	0.63
1:B:95:GLU:HG3	2:B:241:GOL:H31	1.82	0.62
1:D:103:HIS:HB2	2:D:241:GOL:H31	1.81	0.62
1:B:103:HIS:CD2	2:B:241:GOL:H2	2.34	0.61
1:B:110:LYS:HZ1	2:B:241:GOL:C1	2.04	0.60
1:C:8:ASN:HD21	1:C:50:ARG:HH21	1.49	0.60
1:D:123:LEU:HD13	1:D:142:ILE:HG12	1.87	0.57
1:A:94:ASP:HB3	2:A:241:GOL:O2	2.05	0.56

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:110:LYS:CE	2:B:241:GOL:H12	2.35	0.55
1:B:24:TYR:HB3	1:B:51:LEU:HD11	1.88	0.55
1:A:56:LEU:HD21	1:A:60:PHE:HD2	1.70	0.55
1:A:61:ARG:HG2	1:A:67:LEU:HD21	1.89	0.55
1:B:165:ARG:HG3	1:D:165:ARG:HD2	1.91	0.53
1:B:145:THR:HA	1:B:148:ILE:HD12	1.88	0.53
1:A:110:LYS:HZ2	2:A:241:GOL:H31	1.74	0.53
1:A:101:THR:HG22	3:C:262:HOH:O	2.07	0.53
1:B:110:LYS:HZ2	2:B:241:GOL:C1	2.14	0.52
1:B:161:PHE:HB3	1:B:169:LEU:HD23	1.91	0.52
1:A:91:ASP:HB3	2:A:241:GOL:H12	1.92	0.51
1:D:128:LEU:O	1:D:132:ASP:HB2	2.11	0.51
1:A:40:GLU:HB3	1:A:213:ILE:HG23	1.94	0.50
1:B:61:ARG:HG2	1:B:67:LEU:HD21	1.94	0.50
1:C:123:LEU:CB	1:C:145:THR:HG21	2.40	0.50
1:A:123:LEU:HD13	1:A:142:ILE:HA	1.94	0.49
1:B:32:GLY:HA2	1:B:70:PRO:HG3	1.94	0.49
1:A:56:LEU:HD21	1:A:60:PHE:CD2	2.47	0.49
1:B:123:LEU:HB2	1:B:145:THR:CG2	2.43	0.48
1:D:40:GLU:HB3	1:D:213:ILE:HG23	1.95	0.47
1:C:95:GLU:CG	2:C:241:GOL:H2	2.33	0.47
1:C:238:TYR:CG	1:C:238:TYR:O	2.68	0.47
1:A:110:LYS:HZ1	2:A:241:GOL:C1	2.29	0.46
1:B:14:PRO:HB3	1:B:49:TYR:CZ	2.51	0.46
1:B:7:VAL:HG22	1:B:8:ASN:H	1.80	0.45
1:A:14:PRO:HB3	1:A:49:TYR:CZ	2.51	0.45
1:D:121:ASN:O	1:D:125:LYS:HG3	2.16	0.45
1:B:26:VAL:HG11	1:B:235:SER:HB2	1.99	0.45
1:B:20:LYS:HG2	1:B:45:LYS:HB3	1.99	0.45
1:D:168:MET:O	1:D:172:MET:HG2	2.17	0.44
1:D:225:PHE:CZ	1:D:229:LYS:HD2	2.53	0.44
1:D:123:LEU:HD12	1:D:123:LEU:O	2.18	0.43
1:A:42:LYS:HE3	1:A:42:LYS:HB3	1.79	0.43
1:D:119:ARG:HH11	1:D:148:ILE:CD1	2.31	0.43
1:D:123:LEU:CD1	1:D:142:ILE:HG12	2.48	0.43
1:B:95:GLU:CG	2:B:241:GOL:H31	2.48	0.43
1:A:194:LYS:HA	1:A:197:LEU:HD12	2.01	0.43
1:A:95:GLU:HG3	2:A:241:GOL:C2	2.31	0.43
1:C:30:PRO:HB2	1:C:176:TRP:CZ2	2.54	0.43
1:D:21:LEU:HD22	1:D:74:ILE:HG22	2.00	0.43
1:D:26:VAL:HG21	1:D:240:LEU:HD11	2.00	0.43

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:21:LEU:HG	1:A:74:ILE:HG22	2.00	0.42
1:B:95:GLU:OE1	2:B:241:GOL:H11	2.18	0.42
1:C:94:ASP:HB3	2:C:241:GOL:O2	2.20	0.42
1:D:26:VAL:HG11	1:D:235:SER:HB2	2.02	0.42
1:A:120:PHE:CD1	1:A:173:VAL:HG22	2.53	0.42
1:B:120:PHE:CG	1:B:173:VAL:HG22	2.54	0.42
1:A:11:VAL:HG13	1:A:12:ILE:HG13	2.01	0.42
1:D:119:ARG:HH11	1:D:148:ILE:HD13	1.85	0.42
1:A:72:LEU:HD21	1:A:93:LEU:HD21	2.02	0.41
1:D:108:TYR:CE2	1:D:112:GLN:NE2	2.89	0.41
1:B:103:HIS:HD2	2:B:241:GOL:H2	1.81	0.41
1:B:168:MET:O	1:B:172:MET:HG2	2.21	0.40
1:C:51:LEU:O	1:C:53:PRO:HD3	2.21	0.40
1:D:161:PHE:HB3	1:D:169:LEU:HD23	2.03	0.40
1:A:21:LEU:HD22	1:A:44:ILE:HG21	2.04	0.40

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	228/246 (93%)	224 (98%)	4 (2%)	0	100	100
1	B	228/246 (93%)	222 (97%)	6 (3%)	0	100	100
1	C	224/246 (91%)	217 (97%)	7 (3%)	0	100	100
1	D	240/246 (98%)	236 (98%)	4 (2%)	0	100	100
All	All	920/984 (94%)	899 (98%)	21 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	211/224 (94%)	205 (97%)	6 (3%)	43	56
1	B	212/224 (95%)	207 (98%)	5 (2%)	49	62
1	C	209/224 (93%)	206 (99%)	3 (1%)	67	80
1	D	220/224 (98%)	209 (95%)	11 (5%)	24	30
All	All	852/896 (95%)	827 (97%)	25 (3%)	42	54

All (25) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	27	ASP
1	A	56	LEU
1	A	123	LEU
1	A	173	VAL
1	A	183	LEU
1	A	235	SER
1	B	19	ASP
1	B	56	LEU
1	B	123	LEU
1	B	145	THR
1	B	173	VAL
1	C	119	ARG
1	C	153	LEU
1	C	173	VAL
1	D	-1	HIS
1	D	56	LEU
1	D	61	ARG
1	D	69	ILE
1	D	119	ARG
1	D	123	LEU
1	D	134	ASN
1	D	173	VAL
1	D	212	ASP
1	D	215	LYS
1	D	240	LEU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (5) such sidechains are listed below:

Mol	Chain	Res	Type
1	B	100	HIS
1	C	8	ASN
1	D	112	GLN
1	D	134	ASN
1	D	144	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

4 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	GOL	C	241	-	5,5,5	0.61	0	5,5,5	0.52	0
2	GOL	B	241	-	5,5,5	0.54	0	5,5,5	0.62	0
2	GOL	D	241	-	5,5,5	0.47	0	5,5,5	0.37	0
2	GOL	A	241	-	5,5,5	0.59	0	5,5,5	0.42	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the

Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	GOL	C	241	-	-	2/4/4/4	-
2	GOL	B	241	-	-	4/4/4/4	-
2	GOL	D	241	-	-	2/4/4/4	-
2	GOL	A	241	-	-	2/4/4/4	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (10) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	C	241	GOL	C1-C2-C3-O3
2	B	241	GOL	O1-C1-C2-C3
2	C	241	GOL	O2-C2-C3-O3
2	D	241	GOL	C1-C2-C3-O3
2	A	241	GOL	O1-C1-C2-C3
2	A	241	GOL	O1-C1-C2-O2
2	B	241	GOL	O1-C1-C2-O2
2	B	241	GOL	C1-C2-C3-O3
2	D	241	GOL	O2-C2-C3-O3
2	B	241	GOL	O2-C2-C3-O3

There are no ring outliers.

4 monomers are involved in 28 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	C	241	GOL	6	0
2	B	241	GOL	12	0
2	D	241	GOL	1	0
2	A	241	GOL	9	0

5.7 Other polymers ⓘ

There are no such residues in this entry.

5.8 Polymer linkage issues ⓘ

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	230/246 (93%)	0.05	7 (3%) 50 48	14, 29, 48, 64	0
1	B	232/246 (94%)	0.04	5 (2%) 62 59	16, 28, 49, 60	0
1	C	228/246 (92%)	0.37	21 (9%) 9 7	15, 34, 66, 87	0
1	D	242/246 (98%)	0.15	9 (3%) 41 39	14, 33, 56, 72	0
All	All	932/984 (94%)	0.15	42 (4%) 33 32	14, 31, 56, 87	0

All (42) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	C	238	TYR	10.7
1	C	239	TYR	6.4
1	D	2	THR	5.7
1	C	234	HIS	5.4
1	C	189	ARG	5.3
1	B	54	LEU	4.6
1	C	240	LEU	4.6
1	C	191	PHE	4.4
1	C	7	VAL	3.9
1	C	9	ALA	3.7
1	C	194	LYS	3.7
1	C	55	ARG	3.5
1	D	1	GLY	3.4
1	B	234	HIS	3.3
1	C	184	ARG	3.1
1	D	3	TYR	3.1
1	A	55	ARG	3.0
1	D	189	ARG	3.0
1	A	134	ASN	3.0
1	C	12	ILE	2.9
1	C	183	LEU	2.9

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	136	ALA	2.6
1	B	189	ARG	2.6
1	D	134	ASN	2.6
1	B	232	ARG	2.6
1	D	197	LEU	2.5
1	A	49	TYR	2.5
1	A	183	LEU	2.5
1	C	54	LEU	2.4
1	B	7	VAL	2.4
1	D	-1	HIS	2.4
1	C	227	TYR	2.3
1	D	55	ARG	2.3
1	A	234	HIS	2.2
1	C	49	TYR	2.2
1	C	190	LYS	2.2
1	C	131	PHE	2.2
1	C	226	ASP	2.2
1	C	193	GLU	2.1
1	D	193	GLU	2.1
1	A	194	LYS	2.1
1	C	186	VAL	2.1

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates ⓘ

There are no carbohydrates in this entry.

6.4 Ligands ⓘ

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
2	GOL	D	241	6/6	0.82	0.18	39,39,40,40	0
2	GOL	A	241	6/6	0.84	0.20	34,35,36,36	0

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
2	GOL	C	241	6/6	0.85	0.34	27,29,30,30	0
2	GOL	B	241	6/6	0.88	0.33	28,30,31,32	0

6.5 Other polymers [i](#)

There are no such residues in this entry.