



Full wwPDB X-ray Structure Validation Report ⓘ

Oct 17, 2021 – 12:00 AM EDT

PDB ID : 1RD6
Title : Crystal Structure of S. Marcescens Chitinase A Mutant W167A
Authors : Aronson, N.N.; Halloran, B.A.; Alexyev, M.F.; Zhou, X.E.; Wang, Y.; Meehan, E.J.; Chen, L.
Deposited on : 2003-11-05
Resolution : 2.60 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Xtriage (Phenix)	:	1.13
EDS	:	2.23.2
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac	:	5.8.0158
CCP4	:	7.0.044 (Gargrove)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.23.2


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.60 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	141614	3518 (2.60-2.60)
Ramachandran outliers	138981	3455 (2.60-2.60)
Sidechain outliers	138945	3455 (2.60-2.60)
RSRZ outliers	127900	3104 (2.60-2.60)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	563	<div> <div></div> <div>77%</div> <div>18%</div> <div>..</div> </div>

2 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 4413 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Chitinase A.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	540	Total	C	N	O	S	0	0	0
			4128	2624	696	794	14			

There are 8 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	139	PRO	ALA	SEE REMARK 999	UNP P07254
A	144	SER	PRO	SEE REMARK 999	UNP P07254
A	167	ALA	TRP	engineered mutation	UNP P07254
A	226	ILE	VAL	SEE REMARK 999	UNP P07254
A	395	ALA	PRO	SEE REMARK 999	UNP P07254
A	437	ILE	VAL	SEE REMARK 999	UNP P07254
A	473	GLU	LYS	SEE REMARK 999	UNP P07254
A	484	SER	GLY	SEE REMARK 999	UNP P07254


- Molecule 2 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	285	Total	O	0	0
			285	285		

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Chitinase A



4 Data and refinement statistics

Property	Value	Source
Space group	I 2 2 2	Depositor
Cell constants a, b, c, α , β , γ	77.59Å 134.06Å 187.90Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	19.79 – 2.60 48.97 – 2.60	Depositor EDS
% Data completeness (in resolution range)	95.8 (19.79-2.60) 95.7 (48.97-2.60)	Depositor EDS
R_{merge}	0.09	Depositor
R_{sym}	0.09	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.32 (at 2.61Å)	Xtriage
Refinement program	CNS 1.0	Depositor
R, R_{free}	0.195 , 0.235 0.209 , (Not available)	Depositor DCC
R_{free} test set	No test flags present.	wwPDB-VP
Wilson B-factor (Å ²)	34.5	Xtriage
Anisotropy	0.585	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.33 , 46.9	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	4413	wwPDB-VP
Average B, all atoms (Å ²)	35.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.35% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.37	0/4227	0.59	0/5735

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4128	0	4006	77	0
2	A	285	0	0	6	0
All	All	4413	0	4006	77	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 9.

All (77) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:528:VAL:HG13	1:A:533:LEU:HB2	1.62	0.81
1:A:43:VAL:HG22	1:A:54:VAL:HG12	1.65	0.78
1:A:191:PHE:HA	1:A:271:SER:O	1.96	0.65
1:A:54:VAL:HG11	1:A:304:TRP:CZ2	2.32	0.65

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:163:TYR:O	1:A:540:GLU:HA	1.98	0.63
1:A:541:ILE:HD11	1:A:549:LEU:HD22	1.81	0.62
1:A:54:VAL:HG11	1:A:304:TRP:CH2	2.34	0.62
1:A:164:PHE:O	1:A:190:GLY:N	2.33	0.62
1:A:361:SER:HB3	1:A:382:MET:HE3	1.82	0.61
1:A:153:LYS:HG2	1:A:563:GLN:HB3	1.85	0.57
1:A:29:PRO:HG2	1:A:123:SER:HB3	1.86	0.57
1:A:164:PHE:CD2	1:A:179:ILE:HD11	2.39	0.57
1:A:143:GLU:HB3	2:A:790:HOH:O	2.04	0.56
1:A:524:LYS:O	1:A:528:VAL:HG23	2.07	0.55
1:A:165:VAL:HA	1:A:190:GLY:HA3	1.88	0.55
1:A:319:GLY:O	1:A:320:LYS:HB2	2.07	0.53
1:A:361:SER:HB3	1:A:382:MET:CE	2.37	0.53
1:A:407:LEU:O	1:A:421:VAL:HG23	2.08	0.53
1:A:260:LYS:HG2	1:A:268:ILE:HD12	1.91	0.53
1:A:403:HIS:CB	1:A:406:ALA:HB2	2.40	0.52
1:A:204:LEU:HD11	1:A:277:LEU:HD21	1.90	0.52
1:A:540:GLU:HB2	1:A:542:ASP:OD1	2.10	0.52
1:A:189:TYR:O	1:A:191:PHE:N	2.35	0.52
1:A:441:THR:HG22	1:A:524:LYS:HD2	1.93	0.51
1:A:339:MET:HE1	1:A:342:LEU:HD12	1.93	0.51
1:A:492:GLN:O	1:A:504:VAL:HA	2.11	0.51
1:A:317:PRO:HD2	2:A:719:HOH:O	2.12	0.50
1:A:193:PRO:HD2	1:A:227:SER:O	2.12	0.49
1:A:166:GLU:H	1:A:190:GLY:HA3	1.78	0.49
1:A:244:ALA:HB3	1:A:247:ASP:OD2	2.12	0.49
1:A:421:VAL:HG12	1:A:425:ASN:ND2	2.28	0.48
1:A:315:GLU:HB3	2:A:611:HOH:O	2.13	0.47
1:A:226:ILE:HD11	1:A:272:ILE:HD13	1.96	0.47
1:A:443:MET:HG2	1:A:548:ILE:HG21	1.96	0.47
1:A:279:ASP:HB2	2:A:601:HOH:O	2.14	0.47
1:A:483:ALA:HA	1:A:487:MET:HG3	1.96	0.47
1:A:141:LEU:HB2	1:A:258:ALA:HB1	1.96	0.47
1:A:107:ARG:CG	1:A:107:ARG:HH11	2.28	0.46
1:A:110:MET:C	1:A:110:MET:SD	2.93	0.46
1:A:163:TYR:CZ	1:A:188:LEU:HD22	2.51	0.46
1:A:226:ILE:HD13	1:A:226:ILE:HA	1.77	0.45
1:A:312:ILE:HG13	1:A:339:MET:CE	2.46	0.45
1:A:275:TRP:HE3	1:A:315:GLU:HG2	1.81	0.45
1:A:153:LYS:O	1:A:558:ASN:HB3	2.16	0.45
1:A:226:ILE:HD11	1:A:293:PHE:HE1	1.82	0.45

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:366:GLY:O	1:A:370:ILE:HG13	2.17	0.45
1:A:211:PHE:O	1:A:215:GLN:HG2	2.16	0.44
1:A:244:ALA:HB3	1:A:247:ASP:CG	2.38	0.44
1:A:31:ILE:HD13	1:A:110:MET:CE	2.48	0.44
1:A:273:GLY:O	1:A:277:LEU:HB2	2.18	0.44
1:A:443:MET:HG3	1:A:538:SER:HB2	1.99	0.44
1:A:270:PRO:HD3	1:A:307:PHE:CD1	2.53	0.44
1:A:410:PRO:HD3	2:A:670:HOH:O	2.17	0.44
1:A:477:VAL:HG21	1:A:513:ILE:HD13	2.00	0.44
1:A:314:TRP:O	1:A:316:PHE:N	2.52	0.43
1:A:316:PHE:CD1	1:A:364:SER:HB2	2.52	0.43
1:A:165:VAL:HA	1:A:190:GLY:CA	2.49	0.43
1:A:279:ASP:HB2	1:A:280:PRO:HD3	2.01	0.43
1:A:80:LEU:HD23	1:A:85:GLU:HA	2.01	0.42
1:A:450:GLY:O	1:A:465:THR:HG23	2.19	0.42
1:A:102:VAL:HG11	1:A:130:VAL:HG21	2.01	0.42
1:A:128:ILE:O	1:A:128:ILE:HG13	2.19	0.42
1:A:548:ILE:O	1:A:552:MET:HG3	2.18	0.42
1:A:316:PHE:CD1	1:A:316:PHE:N	2.85	0.42
1:A:403:HIS:HB3	1:A:406:ALA:HB2	2.02	0.42
1:A:204:LEU:HD22	1:A:210:SER:OG	2.19	0.42
1:A:43:VAL:HA	1:A:53:LEU:O	2.19	0.42
1:A:448:TRP:HB3	1:A:464:ALA:CB	2.50	0.42
1:A:84:LYS:HD2	2:A:662:HOH:O	2.20	0.41
1:A:107:ARG:NH1	1:A:107:ARG:HG2	2.35	0.41
1:A:434:PRO:O	1:A:533:LEU:HA	2.20	0.41
1:A:439:VAL:O	1:A:536:LEU:HA	2.21	0.41
1:A:200:ILE:HG22	1:A:280:PRO:HG3	2.03	0.40
1:A:271:SER:HA	1:A:311:ASP:HB3	2.03	0.40
1:A:408:ASN:HB2	1:A:499:ALA:HA	2.03	0.40
1:A:131:ALA:CB	1:A:248:PRO:HG3	2.51	0.40
1:A:471:THR:OG1	1:A:477:VAL:HA	2.22	0.40

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries

of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	538/563 (96%)	507 (94%)	25 (5%)	6 (1%)	14	30

All (6) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	540	GLU
1	A	190	GLY
1	A	315	GLU
1	A	316	PHE
1	A	561	GLY
1	A	191	PHE

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	429/447 (96%)	421 (98%)	8 (2%)	57	79

All (8) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	30	THR
1	A	107	ARG
1	A	284	MET
1	A	315	GLU
1	A	316	PHE
1	A	455	GLN
1	A	484	SER
1	A	516	ASP

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (5) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	52	ASN
1	A	182	GLN
1	A	330	GLN
1	A	425	ASN
1	A	485	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	540/563 (95%)	-0.36	7 (1%) 77 73	19, 34, 49, 102	21 (3%)

All (7) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	563	GLN	10.5
1	A	562	VAL	7.4
1	A	24	ALA	2.9
1	A	188	LEU	2.8
1	A	117	ALA	2.7
1	A	561	GLY	2.5
1	A	493	TYR	2.4

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.