



wwPDB X-ray Structure Validation Summary Report ⓘ

May 24, 2021 – 04:25 PM EDT

PDB ID : 4RFO
Title : Crystal structure of the ADCC-Potent Antibody N60-I3 Fab in complex with HIV-1 Clade A/E gp120 and M48u1
Authors : Tolbert, W.D.; Gohain, N.; Pazgier, M.
Deposited on : 2014-09-26
Resolution : 3.20 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix)	:	1.13
EDS	:	2.18
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac	:	5.8.0158
CCP4	:	7.0.044 (Gargrove)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.18


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 3.20 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	1133 (3.20-3.20)
Clashscore	141614	1253 (3.20-3.20)
Ramachandran outliers	138981	1234 (3.20-3.20)
Sidechain outliers	138945	1233 (3.20-3.20)
RSRZ outliers	127900	1095 (3.20-3.20)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	G	353	<div> <div>3%</div> <div>57% 31% 6% 5%</div> </div>
2	N	28	<div> <div>71% 21% . .</div> </div>
3	H	229	<div> <div>3%</div> <div>50% 30% 10% 7%</div> </div>
4	L	221	<div> <div>3%</div> <div>59% 30% 6% . .</div> </div>

2 Entry composition

There are 6 unique types of molecules in this entry. The entry contains 6169 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called HIV-1 clade A/E gp120.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	G	336	Total	C	N	O	S	0	0	0
			2640	1657	458	503	22			

- Molecule 2 is a protein called m48u1 CD4 mimetic peptide.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	N	28	Total	C	N	O	S	0	0	1
			209	133	38	32	6			

- Molecule 3 is a protein called N60-i3 Fab heavy chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	H	214	Total	C	N	O	S	0	0	0
			1614	1027	269	314	4			

- Molecule 4 is a protein called N60-i3 Fab light chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
4	L	212	Total	C	N	O	S	0	0	0
			1579	986	261	327	5			

- Molecule 5 is 2-acetamido-2-deoxy-beta-D-glucopyranose (three-letter code: NAG) (formula: C₈H₁₅NO₆).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
5	G	1	Total	C	N	O	0	0
			14	8	1	5		
5	G	1	Total	C	N	O	0	0
			14	8	1	5		
5	G	1	Total	C	N	O	0	0
			14	8	1	5		
5	G	1	Total	C	N	O	0	0
			14	8	1	5		
5	G	1	Total	C	N	O	0	0
			14	8	1	5		
5	G	1	Total	C	N	O	0	0
			14	8	1	5		
5	G	1	Total	C	N	O	0	0
			14	8	1	5		


- Molecule 6 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	H	1	Total	O	0	0
			1	1		


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: HIV-1 clade A/E gp120




- Molecule 2: m48u1 CD4 mimetic peptide



- Molecule 3: N60-i3 Fab heavy chain



● Molecule 4: N60-i3 Fab light chain



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 2	Depositor
Cell constants a, b, c, α , β , γ	98.31Å 102.58Å 108.04Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	36.01 – 3.20 36.01 – 3.19	Depositor EDS
% Data completeness (in resolution range)	93.8 (36.01-3.20) 93.7 (36.01-3.19)	Depositor EDS
R_{merge}	0.13	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.27 (at 3.18Å)	Xtriage
Refinement program	REFMAC 5.8.0073/Phenix 1.9	Depositor
R, R_{free}	0.220 , 0.277 0.221 , 0.275	Depositor DCC
R_{free} test set	879 reflections (5.02%)	wwPDB-VP
Wilson B-factor (Å ²)	102.1	Xtriage
Anisotropy	0.347	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.22 , 43.9	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	0.025 for k,h,-l	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	6169	wwPDB-VP
Average B, all atoms (Å ²)	116.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.92% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: U2X, NAG, NH2, DPR, MPT

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	G	0.41	1/2694 (0.0%)	0.62	0/3655
2	N	0.43	0/176	0.58	0/231
3	H	0.47	0/1655	0.81	4/2263 (0.2%)
4	L	0.41	0/1618	0.64	0/2208
All	All	0.43	1/6143 (0.0%)	0.68	4/8357 (0.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	G	0	1
2	N	0	2
3	H	0	4
All	All	0	7

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	G	355	ASN	C-N	7.22	1.50	1.34

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	H	100	GLY	C-N-CA	7.95	139.00	122.30
3	H	100(C)	TYR	O-C-N	-7.57	110.59	122.70
3	H	100(C)	TYR	CA-C-N	5.93	130.25	117.20
3	H	100	GLY	O-C-N	-5.33	114.13	123.20

There are no chirality outliers.

5 of 7 planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	G	217	TYR	Peptide
3	H	23	THR	Peptide
3	H	29	ILE	Peptide
2	N	22	THR	Mainchain
2	N	23	U2X	Mainchain

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	G	2640	0	2568	92	0
2	N	209	0	212	4	0
3	H	1614	0	1590	91	0
4	L	1579	0	1524	53	0
5	G	126	0	117	2	0
6	H	1	0	0	0	0
All	All	6169	0	6011	233	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 19.

The worst 5 of 233 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
4:L:167:GLN:HG3	4:L:171:LYS:O	1.59	1.01
3:H:34:TYR:CD1	3:H:96:PRO:HB3	1.98	0.98
3:H:171:GLN:NE2	3:H:177:SER:OG	1.96	0.98
3:H:72:ASP:O	3:H:75:GLN:N	2.00	0.94
3:H:72:ASP:O	3:H:75:GLN:CA	2.19	0.91

There are no symmetry-related clashes.

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	G	328/353 (93%)	271 (83%)	29 (9%)	28 (8%)	1	4
2	N	24/28 (86%)	20 (83%)	3 (12%)	1 (4%)	3	20
3	H	210/229 (92%)	160 (76%)	25 (12%)	25 (12%)	0	2
4	L	210/221 (95%)	164 (78%)	31 (15%)	15 (7%)	1	8
All	All	772/831 (93%)	615 (80%)	88 (11%)	69 (9%)	1	4

5 of 69 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	G	87	GLU
1	G	114	GLN
1	G	121	LYS
1	G	235	GLY
1	G	345	VAL

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	G	302/311 (97%)	270 (89%)	32 (11%)	6	27
2	N	20/20 (100%)	18 (90%)	2 (10%)	7	30
3	H	184/197 (93%)	158 (86%)	26 (14%)	3	16
4	L	180/188 (96%)	160 (89%)	20 (11%)	6	25
All	All	686/716 (96%)	606 (88%)	80 (12%)	5	23

5 of 80 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
3	H	192	GLN
4	L	145	THR
3	H	208	ASP
4	L	38	GLN
4	L	161	THR

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 6 such sidechains are listed below:

Mol	Chain	Res	Type
1	G	490	GLN
3	H	171	GLN
4	L	60	ASN
1	G	280	ASN
1	G	258	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

2 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	U2X	N	23	2	19,20,21	0.33	0	22,25,27	0.67	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	U2X	N	23	2	-	6/10/19/21	0/2/2/2

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

5 of 6 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	N	23	U2X	C4-C3-C7-OH
2	N	23	U2X	CE1-CZ-OH-C7
2	N	23	U2X	CE2-CZ-OH-C7
2	N	23	U2X	C2-C3-C7-OH
2	N	23	U2X	CA-CB-CG-CD1

There are no ring outliers.

No monomer is involved in short contacts.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

9 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
5	NAG	G	501	1	14,14,15	0.50	0	17,19,21	1.26	2 (11%)
5	NAG	G	505	1	14,14,15	0.50	0	17,19,21	2.39	6 (35%)
5	NAG	G	508	1	14,14,15	0.61	0	17,19,21	1.40	2 (11%)
5	NAG	G	503	1	14,14,15	0.98	1 (7%)	17,19,21	2.80	4 (23%)
5	NAG	G	502	1	14,14,15	0.57	0	17,19,21	1.13	2 (11%)

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
5	NAG	G	509	1	14,14,15	0.78	1 (7%)	17,19,21	1.29	3 (17%)
5	NAG	G	504	1	14,14,15	0.60	0	17,19,21	1.56	2 (11%)
5	NAG	G	507	1	14,14,15	0.57	0	17,19,21	1.07	2 (11%)
5	NAG	G	506	1	14,14,15	0.60	0	17,19,21	1.12	2 (11%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	NAG	G	501	1	-	2/6/23/26	0/1/1/1
5	NAG	G	505	1	-	4/6/23/26	0/1/1/1
5	NAG	G	508	1	-	2/6/23/26	0/1/1/1
5	NAG	G	503	1	-	0/6/23/26	0/1/1/1
5	NAG	G	502	1	-	1/6/23/26	0/1/1/1
5	NAG	G	509	1	-	2/6/23/26	0/1/1/1
5	NAG	G	504	1	-	0/6/23/26	0/1/1/1
5	NAG	G	507	1	-	2/6/23/26	0/1/1/1
5	NAG	G	506	1	-	0/6/23/26	0/1/1/1

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
5	G	503	NAG	C1-C2	2.74	1.56	1.52
5	G	509	NAG	C1-C2	2.33	1.55	1.52

The worst 5 of 25 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
5	G	503	NAG	C1-O5-C5	9.74	125.39	112.19
5	G	505	NAG	C2-N2-C7	5.77	131.12	122.90
5	G	504	NAG	C4-C3-C2	4.72	117.93	111.02
5	G	505	NAG	O5-C1-C2	-4.56	104.09	111.29
5	G	508	NAG	O5-C1-C2	-4.33	104.45	111.29

There are no chirality outliers.

5 of 13 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
5	G	505	NAG	C3-C2-N2-C7
5	G	508	NAG	O5-C5-C6-O6
5	G	509	NAG	O5-C5-C6-O6
5	G	509	NAG	C4-C5-C6-O6
5	G	508	NAG	C4-C5-C6-O6

There are no ring outliers.

1 monomer is involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
5	G	503	NAG	2	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	G	336/353 (95%)	-0.47	2 (0%) 89 83	61, 105, 180, 200	0
2	N	24/28 (85%)	-0.66	0 100 100	84, 97, 102, 119	0
3	H	214/229 (93%)	-0.35	7 (3%) 46 30	66, 116, 171, 210	0
4	L	212/221 (95%)	-0.30	7 (3%) 46 30	76, 136, 179, 210	0
All	All	786/831 (94%)	-0.40	16 (2%) 65 51	61, 113, 179, 210	0

The worst 5 of 16 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
3	H	31	SER	4.3
1	G	123	THR	3.6
4	L	144	VAL	3.5
4	L	142	GLY	3.4
3	H	127	SER	3.3

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q < 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	DPR	N	21	7/8	0.93	0.23	87,89,93,95	0
2	U2X	N	23	19/20	0.98	0.21	78,79,86,86	0

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
5	NAG	G	509	14/15	0.89	0.26	145,157,162,167	0
5	NAG	G	503	14/15	0.91	0.21	76,94,101,104	0
5	NAG	G	504	14/15	0.91	0.14	113,125,129,130	0
5	NAG	G	502	14/15	0.91	0.26	116,120,129,143	0
5	NAG	G	508	14/15	0.92	0.17	113,118,126,134	0
5	NAG	G	505	14/15	0.93	0.14	100,112,121,122	0
5	NAG	G	506	14/15	0.93	0.16	100,115,122,126	0
5	NAG	G	501	14/15	0.94	0.10	99,109,124,125	0
5	NAG	G	507	14/15	0.94	0.32	114,121,127,131	0

6.5 Other polymers [i](#)

There are no such residues in this entry.