



wwPDB X-ray Structure Validation Summary Report ⓘ

Jun 15, 2020 – 03:54 am BST

PDB ID : 3RHP
Title : Crystal structure of the C707A mutant of the C-Terminal domain of 10'FOR MYLTETRAHYDROFOLATE DEHYDROGENASE
Authors : Tsybovsky, Y.
Deposited on : 2011-04-11
Resolution : 2.50 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.11
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.11


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.50 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	4661 (2.50-2.50)
Clashscore	141614	5346 (2.50-2.50)
Ramachandran outliers	138981	5231 (2.50-2.50)
Sidechain outliers	138945	5233 (2.50-2.50)
RSRZ outliers	127900	4559 (2.50-2.50)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	517	<div> <div>2%</div> <div> <div></div> <div>75%</div> <div>19%</div> <div>• •</div> </div> </div>
1	B	517	<div> <div>%</div> <div> <div></div> <div>76%</div> <div>19%</div> <div>• •</div> </div> </div>
1	C	517	<div> <div>%</div> <div> <div></div> <div>77%</div> <div>18%</div> <div>• •</div> </div> </div>
1	D	517	<div> <div>%</div> <div> <div></div> <div>77%</div> <div>17%</div> <div>• •</div> </div> </div>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit crite-

ria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	SO4	C	3	-	-	X	-

2 Entry composition [i](#)

There are 4 unique types of molecules in this entry. The entry contains 16351 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Aldehyde dehydrogenase 1 family, member L1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	498	Total	C	N	O	S	0	12	0
			3867	2461	659	727	20			
1	B	498	Total	C	N	O	S	0	11	0
			3871	2461	664	726	20			
1	C	498	Total	C	N	O	S	0	11	0
			3866	2457	669	720	20			
1	D	498	Total	C	N	O	S	0	13	0
			3889	2473	667	729	20			

There are 48 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	386	MET	-	EXPRESSION TAG	UNP Q5HQB2
A	387	ARG	-	EXPRESSION TAG	UNP Q5HQB2
A	388	GLY	-	EXPRESSION TAG	UNP Q5HQB2
A	389	SER	-	EXPRESSION TAG	UNP Q5HQB2
A	390	HIS	-	EXPRESSION TAG	UNP Q5HQB2
A	391	HIS	-	EXPRESSION TAG	UNP Q5HQB2
A	392	HIS	-	EXPRESSION TAG	UNP Q5HQB2
A	393	HIS	-	EXPRESSION TAG	UNP Q5HQB2
A	394	HIS	-	EXPRESSION TAG	UNP Q5HQB2
A	395	THR	-	EXPRESSION TAG	UNP Q5HQB2
A	396	THR	-	EXPRESSION TAG	UNP Q5HQB2
A	707	ALA	CYS	ENGINEERED MUTATION	UNP Q5HQB2
B	386	MET	-	EXPRESSION TAG	UNP Q5HQB2
B	387	ARG	-	EXPRESSION TAG	UNP Q5HQB2
B	388	GLY	-	EXPRESSION TAG	UNP Q5HQB2
B	389	SER	-	EXPRESSION TAG	UNP Q5HQB2
B	390	HIS	-	EXPRESSION TAG	UNP Q5HQB2
B	391	HIS	-	EXPRESSION TAG	UNP Q5HQB2
B	392	HIS	-	EXPRESSION TAG	UNP Q5HQB2
B	393	HIS	-	EXPRESSION TAG	UNP Q5HQB2
B	394	HIS	-	EXPRESSION TAG	UNP Q5HQB2

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
B	395	THR	-	EXPRESSION TAG	UNP Q5HQB2
B	396	THR	-	EXPRESSION TAG	UNP Q5HQB2
B	707	ALA	CYS	ENGINEERED MUTATION	UNP Q5HQB2
C	386	MET	-	EXPRESSION TAG	UNP Q5HQB2
C	387	ARG	-	EXPRESSION TAG	UNP Q5HQB2
C	388	GLY	-	EXPRESSION TAG	UNP Q5HQB2
C	389	SER	-	EXPRESSION TAG	UNP Q5HQB2
C	390	HIS	-	EXPRESSION TAG	UNP Q5HQB2
C	391	HIS	-	EXPRESSION TAG	UNP Q5HQB2
C	392	HIS	-	EXPRESSION TAG	UNP Q5HQB2
C	393	HIS	-	EXPRESSION TAG	UNP Q5HQB2
C	394	HIS	-	EXPRESSION TAG	UNP Q5HQB2
C	395	THR	-	EXPRESSION TAG	UNP Q5HQB2
C	396	THR	-	EXPRESSION TAG	UNP Q5HQB2
C	707	ALA	CYS	ENGINEERED MUTATION	UNP Q5HQB2
D	386	MET	-	EXPRESSION TAG	UNP Q5HQB2
D	387	ARG	-	EXPRESSION TAG	UNP Q5HQB2
D	388	GLY	-	EXPRESSION TAG	UNP Q5HQB2
D	389	SER	-	EXPRESSION TAG	UNP Q5HQB2
D	390	HIS	-	EXPRESSION TAG	UNP Q5HQB2
D	391	HIS	-	EXPRESSION TAG	UNP Q5HQB2
D	392	HIS	-	EXPRESSION TAG	UNP Q5HQB2
D	393	HIS	-	EXPRESSION TAG	UNP Q5HQB2
D	394	HIS	-	EXPRESSION TAG	UNP Q5HQB2
D	395	THR	-	EXPRESSION TAG	UNP Q5HQB2
D	396	THR	-	EXPRESSION TAG	UNP Q5HQB2
D	707	ALA	CYS	ENGINEERED MUTATION	UNP Q5HQB2

- Molecule 2 is SULFATE ION (three-letter code: SO4) (formula: O₄S).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	O	S	0	0
			5	4	1		
2	A	1	Total	O	S	0	0
			5	4	1		
2	A	1	Total	O	S	0	0
			5	4	1		
2	A	1	Total	O	S	0	0
			5	4	1		
2	A	1	Total	O	S	0	0
			5	4	1		
2	B	1	Total	O	S	0	0
			5	4	1		
2	B	1	Total	O	S	0	0
			5	4	1		
2	B	1	Total	O	S	0	0
			5	4	1		
2	B	1	Total	O	S	0	0
			5	4	1		
2	B	1	Total	O	S	0	0
			5	4	1		
2	C	1	Total	O	S	0	0
			5	4	1		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	C	1	Total	O	S	0	0
			5	4	1		
2	C	1	Total	O	S	0	0
			5	4	1		
2	C	1	Total	O	S	0	0
			5	4	1		
2	C	1	Total	O	S	0	0
			5	4	1		
2	C	1	Total	O	S	0	0
			5	4	1		
2	D	1	Total	O	S	0	0
			5	4	1		
2	D	1	Total	O	S	0	0
			5	4	1		
2	D	1	Total	O	S	0	0
			5	4	1		

- Molecule 3 is GLYCEROL (three-letter code: GOL) (formula: $C_3H_8O_3$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	C	O	0	0
			6	3	3		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	B	1	Total	C	O	0	0
			6	3	3		
3	C	1	Total	C	O	0	0
			6	3	3		
3	D	1	Total	C	O	0	0
			6	3	3		


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	176	Total	O	0	0
			176	176		
4	B	170	Total	O	0	0
			170	170		
4	C	185	Total	O	0	0
			185	185		
4	D	183	Total	O	0	0
			183	183		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Aldehyde dehydrogenase 1 family, member L1




- Molecule 1: Aldehyde dehydrogenase 1 family, member L1





- Molecule 1: Aldehyde dehydrogenase 1 family, member L1





● Molecule 1: Aldehyde dehydrogenase 1 family, member L1



4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, α , β , γ	261.40 Å 194.90 Å 97.11 Å 90.00° 108.70° 90.00°	Depositor
Resolution (Å)	48.97 – 2.50 48.94 – 2.50	Depositor EDS
% Data completeness (in resolution range)	98.0 (48.97-2.50) 98.0 (48.94-2.50)	Depositor EDS
R_{merge}	0.15	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.95 (at 2.51 Å)	Xtriage
Refinement program	REFMAC 5.5.0102	Depositor
R, R_{free}	0.213 , 0.257 0.214 , 0.256	Depositor DCC
R_{free} test set	7861 reflections (5.05%)	wwPDB-VP
Wilson B-factor (Å ²)	33.7	Xtriage
Anisotropy	0.191	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.35 , 44.6	EDS
L-test for twinning ²	$\langle L \rangle = 0.51$, $\langle L^2 \rangle = 0.35$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.93	EDS
Total number of atoms	16351	wwPDB-VP
Average B, all atoms (Å ²)	31.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.51% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: GOL, SO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	1.07	4/3985 (0.1%)	1.00	4/5394 (0.1%)
1	B	1.04	1/3993 (0.0%)	0.98	6/5403 (0.1%)
1	C	1.14	2/3991 (0.1%)	1.05	8/5400 (0.1%)
1	D	1.08	3/4015 (0.1%)	1.01	8/5430 (0.1%)
All	All	1.08	10/15984 (0.1%)	1.01	26/21627 (0.1%)

The worst 5 of 10 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	D	742	ARG	CG-CD	6.56	1.68	1.51
1	C	831	GLU	CG-CD	6.39	1.61	1.51
1	B	827	ALA	CA-CB	-6.36	1.39	1.52
1	D	738	ASN	CB-CG	6.30	1.65	1.51
1	A	491	ARG	CB-CG	5.70	1.68	1.52

The worst 5 of 26 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	D	485	ARG	NE-CZ-NH2	-9.13	115.73	120.30
1	D	826	ARG	NE-CZ-NH2	-7.41	116.59	120.30
1	C	484	ASP	CB-CG-OD2	7.34	124.91	118.30
1	C	687	ASP	CB-CG-OD2	7.21	124.79	118.30
1	A	503	LEU	CB-CG-CD1	-6.64	99.71	111.00

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3867	0	3861	59	0
1	B	3871	0	3869	69	0
1	C	3866	0	3867	61	0
1	D	3889	0	3898	57	0
2	A	30	0	0	0	0
2	B	35	0	0	0	0
2	C	35	0	0	2	0
2	D	20	0	0	1	0
3	A	6	0	8	0	0
3	B	6	0	8	1	0
3	C	6	0	8	1	0
3	D	6	0	8	0	0
4	A	176	0	0	2	0
4	B	170	0	0	6	0
4	C	185	0	0	7	0
4	D	183	0	0	8	0
All	All	16351	0	15527	234	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 7.

The worst 5 of 234 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:690:LYS:HE3	1:B:694[B]:MET:CE	1.29	1.56
1:B:690:LYS:CE	1:B:694[B]:MET:CE	1.99	1.37
1:B:690:LYS:CE	1:B:694[B]:MET:HE1	1.72	1.12
1:B:690:LYS:HE3	1:B:694[B]:MET:SD	1.94	1.06
1:D:756:ARG:NH2	4:D:1159:HOH:O	1.88	1.04

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	507/517 (98%)	484 (96%)	23 (4%)	0	100	100
1	B	507/517 (98%)	477 (94%)	29 (6%)	1 (0%)	47	68
1	C	507/517 (98%)	479 (94%)	27 (5%)	1 (0%)	47	68
1	D	509/517 (98%)	470 (92%)	36 (7%)	3 (1%)	25	43
All	All	2030/2068 (98%)	1910 (94%)	115 (6%)	5 (0%)	41	68

All (5) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	C	436	SER
1	D	524	GLY
1	B	436	SER
1	D	703	LYS
1	D	833	GLY

5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	415/425 (98%)	391 (94%)	24 (6%)	20	38
1	B	416/425 (98%)	390 (94%)	26 (6%)	18	34
1	C	415/425 (98%)	401 (97%)	14 (3%)	37	63
1	D	419/425 (99%)	399 (95%)	20 (5%)	25	48
All	All	1665/1700 (98%)	1581 (95%)	84 (5%)	24	46

5 of 84 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	690	LYS
1	B	825	SER
1	D	728	VAL
1	B	694[B]	MET
1	B	741[A]	GLU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 20 such sidechains are listed below:

Mol	Chain	Res	Type
1	B	844	ASN
1	C	541	GLN
1	D	600	GLN
1	B	706	ASN
1	B	745	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

5.6 Ligand geometry ⓘ

28 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	SO4	A	2	-	4,4,4	0.33	0	6,6,6	0.80	0
2	SO4	C	5	-	4,4,4	0.09	0	6,6,6	0.65	0
2	SO4	A	5	-	4,4,4	0.42	0	6,6,6	1.41	0
2	SO4	A	1	-	4,4,4	0.17	0	6,6,6	0.80	0
2	SO4	C	7	-	4,4,4	0.41	0	6,6,6	0.78	0
3	GOL	C	8	-	5,5,5	0.62	0	5,5,5	2.07	2 (40%)
3	GOL	B	10	-	5,5,5	0.37	0	5,5,5	1.87	1 (20%)
3	GOL	A	7	-	5,5,5	0.33	0	5,5,5	1.45	1 (20%)
2	SO4	C	4	-	4,4,4	0.28	0	6,6,6	0.60	0
2	SO4	C	1	-	4,4,4	0.23	0	6,6,6	0.59	0
2	SO4	C	2	-	4,4,4	0.20	0	6,6,6	0.62	0
2	SO4	B	7	-	4,4,4	0.09	0	6,6,6	0.85	0
2	SO4	D	4	-	4,4,4	0.14	0	6,6,6	0.57	0
2	SO4	B	8	-	4,4,4	0.21	0	6,6,6	0.60	0
2	SO4	C	6	-	4,4,4	0.19	0	6,6,6	1.10	0
2	SO4	D	1	-	4,4,4	0.33	0	6,6,6	0.58	0
2	SO4	B	9	-	4,4,4	0.27	0	6,6,6	0.63	0
2	SO4	B	3	-	4,4,4	0.21	0	6,6,6	0.59	0
3	GOL	D	5	-	5,5,5	0.65	0	5,5,5	1.15	0
2	SO4	C	3	-	4,4,4	0.13	0	6,6,6	0.45	0
2	SO4	B	4	-	4,4,4	0.16	0	6,6,6	0.56	0
2	SO4	A	6	-	4,4,4	0.22	0	6,6,6	1.51	1 (16%)
2	SO4	B	5	-	4,4,4	0.33	0	6,6,6	0.67	0
2	SO4	D	2	-	4,4,4	0.46	0	6,6,6	0.59	0
2	SO4	B	6	-	4,4,4	0.23	0	6,6,6	0.52	0
2	SO4	D	3	-	4,4,4	0.18	0	6,6,6	0.79	0
2	SO4	A	3	-	4,4,4	0.40	0	6,6,6	0.69	0
2	SO4	A	4	-	4,4,4	0.20	0	6,6,6	0.88	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	GOL	C	8	-	-	3/4/4/4	-
3	GOL	A	7	-	-	2/4/4/4	-
3	GOL	D	5	-	-	2/4/4/4	-
3	GOL	B	10	-	-	2/4/4/4	-

There are no bond length outliers.

All (5) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	B	10	GOL	C3-C2-C1	-3.80	96.95	111.70
3	C	8	GOL	C3-C2-C1	-3.39	98.51	111.70
3	A	7	GOL	C3-C2-C1	-2.95	100.25	111.70
2	A	6	SO4	O4-S-O3	2.80	121.02	109.06
3	C	8	GOL	O2-C2-C3	2.01	118.00	109.12

There are no chirality outliers.

5 of 9 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	C	8	GOL	C1-C2-C3-O3
3	B	10	GOL	C1-C2-C3-O3
3	A	7	GOL	C1-C2-C3-O3
3	D	5	GOL	O1-C1-C2-C3
3	D	5	GOL	O1-C1-C2-O2

There are no ring outliers.

4 monomers are involved in 5 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	C	8	GOL	1	0
3	B	10	GOL	1	0
2	C	3	SO4	2	0
2	D	2	SO4	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	498/517 (96%)	-0.07	8 (1%) 72 74	16, 31, 46, 57	19 (3%)
1	B	498/517 (96%)	-0.06	6 (1%) 79 80	19, 32, 46, 62	20 (4%)
1	C	498/517 (96%)	-0.23	6 (1%) 79 80	15, 28, 39, 53	16 (3%)
1	D	498/517 (96%)	-0.21	4 (0%) 86 87	16, 31, 44, 62	19 (3%)
All	All	1992/2068 (96%)	-0.15	24 (1%) 79 80	15, 30, 44, 62	74 (3%)

The worst 5 of 24 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	C	435	GLY	3.2
1	B	436	SER	3.0
1	D	738	ASN	2.8
1	A	764	ARG	2.7
1	B	553	ASN	2.6

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
2	SO4	B	7	5/5	0.81	0.34	35,36,39,39	5
2	SO4	A	5	5/5	0.88	0.41	55,56,60,60	5
2	SO4	B	8	5/5	0.88	0.22	30,32,35,36	5
2	SO4	A	4	5/5	0.89	0.21	33,35,36,39	5
3	GOL	D	5	6/6	0.90	0.24	29,34,43,49	0
2	SO4	B	9	5/5	0.90	0.20	33,33,36,36	5
2	SO4	D	3	5/5	0.91	0.21	34,35,36,37	5
2	SO4	C	4	5/5	0.92	0.21	33,33,35,35	5
2	SO4	D	2	5/5	0.92	0.21	40,40,41,42	5
2	SO4	B	6	5/5	0.93	0.16	42,42,43,44	5
2	SO4	C	6	5/5	0.93	0.32	51,54,56,57	5
3	GOL	C	8	6/6	0.93	0.25	34,34,36,39	0
2	SO4	D	1	5/5	0.94	0.14	34,38,39,42	5
3	GOL	B	10	6/6	0.94	0.25	36,39,43,50	0
3	GOL	A	7	6/6	0.94	0.28	39,40,43,49	0
2	SO4	A	6	5/5	0.94	0.34	59,59,62,63	5
2	SO4	A	2	5/5	0.95	0.17	36,36,38,38	5
2	SO4	C	3	5/5	0.95	0.13	47,49,50,50	5
2	SO4	C	5	5/5	0.95	0.17	37,38,40,41	5
2	SO4	B	5	5/5	0.95	0.14	46,48,49,51	5
2	SO4	A	3	5/5	0.96	0.09	41,41,42,44	5
2	SO4	B	3	5/5	0.97	0.14	18,18,21,21	5
2	SO4	C	1	5/5	0.97	0.14	42,42,43,46	5
2	SO4	C	2	5/5	0.97	0.12	30,31,35,36	5
2	SO4	A	1	5/5	0.97	0.18	25,28,29,30	5
2	SO4	D	4	5/5	0.97	0.18	49,54,54,54	5
2	SO4	B	4	5/5	0.98	0.08	25,26,29,31	5
2	SO4	C	7	5/5	0.98	0.12	32,34,37,37	5

6.5 Other polymers [i](#)

There are no such residues in this entry.