



# Full wwPDB X-ray Structure Validation Report ⓘ

May 23, 2020 – 05:16 pm BST

PDB ID : 4RH8  
Title : Crystal structure of the outer membrane lipopolysaccharide transport protein LptE (RlpB) from Escherichia coli in the tetragonal crystal form  
Authors : Malojcic, G.; Kahne, D.  
Deposited on : 2014-10-01  
Resolution : 2.20 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.11  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.11


# 1 Overall quality at a glance ⓘ

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.20 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	4898 (2.20-2.20)
Clashscore	141614	5594 (2.20-2.20)
Ramachandran outliers	138981	5503 (2.20-2.20)
Sidechain outliers	138945	5504 (2.20-2.20)
RSRZ outliers	127900	4800 (2.20-2.20)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	168	<div> <div>2%</div> <div> <div></div> <div>66%</div> <div>7%</div> <div>•</div> <div>•</div> <div>21%</div> </div> </div>
1	B	168	<div> <div>3%</div> <div> <div></div> <div>64%</div> <div>10%</div> <div>5%</div> <div>•</div> <div>21%</div> </div> </div>
1	C	168	<div> <div>17%</div> <div> <div></div> <div>68%</div> <div>10%</div> <div>•</div> <div>20%</div> </div> </div>
1	D	168	<div> <div>11%</div> <div> <div></div> <div>68%</div> <div>9%</div> <div>•</div> <div>21%</div> </div> </div>

## 2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 4376 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called LPS-assembly lipoprotein LptE.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	133	Total	C	N	O	S	0	0	0
			1045	653	188	198	6			
1	B	133	Total	C	N	O	S	0	0	0
			1045	653	188	198	6			
1	C	135	Total	C	N	O	S	0	0	0
			1060	662	191	201	6			
1	D	133	Total	C	N	O	S	0	0	0
			1045	653	188	198	6			

There are 16 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	19	MET	CYS	CONFLICT	UNP P0ADC1
A	183	LEU	THR	ENGINEERED MUTATION	UNP P0ADC1
A	184	VAL	PRO	ENGINEERED MUTATION	UNP P0ADC1
A	185	PRO	ALA	ENGINEERED MUTATION	UNP P0ADC1
B	19	MET	CYS	CONFLICT	UNP P0ADC1
B	183	LEU	THR	ENGINEERED MUTATION	UNP P0ADC1
B	184	VAL	PRO	ENGINEERED MUTATION	UNP P0ADC1
B	185	PRO	ALA	ENGINEERED MUTATION	UNP P0ADC1
C	19	MET	CYS	CONFLICT	UNP P0ADC1
C	183	LEU	THR	ENGINEERED MUTATION	UNP P0ADC1
C	184	VAL	PRO	ENGINEERED MUTATION	UNP P0ADC1
C	185	PRO	ALA	ENGINEERED MUTATION	UNP P0ADC1
D	19	MET	CYS	CONFLICT	UNP P0ADC1
D	183	LEU	THR	ENGINEERED MUTATION	UNP P0ADC1
D	184	VAL	PRO	ENGINEERED MUTATION	UNP P0ADC1
D	185	PRO	ALA	ENGINEERED MUTATION	UNP P0ADC1

- Molecule 2 is FORMIC ACID (three-letter code: FMT) (formula: CH<sub>2</sub>O<sub>2</sub>).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	C	O	0	0
			3	1	2		
2	B	1	Total	C	O	0	0
			3	1	2		
2	C	1	Total	C	O	0	0
			3	1	2		
2	D	1	Total	C	O	0	0
			3	1	2		


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	57	Total	O	0	0
			57	57		
3	B	34	Total	O	0	0
			34	34		
3	C	32	Total	O	0	0
			32	32		
3	D	46	Total	O	0	0
			46	46		


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


#### • Molecule 1: LPS-assembly lipoprotein LptE




#### • Molecule 1: LPS-assembly lipoprotein LptE




#### • Molecule 1: LPS-assembly lipoprotein LptE



#### • Molecule 1: LPS-assembly lipoprotein LptE





## 4 Data and refinement statistics

Property	Value	Source
Space group	P 41 21 2	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	102.46 Å   102.46 Å   166.84 Å 90.00°   90.00°   90.00°	Depositor
Resolution (Å)	87.31 – 2.20 66.45 – 2.20	Depositor EDS
% Data completeness (in resolution range)	100.0 (87.31-2.20) 100.0 (66.45-2.20)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.74 (at 2.20 Å)	Xtriage
Refinement program	REFMAC 5.8.0071	Depositor
R, $R_{free}$	0.225   ,   0.262 0.229   ,   0.266	Depositor DCC
$R_{free}$ test set	2260 reflections (4.93%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	46.6	Xtriage
Anisotropy	0.187	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.34 , 43.7	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.95	EDS
Total number of atoms	4376	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	58.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.70% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality

### 5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: FMT

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.98	1/1057 (0.1%)	1.12	10/1423 (0.7%)
1	B	0.86	0/1057	1.05	7/1423 (0.5%)
1	C	0.83	0/1073	0.99	1/1446 (0.1%)
1	D	0.82	0/1057	0.95	1/1423 (0.1%)
All	All	0.87	1/4244 (0.0%)	1.03	19/5715 (0.3%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	3
1	B	0	2
1	C	0	4
1	D	0	3
All	All	0	12

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	107	THR	CB-CG2	-5.13	1.35	1.52

All (19) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	36	MET	CG-SD-CE	-9.57	84.89	100.20
1	B	69	ARG	CG-CD-NE	7.57	127.69	111.80
1	C	124	ARG	NE-CZ-NH1	7.31	123.96	120.30
1	D	124	ARG	NE-CZ-NH1	7.17	123.89	120.30

*Continued on next page...*



*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	107	THR	N-CA-CB	-7.11	96.80	110.30
1	A	150	ARG	CG-CD-NE	-7.10	96.90	111.80
1	B	150	ARG	CG-CD-NE	-6.68	97.76	111.80
1	B	36	MET	CA-CB-CG	6.57	124.46	113.30
1	A	124	ARG	NE-CZ-NH1	6.56	123.58	120.30
1	B	69	ARG	NE-CZ-NH1	6.33	123.46	120.30
1	B	124	ARG	NE-CZ-NH1	6.01	123.31	120.30
1	A	36	MET	CA-CB-CG	5.86	123.27	113.30
1	A	127	PHE	N-CA-C	5.57	126.02	111.00
1	A	126	PHE	CA-C-N	5.35	128.97	117.20
1	B	91	ARG	N-CA-C	5.33	125.39	111.00
1	B	52	ARG	NE-CZ-NH2	-5.21	117.69	120.30
1	A	91	ARG	N-CA-C	5.19	125.02	111.00
1	A	36	MET	N-CA-CB	-5.09	101.44	110.60
1	A	126	PHE	C-N-CA	5.05	134.31	121.70

There are no chirality outliers.

All (12) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	126	PHE	Peptide
1	A	91	ARG	Peptide
1	A	92	ASN	Peptide
1	B	126	PHE	Peptide
1	B	92	ASN	Peptide
1	C	66	GLU	Peptide
1	C	68	THR	Peptide
1	C	91	ARG	Peptide
1	C	92	ASN	Peptide
1	D	127	PHE	Peptide
1	D	91	ARG	Peptide
1	D	92	ASN	Peptide

## 5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1045	0	1079	15	0
1	B	1045	0	1079	18	0
1	C	1060	0	1092	16	0
1	D	1045	0	1079	8	0
2	A	3	0	2	0	0
2	B	3	0	2	0	0
2	C	3	0	2	1	0
2	D	3	0	1	0	0
3	A	57	0	0	2	0
3	B	34	0	0	2	0
3	C	32	0	0	3	0
3	D	46	0	0	3	0
All	All	4376	0	4336	53	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

All (53) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:123:PHE:CE1	1:C:121:LYS:HD3	2.03	0.93
1:A:91:ARG:HH21	1:A:95:THR:HG21	1.52	0.74
1:B:91:ARG:HH21	1:B:95:THR:HG21	1.51	0.74
1:D:91:ARG:HH21	1:D:95:THR:HG21	1.53	0.73
1:C:91:ARG:HH21	1:C:95:THR:HG21	1.54	0.72
1:A:76:ARG:HB3	1:A:107:THR:HG22	1.73	0.71
1:B:92:ASN:N	1:B:92:ASN:HD22	1.89	0.69
1:A:92:ASN:N	1:A:92:ASN:HD22	1.88	0.69
1:D:144:VAL:HA	1:D:147:MET:HE2	1.77	0.66
1:A:123:PHE:HB2	3:D:341:HOH:O	1.95	0.66
1:D:121:LYS:HD2	3:D:341:HOH:O	1.96	0.65
1:C:95:THR:CG2	3:C:306:HOH:O	2.46	0.63
1:C:98:TYR:CZ	1:C:129:ASN:HB3	2.36	0.61
1:A:28:GLN:HG3	3:A:356:HOH:O	2.01	0.60
1:B:123:PHE:CE1	1:C:121:LYS:CD	2.80	0.60
1:B:95:THR:CG2	3:B:328:HOH:O	2.51	0.58
1:B:150:ARG:NH1	1:B:153:GLU:OE2	2.40	0.55
1:A:150:ARG:NH1	1:A:153:GLU:OE2	2.40	0.55
1:B:92:ASN:N	1:B:92:ASN:ND2	2.56	0.54
1:A:92:ASN:ND2	1:A:92:ASN:N	2.55	0.54
1:C:95:THR:HG23	3:C:306:HOH:O	2.09	0.53
1:B:123:PHE:CZ	1:C:121:LYS:HD3	2.44	0.51

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:43:PRO:O	1:C:52:ARG:NH2	2.36	0.49
1:B:43:PRO:O	1:B:52:ARG:NH2	2.35	0.49
1:B:150:ARG:HH11	1:B:153:GLU:CD	2.16	0.49
1:A:150:ARG:HH11	1:A:153:GLU:CD	2.15	0.48
1:B:36:MET:HG2	1:B:60:VAL:HG11	1.95	0.47
1:B:91:ARG:C	1:B:92:ASN:HD22	2.18	0.47
1:D:142:MET:CE	1:D:145:LYS:HE3	2.45	0.46
1:B:87:ALA:HB1	1:D:79:LYS:NZ	2.30	0.46
1:C:98:TYR:CE1	1:C:129:ASN:HB3	2.51	0.46
1:A:128:ASP:O	1:A:128:ASP:CG	2.54	0.46
1:A:91:ARG:C	1:A:92:ASN:HD22	2.20	0.44
1:C:36:MET:HG2	1:C:60:VAL:HG11	1.98	0.44
1:D:70:LYS:HD3	1:D:70:LYS:HA	1.86	0.44
1:C:139:GLU:CG	1:C:140:GLN:N	2.80	0.44
1:A:150:ARG:NH2	3:A:322:HOH:O	2.51	0.44
1:C:98:TYR:CE2	1:C:129:ASN:HB3	2.53	0.43
1:B:124:ARG:HH21	1:B:146:GLU:CD	2.21	0.43
1:A:65:LYS:O	1:A:65:LYS:HG3	2.17	0.43
1:B:128:ASP:O	1:B:128:ASP:CG	2.56	0.43
1:C:95:THR:HG22	3:C:306:HOH:O	2.14	0.43
1:A:124:ARG:HH21	1:A:146:GLU:CD	2.21	0.43
1:C:129:ASN:HA	1:C:130:PRO:HD3	1.80	0.42
1:A:98:TYR:CD1	1:A:98:TYR:N	2.88	0.42
1:B:95:THR:HG22	3:B:328:HOH:O	2.16	0.41
1:C:77:LEU:O	2:C:201:FMT:O1	2.38	0.41
1:C:139:GLU:HG2	1:C:140:GLN:N	2.35	0.41
1:A:139:GLU:CG	1:A:140:GLN:N	2.84	0.41
1:D:103:THR:OG1	3:D:306:HOH:O	2.22	0.41
1:B:84:LYS:HE2	1:B:140:GLN:NE2	2.36	0.40
1:D:144:VAL:HA	1:D:147:MET:CE	2.48	0.40
1:B:91:ARG:NH2	1:B:95:THR:HG21	2.29	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles ⓘ

### 5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	129/168 (77%)	127 (98%)	1 (1%)	1 (1%)	19	19
1	B	129/168 (77%)	128 (99%)	1 (1%)	0	100	100
1	C	131/168 (78%)	128 (98%)	2 (2%)	1 (1%)	19	19
1	D	129/168 (77%)	125 (97%)	4 (3%)	0	100	100
All	All	518/672 (77%)	508 (98%)	8 (2%)	2 (0%)	34	37

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	127	PHE
1	C	92	ASN

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	117/148 (79%)	107 (92%)	10 (8%)	10	10
1	B	117/148 (79%)	106 (91%)	11 (9%)	8	8
1	C	119/148 (80%)	114 (96%)	5 (4%)	30	38
1	D	117/148 (79%)	110 (94%)	7 (6%)	19	22
All	All	470/592 (79%)	437 (93%)	33 (7%)	15	16

All (33) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	28	GLN
1	A	65	LYS
1	A	69	ARG
1	A	81	SER
1	A	92	ASN
1	A	95	THR

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	A	107	THR
1	A	114	ASP
1	A	128	ASP
1	A	150	ARG
1	B	28	GLN
1	B	57	LEU
1	B	69	ARG
1	B	70	LYS
1	B	81	SER
1	B	92	ASN
1	B	95	THR
1	B	114	ASP
1	B	127	PHE
1	B	128	ASP
1	B	145	LYS
1	C	28	GLN
1	C	70	LYS
1	C	81	SER
1	C	95	THR
1	C	129	ASN
1	D	28	GLN
1	D	52	ARG
1	D	68	THR
1	D	81	SER
1	D	95	THR
1	D	114	ASP
1	D	145	LYS

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (9) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	92	ASN
1	A	140	GLN
1	B	92	ASN
1	B	94	GLN
1	B	140	GLN
1	C	44	ASN
1	C	99	GLN
1	D	92	ASN
1	D	140	GLN

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

## 5.6 Ligand geometry ⓘ

4 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
2	FMT	B	201	-	0,2,2	0.00	-	0,1,1	0.00	-
2	FMT	D	201	-	0,2,2	0.00	-	0,1,1	0.00	-
2	FMT	A	201	-	0,2,2	0.00	-	0,1,1	0.00	-
2	FMT	C	201	-	0,2,2	0.00	-	0,1,1	0.00	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	C	201	FMT	1	0

## 5.7 Other polymers

There are no such residues in this entry.

## 5.8 Polymer linkage issues

There are no chain breaks in this entry.

## 6 Fit of model and data ⓘ

### 6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	133/168 (79%)	0.12	3 (2%) 60 58	27, 42, 79, 135	0
1	B	133/168 (79%)	0.07	5 (3%) 40 38	30, 49, 94, 134	0
1	C	135/168 (80%)	0.87	29 (21%) 0 0	31, 58, 120, 148	0
1	D	133/168 (79%)	0.39	19 (14%) 2 2	35, 56, 116, 133	0
All	All	534/672 (79%)	0.36	56 (10%) 6 5	27, 51, 111, 148	0

All (56) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	C	170	ASP	9.2
1	A	127	PHE	7.7
1	C	67	THR	7.0
1	C	168	ARG	5.7
1	C	69	ARG	5.5
1	C	167	ILE	5.0
1	C	169	SER	4.9
1	C	63	LEU	4.7
1	C	164	ALA	4.5
1	D	110	ILE	4.5
1	D	168	ARG	4.4
1	D	169	SER	4.4
1	C	110	ILE	4.3
1	C	165	ALA	4.2
1	D	170	ASP	4.1
1	D	71	ASP	4.1
1	C	111	PRO	4.0
1	C	71	ASP	3.8
1	C	112	GLY	3.7
1	D	65	LYS	3.7
1	C	166	ASP	3.5

*Continued on next page...*



*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	C	72	VAL	3.2
1	D	128	ASP	3.2
1	C	66	GLU	3.1
1	C	68	THR	3.1
1	D	68	THR	3.1
1	C	109	LEU	3.0
1	C	116	TYR	2.8
1	C	113	ARG	2.8
1	B	127	PHE	2.7
1	C	163	ARG	2.7
1	C	162	ILE	2.7
1	C	65	LYS	2.6
1	C	28	GLN	2.6
1	B	139	GLU	2.5
1	B	123	PHE	2.5
1	D	167	ILE	2.5
1	C	32	THR	2.5
1	C	114	ASP	2.4
1	B	126	PHE	2.4
1	B	141	ASP	2.4
1	C	161	SER	2.4
1	C	64	ASP	2.4
1	D	69	ARG	2.3
1	D	112	GLY	2.2
1	D	111	PRO	2.2
1	D	32	THR	2.2
1	A	126	PHE	2.1
1	D	63	LEU	2.1
1	D	66	GLU	2.1
1	D	33	MET	2.1
1	C	70	LYS	2.1
1	D	116	TYR	2.1
1	D	114	ASP	2.0
1	A	139	GLU	2.0
1	D	113	ARG	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

### 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
2	FMT	B	201	3/3	0.86	0.16	42,42,55,61	0
2	FMT	A	201	3/3	0.87	0.31	40,40,44,60	0
2	FMT	C	201	3/3	0.87	0.09	48,48,63,76	0
2	FMT	D	201	3/3	0.92	0.12	59,59,67,68	0

### 6.5 Other polymers [i](#)

There are no such residues in this entry.