



wwPDB X-ray Structure Validation Summary Report ⓘ

May 22, 2020 – 04:18 am BST

PDB ID : 4RHX
Title : Structures of Mycobacterium tuberculosis 6-oxopurine phosphoribosyltransferase which is a potential target for drug development against this disease
Authors : Eng, W.S.; Hockova, D.; Spacek, P.; West, N.P.; Woods, K.; Naesens, L.M.J.; Keough, D.T.; Guddat, L.W.
Deposited on : 2014-10-03
Resolution : 2.03 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.11
buster-report : 1.1.7 (2018)
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.11


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.03 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	1692 (2.04-2.04)
Clashscore	141614	1773 (2.04-2.04)
Ramachandran outliers	138981	1752 (2.04-2.04)
Sidechain outliers	138945	1752 (2.04-2.04)
RSRZ outliers	127900	1672 (2.04-2.04)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	201	<div> <div>2%</div> <div> <div></div> <div>83%</div> <div>6%</div> <div>10%</div> </div> </div>
1	B	201	<div> <div>3%</div> <div> <div></div> <div>78%</div> <div>8%</div> <div>13%</div> </div> </div>
1	C	201	<div> <div>3%</div> <div> <div></div> <div>78%</div> <div>8%</div> <div>12%</div> </div> </div>
1	D	201	<div> <div>2%</div> <div> <div></div> <div>78%</div> <div>9%</div> <div>13%</div> </div> </div>

2 Entry composition [i](#)


There are 4 unique types of molecules in this entry. The entry contains 6413 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Hypoxanthine-guanine phosphoribosyltransferase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	181	Total	C	N	O	S	0	1	0
			1431	910	244	275	2			
1	B	175	Total	C	N	O	S	0	1	0
			1391	883	241	265	2			
1	C	176	Total	C	N	O	S	0	0	0
			1392	885	238	267	2			
1	D	174	Total	C	N	O	S	0	1	0
			1383	880	237	264	2			

- Molecule 2 is [2-[(2-amino-6-oxo-1,6-dihydro-9H-purin-9-yl)ethyl]{2-[(2-oxoethyl)(2-phosphonoethyl)amino]ethyl}amino)ethyl]phosphonic acid (three-letter code: 3QF) (formula: C₁₅H₂₇N₇O₈P₂).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total	C	N	O	P	0	1
			96	45	21	24	6		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	B	1	Total	C	N	O	P	0	1
			64	30	14	16	4		
2	C	1	Total	C	N	O	P	0	1
			64	30	14	16	4		
2	D	1	Total	C	N	O	P	0	0
			32	15	7	8	2		

- Molecule 3 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	B	2	Total	Mg	0	0
			2	2		
3	A	2	Total	Mg	0	0
			2	2		
3	D	2	Total	Mg	0	0
			2	2		
3	C	2	Total	Mg	0	0
			2	2		


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	186	Total	O	0	0
			186	186		
4	B	119	Total	O	0	0
			119	119		
4	C	109	Total	O	0	0
			109	109		
4	D	138	Total	O	0	0
			138	138		

- Molecule 1: Hypoxanthine-guanine phosphoribosyltransferase




Category	Value	Color
HIS	160	Yellow
VAL	170	Yellow
THR	175	Yellow
GLN	179	Yellow
SER	183	Yellow
SER	184	Yellow
SER	185	Yellow
SER	186	Yellow
ALA	189	Yellow
ILE	192	Yellow
THR	193	Yellow
PRO	194	Yellow
GLY	195	Yellow
GLN	196	Yellow
THR	197	Yellow
A16	198	Red
R49	199	Green
GLU	200	Grey
LEU	201	Grey
S52	202	Yellow
A53	203	Red
L60	204	Yellow
V69	205	Yellow
L70	206	Yellow
L75	207	Yellow
L79	208	Yellow
P82	209	Yellow
T83	210	Yellow
Q84	211	Yellow
P85	212	Yellow
E86	213	Yellow
S92	214	Green
T92	215	Yellow
GLY	216	Grey
SER	217	Grey
SER	218	Grey
THR	219	Grey
SER	220	Grey
SER	221	Grey
SER	222	Grey
GLY	223	Grey
V103	224	Green
R104	225	Yellow
I105	226	Yellow
L406	227	Red
K107	228	Yellow
D108	229	Yellow
L109	230	Red
D110	231	Red
R111	232	Red
S132	233	Yellow



HIS	VAL	THR	GLN	SER	SER	SER	ALA	ILE	THR	PRO	GLY	GLN	THR	A16	L65	L70	Q84	F87	S92	Tyr	GLY	SER	SER	THR	SER	SER	SER	GLY	VAL	V103	K107	D108	L109	D110	R111	S132	R136	L152	R153	P154	P155	A156	A157	V158	H159	ALA	N161	V162	E163
-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	-----	------	------	------



Label	Value	Color
HIS	10.0	Blue
VAL	10.0	Blue
THR	10.0	Blue
GLN	10.0	Blue
SER	10.0	Blue
SER	10.0	Blue
SER	10.0	Blue
ALA	10.0	Blue
ILE	10.0	Blue
THR	10.0	Blue
PRO	10.0	Blue
GLY	10.0	Blue
GLN	10.0	Blue
THR	10.0	Blue
ALA	10.0	Blue
GLU	10.0	Blue
L18	10.0	Red
L27	10.0	Red
L60	10.0	Red
T73	10.0	Red
Q84	10.0	Red
F85	10.0	Red
E86	10.0	Red
S92	10.0	Red
TYR	10.0	Blue
GLY	10.0	Blue
SER	10.0	Blue
SER	10.0	Blue
THR	10.0	Blue
THR	10.0	Blue
SER	10.0	Blue
SER	10.0	Blue
SER	10.0	Blue
GLY	10.0	Blue
VAL	10.0	Blue
V103	10.0	Red
R104	10.0	Red
I105	10.0	Red
L106	10.0	Red
K107	10.0	Red
D108	10.0	Red
L109	10.0	Red
D110	10.0	Red
R111	10.0	Red
D123	10.0	Red
L129	10.0	Red
R136	10.0	Red
R147	10.0	Red
Y166	10.0	Red



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	53.44Å 85.72Å 88.14Å 90.00° 98.77° 90.00°	Depositor
Resolution (Å)	44.97 – 2.03 44.97 – 2.03	Depositor EDS
% Data completeness (in resolution range)	99.6 (44.97-2.03) 93.9 (44.97-2.03)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.04 (at 2.03Å)	Xtriage
Refinement program	PHENIX (phenix.refine: 1.9_1692)	Depositor
R, R_{free}	0.187 , 0.226 0.189 , 0.226	Depositor DCC
R_{free} test set	2000 reflections (3.96%)	wwPDB-VP
Wilson B-factor (Å ²)	21.7	Xtriage
Anisotropy	0.418	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.33 , 54.3	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	6413	wwPDB-VP
Average B, all atoms (Å ²)	35.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 15.32% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: MG, 3QF

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.22	0/1454	0.41	0/1976
1	B	0.22	0/1412	0.42	0/1918
1	C	0.22	0/1413	0.40	0/1920
1	D	0.21	0/1405	0.40	0/1911
All	All	0.22	0/5684	0.41	0/7725

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1431	0	1440	11	0
1	B	1391	0	1401	11	0
1	C	1392	0	1401	15	0
1	D	1383	0	1395	12	0
2	A	96	0	72	6	0
2	B	64	0	48	4	0
2	C	64	0	48	5	0
2	D	32	0	24	2	0
3	A	2	0	0	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	B	2	0	0	0	0
3	C	2	0	0	0	0
3	D	2	0	0	0	0
4	A	186	0	0	3	0
4	B	119	0	0	2	0
4	C	109	0	0	5	0
4	D	138	0	0	2	0
All	All	6413	0	5829	59	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

The worst 5 of 59 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:A:301[B]:3QF:H4	2:A:301[B]:3QF:H10	1.64	0.78
1:A:107[B]:LYS:HG3	1:B:107:LYS:HD3	1.74	0.70
1:D:84:GLN:OE1	1:D:111:ARG:NH1	2.30	0.65
2:B:301[B]:3QF:H8	4:B:518:HOH:O	1.98	0.62
1:C:107:LYS:HD2	1:D:107:LYS:HE2	1.81	0.60

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	178/201 (89%)	173 (97%)	5 (3%)	0	100	100
1	B	170/201 (85%)	167 (98%)	3 (2%)	0	100	100
1	C	170/201 (85%)	166 (98%)	4 (2%)	0	100	100
1	D	171/201 (85%)	168 (98%)	3 (2%)	0	100	100

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
All	All	689/804 (86%)	674 (98%)	15 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	158/174 (91%)	155 (98%)	3 (2%)	57	53
1	B	153/174 (88%)	151 (99%)	2 (1%)	69	67
1	C	154/174 (88%)	152 (99%)	2 (1%)	69	67
1	D	153/174 (88%)	152 (99%)	1 (1%)	84	84
All	All	618/696 (89%)	610 (99%)	8 (1%)	73	67

5 of 8 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	106	LEU
1	D	129	LEU
1	C	65	LEU
1	A	199	ARG
1	B	111	ARG

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
1	B	46	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

5.6 Ligand geometry ⓘ

Of 16 ligands modelled in this entry, 8 are monoatomic - leaving 8 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	3QF	D	301	3	30,33,33	2.85	9 (30%)	35,47,47	1.79	6 (17%)
2	3QF	C	301[B]	3	30,33,33	2.83	10 (33%)	35,47,47	1.80	5 (14%)
2	3QF	A	301[A]	3	30,33,33	2.83	9 (30%)	35,47,47	1.83	6 (17%)
2	3QF	C	301[A]	3	30,33,33	2.82	10 (33%)	35,47,47	1.80	6 (17%)
2	3QF	A	301[B]	3	30,33,33	2.83	9 (30%)	35,47,47	1.82	5 (14%)
2	3QF	A	301[C]	3	30,33,33	2.83	9 (30%)	35,47,47	1.81	5 (14%)
2	3QF	B	301[B]	3	30,33,33	2.81	9 (30%)	35,47,47	1.84	6 (17%)
2	3QF	B	301[A]	3	30,33,33	2.80	9 (30%)	35,47,47	1.82	6 (17%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	3QF	D	301	3	-	10/24/25/25	0/2/2/2
2	3QF	C	301[B]	3	-	5/24/25/25	0/2/2/2
2	3QF	A	301[A]	3	-	5/24/25/25	0/2/2/2
2	3QF	C	301[A]	3	-	3/24/25/25	0/2/2/2

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	3QF	A	301[B]	3	-	3/24/25/25	0/2/2/2
2	3QF	A	301[C]	3	-	5/24/25/25	0/2/2/2
2	3QF	B	301[B]	3	-	4/24/25/25	0/2/2/2
2	3QF	B	301[A]	3	-	4/24/25/25	0/2/2/2

The worst 5 of 74 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	D	301	3QF	PBF-CAT	8.36	1.87	1.78
2	A	301[A]	3QF	PBF-CAT	8.36	1.87	1.78
2	C	301[B]	3QF	PBF-CAT	8.28	1.87	1.78
2	C	301[A]	3QF	PBF-CAT	8.26	1.87	1.78
2	A	301[B]	3QF	PBF-CAT	8.25	1.87	1.78

The worst 5 of 45 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	301[A]	3QF	C2-N3-C4	5.62	121.77	115.36
2	A	301[B]	3QF	C2-N3-C4	5.60	121.75	115.36
2	A	301[C]	3QF	C2-N3-C4	5.60	121.75	115.36
2	B	301[A]	3QF	C2-N3-C4	5.52	121.66	115.36
2	C	301[B]	3QF	C2-N3-C4	5.49	121.63	115.36

There are no chirality outliers.

5 of 39 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	D	301	3QF	CAS-CAP-NBB-CAL
2	D	301	3QF	CAK-CAL-NBB-CAM
2	D	301	3QF	CAT-CAQ-NBC-CAO
2	D	301	3QF	CAQ-CAT-PBF-OAH
2	D	301	3QF	CAQ-CAT-PBF-OAI

There are no ring outliers.

8 monomers are involved in 17 short contacts:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	D	301	3QF	2	0
2	C	301[B]	3QF	4	0
2	A	301[A]	3QF	1	0


Continued on next page...


Continued from previous page...


Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	C	301[A]	3QF	1	0
2	A	301[B]	3QF	3	0
2	A	301[C]	3QF	2	0
2	B	301[B]	3QF	3	0
2	B	301[A]	3QF	1	0


The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.




















5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	181/201 (90%)	-0.14	4 (2%) 62 66	15, 25, 53, 108	0
1	B	175/201 (87%)	-0.06	7 (4%) 38 41	15, 29, 75, 121	0
1	C	176/201 (87%)	0.04	6 (3%) 45 49	21, 38, 74, 120	0
1	D	174/201 (86%)	-0.12	5 (2%) 51 56	17, 30, 68, 111	0
All	All	706/804 (87%)	-0.07	22 (3%) 49 53	15, 30, 72, 121	0

The worst 5 of 22 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	D	201	TYR	6.6
1	A	201	TYR	6.1
1	A	99	SER	4.3
1	B	53	ALA	4.1
1	B	110	ASP	4.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)



In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
3	MG	D	303	1/1	0.94	0.06	26,26,26,26	0
3	MG	A	302	1/1	0.95	0.05	27,27,27,27	0
2	3QF	C	301[A]	32/32	0.96	0.12	18,23,29,30	32
2	3QF	A	301[B]	32/32	0.96	0.12	3,17,21,22	32
2	3QF	A	301[C]	32/32	0.96	0.12	7,17,21,22	32
2	3QF	B	301[B]	32/32	0.96	0.11	16,19,22,23	32
2	3QF	C	301[B]	32/32	0.96	0.12	16,23,29,30	32
2	3QF	B	301[A]	32/32	0.96	0.11	15,19,22,23	32
2	3QF	A	301[A]	32/32	0.96	0.12	8,17,21,22	32
3	MG	C	303	1/1	0.97	0.10	27,27,27,27	0
3	MG	C	302	1/1	0.97	0.04	23,23,23,23	0
3	MG	D	302	1/1	0.97	0.03	19,19,19,19	0
2	3QF	D	301	32/32	0.97	0.09	20,26,34,35	0
3	MG	B	303	1/1	0.97	0.06	21,21,21,21	0
3	MG	A	303	1/1	0.98	0.05	20,20,20,20	0
3	MG	B	302	1/1	0.99	0.05	18,18,18,18	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.



Electron density around 3QF C 301 (A):

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around 3QF A 301 (B):

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around 3QF A 301 (C):

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around 3QF B 301 (B):

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)





Electron density around 3QF C 301 (B):

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around 3QF B 301 (A):

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around 3QF A 301 (A):

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



Electron density around 3QF D 301:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers ⓘ

There are no such residues in this entry.