



Full wwPDB X-ray Structure Validation Report ⓘ

Aug 20, 2020 – 10:01 PM BST

PDB ID : 6RHV
Title : Crystal structure of mouse CD11b I-domain (CD11b-I) in complex with Staphylococcus aureus octameric bi-component leukocidin LukGH (LukH K319A mutant)
Authors : Trstenjak, N.; Milic, D.; Djinovic-Carugo, K.; Badarau, A.
Deposited on : 2019-04-23
Resolution : 2.29 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.13.1
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.13.1


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.29 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	5042 (2.30-2.30)
Clashscore	141614	5643 (2.30-2.30)
Ramachandran outliers	138981	5575 (2.30-2.30)
Sidechain outliers	138945	5575 (2.30-2.30)
RSRZ outliers	127900	4938 (2.30-2.30)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	G	309	<div> <div>6%</div> <div> <div></div> <div>80%</div> <div>11%</div> <div>8%</div> </div> </div>
2	H	324	<div> <div>10%</div> <div> <div></div> <div>81%</div> <div>8%</div> <div>11%</div> </div> </div>
3	C	195	<div> <div>10%</div> <div> <div></div> <div>80%</div> <div>7%</div> <div>13%</div> </div> </div>

2 Entry composition

There are 6 unique types of molecules in this entry. The entry contains 12330 atoms, of which 5905 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Beta-channel forming cytolysin.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	G	283	Total	C	H	N	O	S	0	2	0
			4538	1459	2219	402	453	5			

- Molecule 2 is a protein called Beta-channel forming cytolysin.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	H	288	Total	C	H	N	O	0	1	0
			4681	1496	2309	410	466			

There is a discrepancy between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
H	319	ALA	LYS	engineered mutation	UNP A0A0D6HC73

- Molecule 3 is a protein called Integrin alpha-M.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
3	C	170	Total	C	H	N	O	S	0	1	0
			2694	856	1341	240	254	3			

- Molecule 4 is DIMETHYL SULFOXIDE (three-letter code: DMS) (formula: C₂H₆OS).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
4	G	1	Total	C	H	O	S	0	0
			10	2	6	1	1		
4	G	1	Total	C	H	O	S	0	0
			10	2	6	1	1		
4	G	1	Total	C	H	O	S	0	0
			10	2	6	1	1		
4	H	1	Total	C	H	O	S	0	0
			10	2	6	1	1		
4	H	1	Total	C	H	O	S	0	0
			10	2	6	1	1		
4	C	1	Total	C	H	O	S	0	0
			10	2	6	1	1		

- Molecule 5 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	H	1	Total	Mg	0	0
			1	1		

- Molecule 6 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	G	145	Total	O	0	0
			145	145		
6	H	169	Total	O	0	0
			169	169		

Continued on next page...


Continued from previous page...

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	C	42	Total	O	0	0
			42	42		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Beta-channel forming cytolsin



- Molecule 2: Beta-channel forming cytolsin



- Molecule 3: Integrin alpha-M



4 Data and refinement statistics

Property	Value	Source
Space group	P 4 21 2	Depositor
Cell constants a, b, c, α , β , γ	130.49Å 130.49Å 109.05Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	46.94 – 2.29 46.94 – 2.29	Depositor EDS
% Data completeness (in resolution range)	99.6 (46.94-2.29) 99.6 (46.94-2.29)	Depositor EDS
R_{merge}	0.19	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.07 (at 2.29Å)	Xtriage
Refinement program	PHENIX 1.14rc1_3177	Depositor
R, R_{free}	0.188 , 0.227 0.187 , 0.227	Depositor DCC
R_{free} test set	2100 reflections (4.89%)	wwPDB-VP
Wilson B-factor (Å ²)	44.5	Xtriage
Anisotropy	0.022	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.39 , 45.9	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	12330	wwPDB-VP
Average B, all atoms (Å ²)	60.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.48% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: MG, DMS

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	G	0.26	0/2382	0.46	0/3221
2	H	0.26	0/2429	0.45	0/3281
3	C	0.26	0/1382	0.41	0/1861
All	All	0.26	0/6193	0.45	0/8363

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	G	2319	2219	2220	14	0
2	H	2372	2309	2308	12	0
3	C	1353	1341	1341	7	0
4	C	4	6	6	0	0
4	G	12	18	18	0	0
4	H	8	12	12	0	0
5	H	1	0	0	0	0
6	C	42	0	0	2	0
6	G	145	0	0	1	0
6	H	169	0	0	1	0
All	All	6425	5905	5905	30	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (30) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:G:181:ASP:OD2	2:H:167:LYS:NZ	2.28	0.66
1:G:244:LYS:NZ	1:G:291:ASP:OD2	2.31	0.64
3:C:271[A]:VAL:HG23	3:C:299:VAL:HG11	1.81	0.63
1:G:127:ILE:HG13	1:G:132:LEU:HD13	1.83	0.61
2:H:119:ARG:NH2	3:C:254:ASP:OD1	2.35	0.56
1:G:126:SER:C	1:G:127:ILE:HD12	2.27	0.54
2:H:148:GLY:O	2:H:160:THR:N	2.40	0.54
1:G:38:LEU:HD11	1:G:249:VAL:HG21	1.90	0.54
2:H:126:LEU:HD11	2:H:252:SER:HB3	1.89	0.54
3:C:218:VAL:HG11	3:C:237:LEU:HD13	1.91	0.52
1:G:65:LEU:HD22	1:G:224:SER:HA	1.90	0.52
1:G:125:PHE:CE1	1:G:127:ILE:HD11	2.44	0.52
2:H:175:ASN:HB3	2:H:198:LEU:HD23	1.92	0.51
1:G:291:ASP:O	1:G:295:HIS:N	2.45	0.50
1:G:170:VAL:HG21	1:G:227:PHE:CZ	2.48	0.47
3:C:183:HIS:CE1	3:C:202:ILE:HG23	2.50	0.47
2:H:154:THR:OG1	2:H:155:LYS:N	2.47	0.47
2:H:198:LEU:HD12	2:H:212:LEU:HD21	1.98	0.46
1:G:114:LYS:NZ	6:G:505:HOH:O	2.44	0.46
2:H:287:GLU:HB3	2:H:290:LYS:HB2	1.97	0.46
1:G:135:ASN:HA	2:H:151:PHE:O	2.16	0.45
1:G:127:ILE:HD12	1:G:127:ILE:N	2.32	0.45
2:H:91:LEU:HD11	2:H:105:LEU:HD11	1.98	0.45
2:H:80:LEU:HD11	2:H:248:LEU:HD23	2.00	0.43
3:C:174:MET:HB2	3:C:222:LEU:HD11	2.00	0.43
3:C:253:LYS:NZ	6:C:507:HOH:O	2.51	0.43
1:G:250:HIS:CE1	1:G:285:SER:HG	2.27	0.42
3:C:165:LYS:NZ	6:C:504:HOH:O	2.47	0.41
1:G:125:PHE:HA	1:G:134:GLY:HA2	2.03	0.41
2:H:240:ARG:NH1	6:H:513:HOH:O	2.51	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	G	281/309 (91%)	265 (94%)	16 (6%)	0	100	100
2	H	287/324 (89%)	276 (96%)	11 (4%)	0	100	100
3	C	169/195 (87%)	157 (93%)	12 (7%)	0	100	100
All	All	737/828 (89%)	698 (95%)	39 (5%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	G	260/281 (92%)	242 (93%)	18 (7%)	15	20
2	H	266/300 (89%)	258 (97%)	8 (3%)	41	57
3	C	149/172 (87%)	146 (98%)	3 (2%)	55	72
All	All	675/753 (90%)	646 (96%)	29 (4%)	29	40

All (29) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	G	19	LYS
1	G	20	MET
1	G	66	ARG
1	G	107	GLN
1	G	110	SER

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	G	111	ARG
1	G	124	ASP
1	G	126	SER
1	G	135	ASN
1	G	178	MET
1	G	204	ASN
1	G	210	LYS
1	G	216	LYS
1	G	264	ARG
1	G	281	GLU
1	G	282	GLU
1	G	293	LYS
1	G	304	ASN
2	H	39	ASP
2	H	45	LYS
2	H	106	LYS
2	H	121	ARG
2	H	150	LYS
2	H	151	PHE
2	H	159	ARG
2	H	207	ARG
3	C	203	LYS
3	C	244	GLU
3	C	301	ASN

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 7 ligands modelled in this entry, 1 is monoatomic - leaving 6 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
4	DMS	H	402	-	3,3,3	0.68	0	3,3,3	0.57	0
4	DMS	H	401	-	3,3,3	0.66	0	3,3,3	0.51	0
4	DMS	G	401	-	3,3,3	0.63	0	3,3,3	0.40	0
4	DMS	G	402	-	3,3,3	0.66	0	3,3,3	0.50	0
4	DMS	C	401	-	3,3,3	0.65	0	3,3,3	0.48	0
4	DMS	G	403	-	3,3,3	0.64	0	3,3,3	0.57	0

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	G	283/309 (91%)	0.43	17 (6%) 21 28	31, 47, 94, 131	0
2	H	288/324 (88%)	0.73	34 (11%) 4 6	32, 44, 95, 141	0
3	C	170/195 (87%)	0.69	19 (11%) 5 7	40, 60, 100, 116	0
All	All	741/828 (89%)	0.61	70 (9%) 8 11	31, 49, 99, 141	0

All (70) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	G	127	ILE	10.6
1	G	267	PHE	7.5
2	H	158	GLY	6.4
1	G	125	PHE	6.3
1	G	132	LEU	5.7
2	H	154	THR	5.4
2	H	41	GLY	5.2
3	C	234	ALA	5.1
2	H	152	ASP	4.9
2	H	153	SER	4.8
2	H	38	ASP	4.6
1	G	133	THR	4.6
2	H	157	ILE	4.5
1	G	134	GLY	4.4
2	H	151	PHE	4.4
2	H	39	ASP	4.4
2	H	149	GLY	4.3
2	H	40	ILE	4.3
2	H	150	LYS	4.0
1	G	126	SER	4.0
1	G	123	GLY	3.6
1	G	124	ASP	3.6
3	C	236	ILE	3.6

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
2	H	44	GLY	3.4
2	H	156	GLY	3.4
1	G	73	TYR	3.3
1	G	268	TRP	3.2
1	G	135	ASN	3.0
3	C	194	SER	2.9
3	C	169	THR	2.9
2	H	298	THR	2.8
2	H	37	PRO	2.8
2	H	159	ARG	2.8
2	H	299	TYR	2.8
2	H	148	GLY	2.7
3	C	170	LEU	2.7
3	C	299	VAL	2.7
2	H	160	THR	2.7
1	G	69	ASP	2.6
2	H	262	PHE	2.6
2	H	115	PHE	2.5
2	H	260	THR	2.5
2	H	311	VAL	2.5
3	C	133	SER	2.5
2	H	43	ASN	2.5
2	H	191	TRP	2.4
3	C	265	ILE	2.4
3	C	229	ALA	2.4
2	H	264	VAL	2.3
1	G	270	TYR	2.3
3	C	199	VAL	2.3
3	C	163	GLN	2.3
1	G	136	ILE	2.3
2	H	113	VAL	2.3
2	H	117	VAL	2.3
2	H	187	TRP	2.3
2	H	297	VAL	2.3
3	C	166	LYS	2.2
3	C	223	PHE	2.2
1	G	138	LYS	2.2
2	H	188	HIS	2.2
3	C	162	GLU	2.2
3	C	270	GLY	2.2
3	C	189	PHE	2.2
2	H	42	LYS	2.1

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
3	C	159	THR	2.1
3	C	232	ASN	2.1
2	H	116	GLN	2.1
3	C	156	PHE	2.1
1	G	119	TYR	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
4	DMS	G	403	4/4	0.81	0.34	76,92,92,92	0
4	DMS	C	401	4/4	0.90	0.34	79,98,106,106	0
4	DMS	H	402	4/4	0.92	0.29	76,92,95,95	0
4	DMS	H	401	4/4	0.93	0.22	75,90,91,91	0
4	DMS	G	402	4/4	0.98	0.23	75,90,92,92	0
4	DMS	G	401	4/4	0.99	0.20	52,63,65,65	0
5	MG	H	403	1/1	0.99	0.08	36,36,36,36	0

6.5 Other polymers [i](#)

There are no such residues in this entry.