



# wwPDB X-ray Structure Validation Summary Report ⓘ

Jan 5, 2022 – 04:11 PM EST

PDB ID : 7RIW  
Title : RNA polymerase II elongation complex scaffold 2, without polyamide  
Authors : Oh, J.; Dervan, P.B.; Wang, D.  
Deposited on : 2021-07-20  
Resolution : 3.20 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Xtriage (Phenix)	:	1.13
EDS	:	2.25
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac	:	5.8.0158
CCP4	:	7.0.044 (Gargrove)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.25


# 1 Overall quality at a glance i

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 3.20 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.












Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R <sub>free</sub>	130704	1133 (3.20-3.20)
Clashscore	141614	1253 (3.20-3.20)
Ramachandran outliers	138981	1234 (3.20-3.20)
Sidechain outliers	138945	1233 (3.20-3.20)
RSRZ outliers	127900	1095 (3.20-3.20)
RNA backbone	3102	1010 (3.50-2.90)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	R	10	<div><div></div><div>50%40%10%</div></div>
2	T	30	<div><div>7%</div><div>30%57%13%</div></div>
3	A	1733	<div><div>2%</div><div>57%22%20%</div></div>
4	B	1224	<div><div>%</div><div>63%27%8%</div></div>

Continued on next page...

*Continued from previous page...*

Mol	Chain	Length	Quality of chain
5	C	318	
6	E	215	
7	F	155	
8	H	146	
9	I	122	
10	J	70	
11	K	120	
12	L	70	
13	N	20	

## 2 Entry composition

There are 15 unique types of molecules in this entry. The entry contains 29078 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a RNA chain called RNA.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	R	9	Total	C	N	O	P	0	0	0
			193	87	38	60	8			

- Molecule 2 is a DNA chain called Template strand DNA.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	T	26	Total	C	N	O	P	0	0	0
			524	251	85	162	26			

- Molecule 3 is a protein called DNA-directed RNA polymerase II subunit RPB1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	A	1384	Total	C	N	O	S	0	0	0
			10827	6830	1895	2042	60			

- Molecule 4 is a protein called DNA-directed RNA polymerase II subunit RPB2.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
4	B	1126	Total	C	N	O	S	0	0	0
			8871	5614	1554	1650	53			

- Molecule 5 is a protein called DNA-directed RNA polymerase II subunit RPB3.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
5	C	267	Total	C	N	O	S	0	0	0
			2101	1320	349	419	13			

- Molecule 6 is a protein called DNA-directed RNA polymerases I, II, and III subunit RPABC1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
6	E	212	Total	C	N	O	S	0	0	0
			1731	1100	305	315	11			

- Molecule 7 is a protein called DNA-directed RNA polymerases I, II, and III subunit RPABC2.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
7	F	86	Total	C	N	O	S	0	0	0
			684	437	115	129	3			

- Molecule 8 is a protein called DNA-directed RNA polymerases I, II, and III subunit RPABC3.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
8	H	133	Total	C	N	O	S	0	0	0
			1064	670	179	211	4			

- Molecule 9 is a protein called DNA-directed RNA polymerase II subunit RPB9.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
9	I	118	Total	C	N	O	S	0	0	0
			952	585	173	184	10			

- Molecule 10 is a protein called DNA-directed RNA polymerases I, II, and III subunit RPABC5.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
10	J	65	Total	C	N	O	S	0	0	0
			532	339	93	94	6			

- Molecule 11 is a protein called DNA-directed RNA polymerase II subunit RPB11.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
11	K	114	Total	C	N	O	S	0	0	0
			919	590	156	171	2			

- Molecule 12 is a protein called DNA-directed RNA polymerases I, II, and III subunit RPABC4.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
12	L	43	Total	C	N	O	S	0	0	0
			337	208	66	59	4			

- Molecule 13 is a DNA chain called Non-template strand DNA.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
13	N	16	Total	C	N	O	P	0	0	0
			334	157	71	90	16			


- Molecule 14 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
14	R	1	Total	Mg	0	0
			1	1		


- Molecule 15 is ZINC ION (three-letter code: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
15	A	2	Total	Zn	0	0
			2	2		
15	B	1	Total	Zn	0	0
			1	1		
15	C	1	Total	Zn	0	0
			1	1		
15	I	2	Total	Zn	0	0
			2	2		
15	J	1	Total	Zn	0	0
			1	1		
15	L	1	Total	Zn	0	0
			1	1		






### • Molecule 4: DNA-directed RNA polymerase II subunit RPB2








• Molecule 5: DNA-directed RNA polymerase II subunit RPB3


Chain C: 61% 22% 16%




• Molecule 6: DNA-directed RNA polymerases I, II, and III subunit RPABC1

Chain E: 3% 69% 28%







- Molecule 7: DNA-directed RNA polymerases I, II, and III subunit RPABC2



- Molecule 8: DNA-directed RNA polymerases I, II, and III subunit RPABC3



- Molecule 9: DNA-directed RNA polymerase II subunit RPB9




- Molecule 10: DNA-directed RNA polymerases I, II, and III subunit RPABC5



- Molecule 11: DNA-directed RNA polymerase II subunit RPB11





● Molecule 12: DNA-directed RNA polymerases I, II, and III subunit RPABC4



● Molecule 13: Non-template strand DNA



## 4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	165.01Å 222.44Å 192.30Å 90.00° 99.61° 90.00°	Depositor
Resolution (Å)	47.80 – 3.20 47.81 – 3.20	Depositor EDS
% Data completeness (in resolution range)	99.8 (47.80-3.20) 99.8 (47.81-3.20)	Depositor EDS
$R_{merge}$	0.41	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.18 (at 3.19Å)	Xtriage
Refinement program	PHENIX 1.13	Depositor
R, $R_{free}$	0.228 , 0.265 0.228 , 0.265	Depositor DCC
$R_{free}$ test set	1944 reflections (1.73%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	82.4	Xtriage
Anisotropy	0.473	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.27 , 37.4	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.45$ , $\langle L^2 \rangle = 0.28$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.93	EDS
Total number of atoms	29078	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	100.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.64% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: MG, ZN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z  > 5$	RMSZ	# $ Z  > 5$
1	R	0.31	0/216	0.92	0/336
2	T	0.63	0/583	1.07	0/896
3	A	0.27	0/11019	0.45	0/14906
4	B	0.27	0/9042	0.45	0/12203
5	C	0.28	0/2139	0.45	0/2899
6	E	0.26	0/1767	0.44	0/2378
7	F	0.25	0/696	0.44	0/943
8	H	0.26	0/1082	0.48	0/1466
9	I	0.29	0/970	0.48	0/1308
10	J	0.24	0/541	0.46	0/727
11	K	0.27	0/937	0.44	0/1265
12	L	0.28	0/339	0.47	0/450
13	N	0.62	0/377	0.83	0/580
All	All	0.29	0/29708	0.49	0/40357

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	R	193	0	99	6	0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
2	T	524	0	296	25	0
3	A	10827	0	10873	280	0
4	B	8871	0	8820	251	0
5	C	2101	0	2056	53	0
6	E	1731	0	1758	36	0
7	F	684	0	692	15	0
8	H	1064	0	1029	35	0
9	I	952	0	898	31	0
10	J	532	0	542	14	0
11	K	919	0	929	25	0
12	L	337	0	352	6	0
13	N	334	0	178	9	0
14	R	1	0	0	0	0
15	A	2	0	0	0	0
15	B	1	0	0	0	0
15	C	1	0	0	0	0
15	I	2	0	0	0	0
15	J	1	0	0	0	0
15	L	1	0	0	0	0
All	All	29078	0	28522	693	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 12.

The worst 5 of 693 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:A:1329:THR:HG22	3:A:1331:SER:H	1.09	1.10
4:B:570:VAL:HG23	4:B:573:GLN:HB2	1.43	0.98
3:A:122:MET:O	3:A:126:LEU:HD12	1.68	0.92
11:K:10:PHE:CD1	11:K:11:LEU:HD13	2.08	0.89
3:A:444:PHE:HE2	3:A:470:LEU:HD23	1.38	0.88

There are no symmetry-related clashes.

## 5.3 Torsion angles

### 5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
3	A	1370/1733 (79%)	1307 (95%)	63 (5%)	0	100	100
4	B	1106/1224 (90%)	1066 (96%)	40 (4%)	0	100	100
5	C	265/318 (83%)	256 (97%)	9 (3%)	0	100	100
6	E	210/215 (98%)	199 (95%)	11 (5%)	0	100	100
7	F	84/155 (54%)	84 (100%)	0	0	100	100
8	H	129/146 (88%)	118 (92%)	11 (8%)	0	100	100
9	I	116/122 (95%)	110 (95%)	6 (5%)	0	100	100
10	J	63/70 (90%)	61 (97%)	2 (3%)	0	100	100
11	K	112/120 (93%)	108 (96%)	4 (4%)	0	100	100
12	L	41/70 (59%)	40 (98%)	1 (2%)	0	100	100
All	All	3496/4173 (84%)	3349 (96%)	147 (4%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
3	A	1194/1520 (79%)	1157 (97%)	37 (3%)	40	72
4	B	955/1061 (90%)	919 (96%)	36 (4%)	33	67
5	C	235/274 (86%)	231 (98%)	4 (2%)	60	83
6	E	193/197 (98%)	187 (97%)	6 (3%)	40	72

*Continued on next page...*

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
7	F	73/137 (53%)	71 (97%)	2 (3%)	44	75
8	H	116/128 (91%)	107 (92%)	9 (8%)	12	43
9	I	110/116 (95%)	110 (100%)	0	100	100
10	J	60/65 (92%)	59 (98%)	1 (2%)	60	83
11	K	99/102 (97%)	97 (98%)	2 (2%)	55	80
12	L	37/57 (65%)	37 (100%)	0	100	100
All	All	3072/3657 (84%)	2975 (97%)	97 (3%)	39	71

5 of 97 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
4	B	722	ASP
4	B	1181	GLU
4	B	953	LEU
4	B	1147	LEU
5	C	193	TYR

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 6 such sidechains are listed below:

Mol	Chain	Res	Type
4	B	1117	GLN
6	E	143	ASN
8	H	33	GLN
3	A	339	ASN
3	A	71	GLN

### 5.3.3 RNA ⓘ

Mol	Chain	Analysed	Backbone Outliers	Pucker Outliers
1	R	8/10 (80%)	0	0

There are no RNA backbone outliers to report.

There are no RNA pucker outliers to report.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.



## 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 5.6 Ligand geometry [i](#)

Of 9 ligands modelled in this entry, 9 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data ⓘ

### 6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	R	9/10 (90%)	-0.45	0 100 100	81, 87, 139, 143	0
2	T	26/30 (86%)	-0.05	2 (7%) 13 7	87, 164, 246, 261	0
3	A	1384/1733 (79%)	-0.02	36 (2%) 56 40	47, 94, 162, 198	0
4	B	1126/1224 (91%)	-0.01	14 (1%) 79 67	41, 81, 131, 176	0
5	C	267/318 (83%)	-0.21	0 100 100	55, 84, 119, 158	0
6	E	212/215 (98%)	0.01	6 (2%) 53 37	76, 130, 181, 195	0
7	F	86/155 (55%)	-0.12	1 (1%) 79 67	68, 99, 142, 176	0
8	H	133/146 (91%)	0.54	13 (9%) 7 4	89, 124, 164, 195	0
9	I	118/122 (96%)	-0.27	1 (0%) 86 78	63, 98, 128, 144	0
10	J	65/70 (92%)	-0.30	0 100 100	57, 75, 110, 127	0
11	K	114/120 (95%)	-0.14	0 100 100	57, 89, 115, 134	0
12	L	43/70 (61%)	0.53	3 (6%) 16 9	61, 146, 189, 198	0
13	N	16/20 (80%)	0.19	1 (6%) 20 11	153, 205, 239, 242	0
All	All	3599/4233 (85%)	-0.03	77 (2%) 63 49	41, 92, 162, 261	0

The worst 5 of 77 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
3	A	105	CYS	4.5
4	B	869	SER	4.5
3	A	183	GLY	4.3
3	A	103	CYS	4.0
4	B	867	GLY	4.0

### 6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

### 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
15	ZN	A	1802	1/1	0.78	0.11	135,135,135,135	0
15	ZN	A	1801	1/1	0.88	0.17	205,205,205,205	0
14	MG	R	2001	1/1	0.88	0.11	104,104,104,104	0
15	ZN	L	101	1/1	0.89	0.14	215,215,215,215	0
15	ZN	B	1301	1/1	0.95	0.09	149,149,149,149	0
15	ZN	J	101	1/1	0.98	0.18	63,63,63,63	0
15	ZN	I	201	1/1	0.98	0.10	112,112,112,112	0
15	ZN	C	401	1/1	0.99	0.10	91,91,91,91	0
15	ZN	I	202	1/1	0.99	0.12	81,81,81,81	0

### 6.5 Other polymers [i](#)

There are no such residues in this entry.