



Full wwPDB X-ray Structure Validation Report ⓘ

May 13, 2020 – 01:41 pm BST

PDB ID : 3RJW
Title : Crystal structure of histone lysine methyltransferase g9a with an inhibitor
Authors : Dong, A.; Wasney, G.A.; Tempel, W.; Liu, F.; Barsyte, D.; Allali-Hassani, A.; Chen, X.; Chau, I.; Hajian, T.; Senisterra, G.; Chavda, N.; Arora, K.; Siarheyeva, A.; Kireev, D.B.; Herold, J.M.; Bochkarev, A.; Bountra, C.; Weigelt, J.; Edwards, A.M.; Frye, S.V.; Arrowsmith, C.H.; Brown, P.J.; Jin, J.; Vedadi, M.; Structural Genomics Consortium (SGC)
Deposited on : 2011-04-15
Resolution : 2.56 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix)	:	1.13
EDS	:	2.11
buster-report	:	1.1.7 (2018)
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac	:	5.8.0158
CCP4	:	7.0.044 (Gargrove)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.11


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.56 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	1279 (2.58-2.54)
Clashscore	141614	1327 (2.58-2.54)
Ramachandran outliers	138981	1312 (2.58-2.54)
Sidechain outliers	138945	1312 (2.58-2.54)
RSRZ outliers	127900	1269 (2.58-2.54)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	283	<div> <div>2%</div> <div> <div></div> <div>93%</div> <div>5%</div> </div> </div>
1	B	283	<div> <div>4%</div> <div> <div></div> <div>92%</div> <div>5%</div> </div> </div>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
5	UNX	A	4260	-	-	-	X

2 Entry composition

There are 6 unique types of molecules in this entry. The entry contains 4301 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Histone-lysine N-methyltransferase EHMT2.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	268	Total	C	N	O	S	0	0	0
			2027	1261	353	390	23			
1	B	270	Total	C	N	O	S	0	0	0
			2064	1289	356	396	23			


There are 4 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	911	GLY	-	EXPRESSION TAG	UNP Q96KQ7
A	912	SER	-	EXPRESSION TAG	UNP Q96KQ7
B	911	GLY	-	EXPRESSION TAG	UNP Q96KQ7
B	912	SER	-	EXPRESSION TAG	UNP Q96KQ7

- Molecule 2 is ZINC ION (three-letter code: ZN) (formula: Zn).


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	B	4	Total	Zn	0	0
			4	4		
2	A	4	Total	Zn	0	0
			4	4		

- Molecule 3 is S-ADENOSYL-L-HOMOCYSTEINE (three-letter code: SAH) (formula: C₁₄H₂₀N₆O₅S).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
3	A	1	Total	C	N	O	S	0	0
			26	14	6	5	1		
3	B	1	Total	C	N	O	S	0	0
			26	14	6	5	1		

- Molecule 4 is 2-cyclohexyl-6-methoxy-N-[1-(1-methylethyl)piperidin-4-yl]-7-(3-pyrrolidin-1-ylpropoxy)quinazolin-4-amine (three-letter code: CIQ) (formula: C₃₀H₄₇N₅O₂).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
4	A	1	Total	C	N	O	0	0
			37	30	5	2		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
4	B	1	Total	C	N	O	0	0
			37	30	5	2		

- Molecule 5 is UNKNOWN ATOM OR ION (three-letter code: UNX) (formula: X).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	B	13	Total	X	0	0
			13	13		
5	A	16	Total	X	0	0
			16	16		


- Molecule 6 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	24	Total	O	0	0
			24	24		
6	B	23	Total	O	0	0
			23	23		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Histone-lysine N-methyltransferase EHMT2



- Molecule 1: Histone-lysine N-methyltransferase EHMT2



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	56.58Å 78.08Å 70.25Å 90.00° 92.16° 90.00°	Depositor
Resolution (Å)	43.25 – 2.56 43.25 – 2.56	Depositor EDS
% Data completeness (in resolution range)	(Not available) (43.25-2.56) 99.8 (43.25-2.56)	Depositor EDS
R_{merge}	0.10	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.79 (at 2.54Å)	Xtriage
Refinement program	BUSTER 2.8.0	Depositor
R, R_{free}	0.195 , 0.245 0.214 , 0.271	Depositor DCC
R_{free} test set	1008 reflections (5.10%)	wwPDB-VP
Wilson B-factor (Å ²)	47.6	Xtriage
Anisotropy	0.129	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.30 , 41.2	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	0.031 for h,-k,-l	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	4301	wwPDB-VP
Average B, all atoms (Å ²)	61.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 29.53 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 1.5263e-03. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

¹ Intensities estimated from amplitudes.

² Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality ⓘ

5.1 Standard geometry ⓘ

Bond lengths and bond angles in the following residue types are not validated in this section: UNX, ZN, SAH, CIQ

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.47	0/2071	0.63	0/2814
1	B	0.49	0/2109	0.63	0/2864
All	All	0.48	0/4180	0.63	0/5678

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2027	0	1766	3	0
1	B	2064	0	1825	6	0
2	A	4	0	0	0	0
2	B	4	0	0	0	0
3	A	26	0	19	1	0
3	B	26	0	19	1	0
4	A	37	0	47	2	0
4	B	37	0	47	2	0
5	A	16	0	0	0	0
5	B	13	0	0	0	0
6	A	24	0	0	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
6	B	23	0	0	0	0
All	All	4301	0	3723	11	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 1.

All (11) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:1078:ASP:O	4:B:2000:CIQ:HBG	2.02	0.60
1:B:1013:LEU:HD22	1:B:1125:PHE:HE2	1.71	0.55
1:B:1158:PHE:HA	4:B:2000:CIQ:HAAB	1.91	0.52
1:A:1158:PHE:HA	4:A:2000:CIQ:HAAB	1.94	0.49
1:B:921:ILE:HA	1:B:938:VAL:HG12	2.01	0.43
3:A:1198:SAH:HG2	3:A:1198:SAH:O4'	2.18	0.42
3:B:1198:SAH:O4'	3:B:1198:SAH:HG2	2.19	0.42
1:A:1083:ASP:OD2	4:A:2000:CIQ:HAE	2.19	0.41
1:B:1067:TYR:HA	1:B:1108:SER:OG	2.21	0.40
1:A:1067:TYR:HA	1:A:1108:SER:OG	2.21	0.40
1:B:1132:ARG:CZ	1:B:1133:PHE:HE2	2.34	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	264/283 (93%)	250 (95%)	14 (5%)	0	100	100
1	B	266/283 (94%)	252 (95%)	14 (5%)	0	100	100
All	All	530/566 (94%)	502 (95%)	28 (5%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	202/253 (80%)	201 (100%)	1 (0%)	88	93
1	B	209/253 (83%)	208 (100%)	1 (0%)	88	93
All	All	411/506 (81%)	409 (100%)	2 (0%)	88	93

All (2) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	961	THR
1	B	961	THR

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

Of 41 ligands modelled in this entry, 29 are unknown and 8 are monoatomic - leaving 4 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and

the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
4	CIQ	B	2000	-	39,41,41	0.97	2 (5%)	54,56,56	1.98	11 (20%)
4	CIQ	A	2000	-	39,41,41	0.99	3 (7%)	54,56,56	2.05	15 (27%)
3	SAH	A	1198	-	21,28,28	1.36	3 (14%)	20,40,40	1.72	3 (15%)
3	SAH	B	1198	-	21,28,28	1.23	2 (9%)	20,40,40	1.72	3 (15%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	CIQ	B	2000	-	-	8/21/46/46	0/5/5/5
4	CIQ	A	2000	-	-	8/21/46/46	0/5/5/5
3	SAH	A	1198	-	-	2/7/31/31	0/3/3/3
3	SAH	B	1198	-	-	2/7/31/31	0/3/3/3

All (10) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	A	2000	CIQ	C6-C5	-3.61	1.40	1.44
4	B	2000	CIQ	C6-C5	-3.61	1.40	1.44
3	A	1198	SAH	O4'-C1'	3.47	1.45	1.41
3	B	1198	SAH	O4'-C1'	3.15	1.45	1.41
3	B	1198	SAH	C8-N7	-2.64	1.30	1.34
3	A	1198	SAH	C8-N7	-2.51	1.30	1.34
4	B	2000	CIQ	CAD-CBB	2.38	1.40	1.36
4	A	2000	CIQ	CAE-CBA	2.13	1.40	1.36
3	A	1198	SAH	C2-N3	2.13	1.35	1.32
4	A	2000	CIQ	CAD-CBB	2.06	1.40	1.36

All (32) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	A	2000	CIQ	C2-N3-C4	6.43	120.90	116.54

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	B	2000	CIQ	C2-N3-C4	6.08	120.67	116.54
4	B	2000	CIQ	C6-C5-C4	5.12	119.10	115.88
3	B	1198	SAH	N3-C2-N1	-5.03	120.82	128.68
4	B	2000	CIQ	OAY-CBA-CAE	-4.39	119.56	125.24
4	A	2000	CIQ	CAA-OAY-CBA	-4.27	111.09	117.53
3	A	1198	SAH	N3-C2-N1	-4.24	122.05	128.68
4	B	2000	CIQ	CAE-C5-C6	-4.20	121.10	124.88
4	A	2000	CIQ	C6-C5-C4	4.16	118.50	115.88
4	A	2000	CIQ	OAY-CBA-CAE	-4.11	119.92	125.24
4	A	2000	CIQ	N3-C2-N1	-3.97	120.44	126.09
4	B	2000	CIQ	CAA-OAY-CBA	-3.89	111.65	117.53
4	B	2000	CIQ	N3-C2-N1	-3.77	120.73	126.09
4	B	2000	CIQ	OAY-CBA-CBB	3.62	120.46	115.41
4	A	2000	CIQ	OAY-CBA-CBB	3.57	120.38	115.41
4	B	2000	CIQ	C5-C4-N3	-3.56	119.04	122.81
4	A	2000	CIQ	CAL-OAZ-CBB	-3.56	108.98	117.69
3	A	1198	SAH	O4'-C1'-C2'	-3.26	102.16	106.93
4	A	2000	CIQ	OAZ-CBB-CAD	-3.22	119.73	125.19
4	A	2000	CIQ	C5-C4-N3	-3.03	119.60	122.81
3	A	1198	SAH	C4-C5-N7	-3.02	106.25	109.40
4	A	2000	CIQ	CAE-C5-C6	-3.01	122.16	124.88
3	B	1198	SAH	O4'-C1'-C2'	-2.82	102.80	106.93
3	B	1198	SAH	C4-C5-N7	-2.79	106.49	109.40
4	A	2000	CIQ	C6-NAX-CBH	-2.73	118.97	124.16
4	B	2000	CIQ	C6-NAX-CBH	-2.59	119.24	124.16
4	A	2000	CIQ	C2-N1-C6	2.54	121.59	117.20
4	B	2000	CIQ	C2-N1-C6	2.47	121.47	117.20
4	B	2000	CIQ	OAZ-CBB-CAD	-2.32	121.27	125.19
4	A	2000	CIQ	OAZ-CBB-CBA	2.27	120.41	115.73
4	A	2000	CIQ	CAK-CAS-NBJ	-2.24	108.20	113.84
4	A	2000	CIQ	C5-C6-NAX	2.17	122.67	120.63

There are no chirality outliers.

All (20) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	A	2000	CIQ	N1-C6-NAX-CBH
4	A	2000	CIQ	CBB-CBA-OAY-CAA
4	A	2000	CIQ	CAE-CBA-OAY-CAA
4	B	2000	CIQ	CBB-CBA-OAY-CAA
4	B	2000	CIQ	CAE-CBA-OAY-CAA
3	A	1198	SAH	CA-CB-CG-SD

Continued on next page...

Continued from previous page...



Mol	Chain	Res	Type	Atoms
4	B	2000	CIQ	CAL-CAK-CAS-NBJ
4	A	2000	CIQ	C5-C6-NAX-CBH
4	A	2000	CIQ	CBA-CBB-OAZ-CAL
3	B	1198	SAH	CA-CB-CG-SD
4	B	2000	CIQ	CBA-CBB-OAZ-CAL
4	A	2000	CIQ	CAL-CAK-CAS-NBJ
4	B	2000	CIQ	CAB-CBG-NBK-CAU
4	A	2000	CIQ	CAD-CBB-OAZ-CAL
4	B	2000	CIQ	N3-C2-CBI-CAN
3	A	1198	SAH	CB-CG-SD-C5'
4	B	2000	CIQ	CAD-CBB-OAZ-CAL
3	B	1198	SAH	CB-CG-SD-C5'
4	B	2000	CIQ	CAS-CAK-CAL-OAZ
4	A	2000	CIQ	N3-C2-CBI-CAN


There are no ring outliers.

4 monomers are involved in 6 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	B	2000	CIQ	2	0
4	A	2000	CIQ	2	0
3	A	1198	SAH	1	0
3	B	1198	SAH	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues ⓘ

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	268/283 (94%)	0.15	7 (2%) 56 64	38, 62, 107, 192	0
1	B	270/283 (95%)	0.06	10 (3%) 41 50	34, 56, 91, 157	0
All	All	538/566 (95%)	0.11	17 (3%) 47 57	34, 59, 102, 192	0

All (17) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	1183	ALA	5.7
1	A	1182	ILE	4.9
1	A	1181	ALA	3.7
1	A	1158	PHE	3.7
1	A	1185	GLU	3.3
1	B	1166	PHE	3.1
1	A	1179	ALA	3.1
1	B	1163	SER	2.8
1	B	1190	ALA	2.7
1	B	1091	ASN	2.5
1	B	991	SER	2.5
1	B	921	ILE	2.5
1	A	1159	TRP	2.4
1	B	1184	LEU	2.3
1	B	1183	ALA	2.2
1	B	1159	TRP	2.1
1	B	1077	ALA	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates ⓘ

There are no carbohydrates in this entry.

6.4 Ligands ⓘ

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
5	UNX	A	4260	1/1	0.63	0.47	30,30,30,30	0
5	UNX	B	4261	1/1	0.73	0.33	30,30,30,30	0
5	UNX	A	9	1/1	0.73	0.19	30,30,30,30	0
5	UNX	B	2	1/1	0.76	0.26	30,30,30,30	0
5	UNX	A	7	1/1	0.77	0.25	30,30,30,30	0
5	UNX	B	4264	1/1	0.78	0.26	30,30,30,30	0
5	UNX	A	4258	1/1	0.82	0.17	30,30,30,30	0
5	UNX	A	4255	1/1	0.82	0.34	30,30,30,30	0
5	UNX	B	4262	1/1	0.86	0.24	30,30,30,30	0
5	UNX	A	4302	1/1	0.87	0.26	30,30,30,30	0
5	UNX	A	4259	1/1	0.89	0.20	30,30,30,30	0
5	UNX	B	4304	1/1	0.89	0.25	30,30,30,30	0
5	UNX	B	6	1/1	0.89	0.30	30,30,30,30	0
5	UNX	A	4248	1/1	0.89	0.24	30,30,30,30	0
5	UNX	A	4256	1/1	0.90	0.25	30,30,30,30	0
5	UNX	B	1	1/1	0.90	0.09	30,30,30,30	0
5	UNX	B	4306	1/1	0.90	0.20	30,30,30,30	0
5	UNX	A	4	1/1	0.91	0.21	30,30,30,30	0
5	UNX	A	4257	1/1	0.91	0.17	30,30,30,30	0
5	UNX	B	3	1/1	0.92	0.54	30,30,30,30	0
5	UNX	B	4266	1/1	0.92	0.11	30,30,30,30	0
5	UNX	B	4305	1/1	0.92	0.13	30,30,30,30	0
3	SAH	A	1198	26/26	0.93	0.18	57,66,74,74	0
5	UNX	A	10	1/1	0.93	0.25	30,30,30,30	0
4	CIQ	A	2000	37/37	0.93	0.24	44,79,172,177	0
3	SAH	B	1198	26/26	0.93	0.15	50,54,59,59	0
5	UNX	A	4247	1/1	0.94	0.31	30,30,30,30	0
4	CIQ	B	2000	37/37	0.95	0.26	33,58,139,212	0
5	UNX	A	5	1/1	0.95	0.18	30,30,30,30	0
5	UNX	B	4303	1/1	0.96	0.28	30,30,30,30	0
5	UNX	A	8	1/1	0.97	0.11	30,30,30,30	0
5	UNX	A	4263	1/1	0.97	0.12	30,30,30,30	0

Continued on next page...




Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
2	ZN	A	1396	1/1	0.97	0.10	68,68,68,68	0
2	ZN	B	1397	1/1	0.98	0.11	69,69,69,69	0
5	UNX	B	4265	1/1	0.98	0.28	30,30,30,30	0
2	ZN	A	1395	1/1	0.98	0.14	52,52,52,52	0
2	ZN	A	1397	1/1	0.98	0.06	102,102,102,102	0
2	ZN	B	1396	1/1	0.99	0.17	46,46,46,46	0
2	ZN	B	1395	1/1	1.00	0.15	48,48,48,48	0
2	ZN	A	1394	1/1	1.00	0.11	56,56,56,56	0
2	ZN	B	1394	1/1	1.00	0.16	51,51,51,51	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.



Electron density around SAH A 1198:

2mF_o-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)







Electron density around CIQ A 2000:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)

**Electron density around SAH B 1198:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)





6.5 Other polymers [i](#)

There are no such residues in this entry.