



Full wwPDB X-ray Structure Validation Report ⓘ

Jul 22, 2021 – 04:04 PM EDT

PDB ID : 7RJ1
Title : Crystal structure of Aro7p chorismate mutase from *Candida albicans*, complex with L-Trp
Authors : Stogios, P.J.; Evdokimova, E.; Tan, K.; Di Leo, R.; Savchenko, A.; Joachimiak, A.; Satchell, K.J.F.; Center for Structural Genomics of Infectious Diseases (CSGID)
Deposited on : 2021-07-20
Resolution : 2.16 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.22
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.22


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.16 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	1479 (2.16-2.16)
Clashscore	141614	1585 (2.16-2.16)
Ramachandran outliers	138981	1560 (2.16-2.16)
Sidechain outliers	138945	1559 (2.16-2.16)
RSRZ outliers	127900	1456 (2.16-2.16)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	269	 85% 13% .
1	B	269	 82% 16% ..
1	C	269	 81% 16% ..
1	D	269	 80% 17% .

2 Entry composition

There are 6 unique types of molecules in this entry. The entry contains 9162 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Chorismate mutase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	265	Total	C	N	O	S	0	1	0
			2164	1392	352	413	7			
1	B	265	Total	C	N	O	S	0	0	0
			2156	1387	349	413	7			
1	C	263	Total	C	N	O	S	0	0	0
			2138	1376	345	410	7			
1	D	261	Total	C	N	O	S	0	0	0
			2123	1367	342	407	7			

There are 4 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	0	SER	-	expression tag	UNP Q59TS4
B	0	SER	-	expression tag	UNP Q59TS4
C	0	SER	-	expression tag	UNP Q59TS4
D	0	SER	-	expression tag	UNP Q59TS4

- Molecule 2 is TRYPTOPHAN (three-letter code: TRP) (formula: $C_{11}H_{12}N_2O_2$).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
2	A	1	Total	C	N	O	0	0
			15	11	2	2		
2	B	1	Total	C	N	O	0	0
			15	11	2	2		
2	C	1	Total	C	N	O	0	0
			15	11	2	2		
2	D	1	Total	C	N	O	0	0
			15	11	2	2		

- Molecule 3 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	2	Total	Mg	0	0
			2	2		
3	B	1	Total	Mg	0	0
			1	1		
3	C	2	Total	Mg	0	0
			2	2		

- Molecule 4 is 1,2-ETHANEDIOL (three-letter code: EDO) (formula: C₂H₆O₂).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	C	O	0	0
			4	2	2		
4	C	1	Total	C	O	0	0
			4	2	2		

- Molecule 5 is 2-AMINO-2-HYDROXYMETHYL-PROPANE-1,3-DIOL (three-letter code: TRS) (formula: $C_4H_{12}NO_3$).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
5	A	1	Total	C	N	O	0	0
			8	4	1	3		


- Molecule 6 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	159	Total 165	O 165	0	6
6	B	118	Total 126	O 126	0	8
6	C	113	Total 114	O 114	0	1
6	D	93	Total 95	O 95	0	2


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


• Molecule 1: Chorismate mutase




• Molecule 1: Chorismate mutase




• Molecule 1: Chorismate mutase



• Molecule 1: Chorismate mutase





4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	61.75Å 93.15Å 99.71Å 90.00° 106.96° 90.00°	Depositor
Resolution (Å)	41.85 – 2.16 49.88 – 2.16	Depositor EDS
% Data completeness (in resolution range)	97.1 (41.85-2.16) 92.5 (49.88-2.16)	Depositor EDS
R_{merge}	0.08	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.61 (at 2.16Å)	Xtriage
Refinement program	PHENIX 1.19.2_4158	Depositor
R, R_{free}	0.184 , 0.241 0.185 , 0.241	Depositor DCC
R_{free} test set	2010 reflections (3.57%)	wwPDB-VP
Wilson B-factor (Å ²)	37.5	Xtriage
Anisotropy	0.185	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.34 , 57.0	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	0.027 for h,-k,-h-l	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	9162	wwPDB-VP
Average B, all atoms (Å ²)	58.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.95% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: TRS, MG, EDO

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.27	0/2209	0.50	0/2987
1	B	0.29	0/2198	0.51	0/2973
1	C	0.26	0/2180	0.46	0/2950
1	D	0.26	0/2165	0.46	0/2929
All	All	0.27	0/8752	0.48	0/11839

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2164	0	2193	21	0
1	B	2156	0	2181	31	0
1	C	2138	0	2160	27	0
1	D	2123	0	2145	25	0
2	A	15	0	9	2	0
2	B	15	0	9	2	0
2	C	15	0	9	0	0
2	D	15	0	9	0	0
3	A	2	0	0	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	B	1	0	0	0	0
3	C	2	0	0	0	0
4	A	4	0	6	0	0
4	C	4	0	6	0	0
5	A	8	0	12	1	0
6	A	165	0	0	1	0
6	B	126	0	0	1	0
6	C	114	0	0	2	0
6	D	95	0	0	2	0
All	All	9162	0	8739	100	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

All (100) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:260:VAL:O	1:B:260:VAL:HG23	1.78	0.84
1:C:229:GLU:HG3	1:C:233:LYS:HE3	1.67	0.77
1:B:196:GLU:OE2	1:B:244:LYS:NZ	2.22	0.73
1:A:140:SER:HG	2:A:301:TRP:N	1.92	0.67
1:D:202:ARG:NH2	6:D:401:HOH:O	2.26	0.67
1:A:172:PRO:HB3	1:A:265:LYS:HD2	1.77	0.66
1:C:245:LYS:NZ	6:C:402:HOH:O	2.32	0.62
1:D:174:TYR:O	1:D:178:ILE:HG13	2.00	0.61
1:B:5:LYS:NZ	6:B:502[B]:HOH:O	2.34	0.60
1:B:235:TYR:HA	1:B:239:ILE:HB	1.82	0.60
1:C:191:THR:HG23	1:C:244:LYS:HE2	1.84	0.59
1:C:260:VAL:HG12	1:C:263:VAL:HG23	1.85	0.58
1:D:15:ILE:HG21	1:D:162:VAL:HG22	1.84	0.58
1:B:183:VAL:HG13	1:B:248:ILE:HG21	1.85	0.57
1:D:89:LYS:NZ	6:D:405:HOH:O	2.36	0.57
1:A:139:GLY:HA3	2:A:301:TRP:CZ3	2.39	0.56
1:B:260:VAL:O	1:B:260:VAL:CG2	2.51	0.56
1:D:226:VAL:HG11	1:D:231:ILE:HD11	1.88	0.55
1:C:259:ASP:O	1:C:261:GLU:N	2.37	0.55
1:C:178:ILE:HD13	1:C:186:ILE:HD11	1.87	0.54
1:B:199:ILE:HG22	1:B:202:ARG:NH2	2.23	0.54
1:D:167:TYR:HE2	1:D:171:LYS:HE3	1.74	0.53
1:A:71:GLN:HA	1:A:91:PRO:HA	1.90	0.53
1:C:172:PRO:HG3	1:C:265:LYS:HE2	1.89	0.53

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:D:0:SER:HB3	1:D:14:ASN:ND2	2.24	0.53
1:B:172:PRO:HB3	1:B:265:LYS:HE2	1.91	0.52
1:C:236:LYS:NZ	1:C:237:ASP:OD2	2.42	0.52
1:D:244:LYS:O	1:D:248:ILE:HG12	2.10	0.52
1:D:166:LYS:HD2	1:D:190:ILE:HG22	1.92	0.51
1:C:171:LYS:O	1:C:175:ILE:HG13	2.09	0.51
1:D:115:ILE:HA	1:D:118:PHE:CE2	2.46	0.51
1:A:16[B]:ARG:HH12	1:A:195:VAL:HG11	1.76	0.51
1:B:220:GLN:CG	1:C:237:ASP:OD2	2.58	0.51
1:A:205:VAL:HG12	1:B:214:PRO:HB2	1.92	0.51
1:B:27:VAL:HG22	1:B:231:ILE:HD13	1.93	0.50
1:B:115:ILE:HA	1:B:118:PHE:CE2	2.46	0.50
1:A:131:GLN:NE2	6:A:403:HOH:O	2.34	0.50
1:C:34:SER:O	1:C:129:CYS:HB3	2.13	0.49
1:B:140:SER:HG	2:B:301:TRP:N	2.11	0.49
1:A:109:ILE:HB	1:A:160:LYS:HE2	1.94	0.49
1:D:167:TYR:CE2	1:D:171:LYS:HE3	2.48	0.49
1:A:60:TRP:O	1:A:64:GLN:HG2	2.12	0.49
1:B:220:GLN:HG2	1:C:237:ASP:OD2	2.12	0.48
1:C:229:GLU:O	1:C:233:LYS:HG2	2.13	0.48
1:A:16[B]:ARG:HH12	1:A:195:VAL:CG1	2.26	0.48
1:B:139:GLY:HA3	2:B:301:TRP:CZ3	2.47	0.48
1:B:139:GLY:O	1:B:143:THR:HG23	2.12	0.48
1:A:235:TYR:HA	1:A:239:ILE:HB	1.95	0.48
1:D:60:TRP:O	1:D:64:GLN:HG2	2.14	0.47
1:B:53:PHE:CE2	1:B:60:TRP:HB2	2.50	0.47
1:C:36:PHE:CZ	1:C:140:SER:HB3	2.50	0.47
1:D:12:LEU:O	1:D:16:ARG:HG2	2.14	0.47
1:C:174:TYR:O	1:C:178:ILE:HG12	2.13	0.47
1:B:4:MET:O	1:B:5:LYS:HD2	2.15	0.47
1:A:166:LYS:HD3	1:A:190:ILE:HG22	1.97	0.46
1:B:177:LEU:CD1	1:B:182:ASP:HB3	2.45	0.46
1:D:229:GLU:H	1:D:229:GLU:CD	2.19	0.46
1:A:259:ASP:C	1:A:261:GLU:H	2.18	0.46
5:A:305:TRS:N	1:B:134:GLN:HG3	2.31	0.46
1:D:105:TYR:OH	1:D:250:TYR:O	2.24	0.46
1:C:178:ILE:HG21	1:C:255:LEU:HG	1.98	0.46
1:A:122:GLU:O	1:A:126:GLN:HG3	2.16	0.46
1:B:175:ILE:HD12	1:B:175:ILE:H	1.81	0.45
1:D:178:ILE:HG23	1:D:252:LEU:HA	1.98	0.45
1:B:229:GLU:H	1:B:229:GLU:CD	2.19	0.45

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:235:TYR:HA	1:C:239:ILE:HB	1.98	0.45
1:B:229:GLU:OE1	1:B:229:GLU:N	2.47	0.45
1:D:244:LYS:HA	1:D:247:GLU:OE1	2.17	0.45
1:B:198:LYS:HA	1:B:198:LYS:HD3	1.64	0.45
1:C:171:LYS:NZ	6:C:406:HOH:O	2.50	0.44
1:C:227:LYS:HA	1:C:227:LYS:HD3	1.64	0.44
1:B:11:ASP:HB3	1:B:14:ASN:HB2	1.99	0.44
1:C:187:GLU:O	1:C:191:THR:OG1	2.22	0.44
1:A:217:LYS:HA	1:A:217:LYS:HD3	1.76	0.44
1:D:235:TYR:HA	1:D:239:ILE:HB	2.00	0.44
1:A:15:ILE:HG21	1:A:162:VAL:HG22	2.00	0.43
1:A:214:PRO:HB2	1:B:205:VAL:HG12	2.01	0.43
1:D:259:ASP:OD2	1:D:260:VAL:N	2.51	0.43
1:C:67:VAL:HG13	1:C:89:LYS:HG3	2.01	0.43
1:D:76:GLU:OE1	1:D:96:ILE:N	2.49	0.42
1:D:227:LYS:HA	1:D:228:PRO:HD3	1.92	0.42
1:C:244:LYS:O	1:C:248:ILE:HG13	2.18	0.42
1:C:12:LEU:HD12	1:C:12:LEU:H	1.85	0.42
1:A:259:ASP:O	1:A:261:GLU:N	2.48	0.42
1:C:255:LEU:HD13	1:C:262:LEU:HD22	2.00	0.42
1:A:70:SER:HB3	1:A:88:LEU:HD22	2.01	0.41
1:B:263:VAL:HA	1:B:266:TYR:HB2	2.02	0.41
1:D:71:GLN:HA	1:D:91:PRO:HA	2.00	0.41
1:D:171:LYS:O	1:D:175:ILE:HG13	2.19	0.41
1:D:182:ASP:OD2	1:D:182:ASP:N	2.53	0.41
1:D:173:LEU:O	1:D:177:LEU:HG	2.20	0.41
1:B:60:TRP:O	1:B:64:GLN:HG2	2.20	0.41
1:A:115:ILE:HA	1:A:118:PHE:CE2	2.56	0.41
1:C:199:ILE:HG12	1:C:202:ARG:NH2	2.36	0.41
1:B:162:VAL:O	1:B:166:LYS:HG3	2.20	0.41
1:B:171:LYS:O	1:B:175:ILE:HD12	2.21	0.41
1:B:265:LYS:N	1:B:265:LYS:HD2	2.35	0.41
1:C:89:LYS:HB3	1:C:89:LYS:HE3	1.66	0.41
1:C:32:GLU:O	1:C:35:GLN:HG3	2.21	0.40
1:A:53:PHE:CE2	1:A:60:TRP:HB2	2.57	0.40

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	262/269 (97%)	250 (95%)	12 (5%)	0	100	100
1	B	261/269 (97%)	254 (97%)	7 (3%)	0	100	100
1	C	259/269 (96%)	248 (96%)	11 (4%)	0	100	100
1	D	257/269 (96%)	246 (96%)	11 (4%)	0	100	100
All	All	1039/1076 (97%)	998 (96%)	41 (4%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	246/248 (99%)	239 (97%)	7 (3%)	43	44
1	B	245/248 (99%)	237 (97%)	8 (3%)	38	37
1	C	243/248 (98%)	233 (96%)	10 (4%)	30	29
1	D	241/248 (97%)	235 (98%)	6 (2%)	47	49
All	All	975/992 (98%)	944 (97%)	31 (3%)	39	38

All (31) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	24	ASP
1	A	131	GLN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	158	PHE
1	A	176	LYS
1	A	220	GLN
1	A	227	LYS
1	A	266	TYR
1	B	106	SER
1	B	158	PHE
1	B	187	GLU
1	B	189	SER
1	B	220	GLN
1	B	221	ASN
1	B	229	GLU
1	B	261	GLU
1	C	5	LYS
1	C	126	GLN
1	C	158	PHE
1	C	209	SER
1	C	215	SER
1	C	216	LEU
1	C	223	GLN
1	C	260	VAL
1	C	261	GLU
1	C	266	TYR
1	D	133	ASP
1	D	155	ARG
1	D	158	PHE
1	D	176	LYS
1	D	233	LYS
1	D	266	TYR

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 12 ligands modelled in this entry, 5 are monoatomic - leaving 7 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	TRP	D	301	-	12,16,16	0.65	0	12,22,22	0.84	0
5	TRS	A	305	-	7,7,7	0.36	0	9,9,9	0.63	0
2	TRP	B	301	-	12,16,16	0.69	0	12,22,22	0.85	0
4	EDO	C	304	-	3,3,3	0.45	0	2,2,2	0.33	0
2	TRP	A	301	-	12,16,16	0.65	0	12,22,22	0.86	0
2	TRP	C	301	-	12,16,16	0.66	0	12,22,22	0.81	0
4	EDO	A	304	-	3,3,3	0.45	0	2,2,2	0.31	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	TRP	D	301	-	-	0/3/8/8	0/2/2/2
5	TRS	A	305	-	-	3/9/9/9	-
2	TRP	B	301	-	-	0/3/8/8	0/2/2/2
4	EDO	C	304	-	-	0/1/1/1	-
2	TRP	A	301	-	-	0/3/8/8	0/2/2/2
2	TRP	C	301	-	-	0/3/8/8	0/2/2/2
4	EDO	A	304	-	-	1/1/1/1	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (4) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
5	A	305	TRS	C1-C-C2-O2
5	A	305	TRS	C3-C-C2-O2
5	A	305	TRS	N-C-C2-O2
4	A	304	EDO	O1-C1-C2-O2

There are no ring outliers.

3 monomers are involved in 5 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
5	A	305	TRS	1	0
2	B	301	TRP	2	0
2	A	301	TRP	2	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	265/269 (98%)	-0.10	4 (1%) 73 79	31, 44, 70, 146	0
1	B	265/269 (98%)	0.07	12 (4%) 33 42	32, 50, 103, 135	0
1	C	263/269 (97%)	0.07	7 (2%) 54 63	35, 59, 90, 148	0
1	D	261/269 (97%)	0.23	15 (5%) 23 32	35, 61, 106, 154	0
All	All	1054/1076 (97%)	0.06	38 (3%) 42 51	31, 53, 98, 154	0

All (38) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	D	266	TYR	6.6
1	C	222	VAL	6.3
1	D	178	ILE	5.8
1	B	260	VAL	5.2
1	B	262	LEU	4.5
1	D	179	LEU	4.2
1	D	223	GLN	4.0
1	D	186	ILE	4.0
1	C	221	ASN	3.8
1	C	223	GLN	3.7
1	B	199	ILE	3.2
1	A	220	GLN	3.1
1	D	216	LEU	3.0
1	C	216	LEU	3.0
1	D	183	VAL	2.9
1	D	167	TYR	2.9
1	D	175	ILE	2.9
1	C	266	TYR	2.8
1	A	266	TYR	2.7
1	B	220	GLN	2.7
1	D	12	LEU	2.7

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	D	214	PRO	2.7
1	C	212	VAL	2.6
1	A	261	GLU	2.5
1	B	217	LYS	2.5
1	B	183	VAL	2.5
1	B	178	ILE	2.4
1	D	261	GLU	2.4
1	B	0	SER	2.3
1	B	181	LYS	2.2
1	D	181	LYS	2.2
1	B	167	TYR	2.2
1	D	190	ILE	2.1
1	A	221	ASN	2.1
1	D	255	LEU	2.1
1	C	12	LEU	2.0
1	B	166	LYS	2.0
1	B	186	ILE	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
3	MG	C	302	1/1	0.56	0.12	83,83,83,83	0
3	MG	B	302	1/1	0.78	0.10	64,64,64,64	0
3	MG	A	303	1/1	0.78	0.21	70,70,70,70	0
4	EDO	C	304	4/4	0.80	0.21	58,73,78,79	0
3	MG	C	303	1/1	0.82	0.23	79,79,79,79	0
4	EDO	A	304	4/4	0.84	0.21	58,60,61,73	0

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
5	TRS	A	305	8/8	0.86	0.18	55,65,82,83	0
2	TRP	C	301	15/15	0.90	0.23	47,52,61,62	0
2	TRP	D	301	15/15	0.92	0.19	70,75,83,91	0
2	TRP	B	301	15/15	0.94	0.16	44,50,59,64	0
2	TRP	A	301	15/15	0.94	0.13	50,54,62,64	0
3	MG	A	302	1/1	0.96	0.11	58,58,58,58	0

6.5 Other polymers [i](#)

There are no such residues in this entry.