



# wwPDB X-ray Structure Validation Summary Report ⓘ

Oct 31, 2021 – 04:08 AM EDT

PDB ID : 2RKW  
Title : Intermediate position of ATP on its trail to the binding pocket inside the subunit B mutant R416W of the energy converter A1Ao ATP synthase  
Authors : Kumar, A.; Manimekalai, M.S.S.; Balakrishna, A.M.; Hunke, C.; Gruber, G.  
Deposited on : 2007-10-18  
Resolution : 2.81 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Xtriage (Phenix)	:	1.13
EDS	:	2.23.2
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac	:	5.8.0158
CCP4	:	7.0.044 (Gargrove)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.23.2


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

## *X-RAY DIFFRACTION*

The reported resolution of this entry is 2.81 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	3617 (2.84-2.80)
Clashscore	141614	4060 (2.84-2.80)
Ramachandran outliers	138981	3978 (2.84-2.80)
Sidechain outliers	138945	3980 (2.84-2.80)
RSRZ outliers	127900	3552 (2.84-2.80)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	469	 6% 52% 29% 9% 10%
1	B	469	 5% 48% 32% 8% 10%

## 2 Entry composition

There is only 1 type of molecule in this entry. The entry contains 6512 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called V-type ATP synthase beta chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	422	Total	C	N	O	S	0	0	0
			3257	2066	561	619	11			
1	B	422	Total	C	N	O	S	0	0	0
			3255	2062	562	620	11			


There are 22 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-8	MET	-	expression tag	UNP Q60187
A	-7	LYS	-	expression tag	UNP Q60187
A	-6	HIS	-	expression tag	UNP Q60187
A	-5	HIS	-	expression tag	UNP Q60187
A	-4	HIS	-	expression tag	UNP Q60187
A	-3	HIS	-	expression tag	UNP Q60187
A	-2	HIS	-	expression tag	UNP Q60187
A	-1	HIS	-	expression tag	UNP Q60187
A	0	PRO	-	expression tag	UNP Q60187
A	2	VAL	ALA	SEE REMARK 999	UNP Q60187
A	416	TRP	ARG	engineered mutation	UNP Q60187
B	-8	MET	-	expression tag	UNP Q60187
B	-7	LYS	-	expression tag	UNP Q60187
B	-6	HIS	-	expression tag	UNP Q60187
B	-5	HIS	-	expression tag	UNP Q60187
B	-4	HIS	-	expression tag	UNP Q60187
B	-3	HIS	-	expression tag	UNP Q60187
B	-2	HIS	-	expression tag	UNP Q60187
B	-1	HIS	-	expression tag	UNP Q60187
B	0	PRO	-	expression tag	UNP Q60187
B	2	VAL	ALA	SEE REMARK 999	UNP Q60187
B	416	TRP	ARG	engineered mutation	UNP Q60187



- Molecule 1: V-type ATP synthase beta chain





## 4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	73.05Å 96.09Å 129.94Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	34.73 – 2.81 34.15 – 2.81	Depositor EDS
% Data completeness (in resolution range)	86.9 (34.73-2.81) 86.9 (34.15-2.81)	Depositor EDS
$R_{merge}$	0.13	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	4.33 (at 2.81Å)	Xtriage
Refinement program	REFMAC 5.2.0019	Depositor
R, $R_{free}$	0.271 , 0.330 0.269 , 0.337	Depositor DCC
$R_{free}$ test set	1033 reflections (5.17%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	80.2	Xtriage
Anisotropy	0.106	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.35 , 50.1	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.47$ , $\langle L^2 \rangle = 0.30$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.91	EDS
Total number of atoms	6512	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	79.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.89% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.39	0/3320	0.63	1/4504 (0.0%)
1	B	0.41	0/3316	0.63	0/4497
All	All	0.40	0/6636	0.63	1/9001 (0.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	B	0	3

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed( $^{\circ}$ )	Ideal( $^{\circ}$ )
1	A	393	LEU	CA-CB-CG	5.48	127.90	115.30

There are no chirality outliers.

All (3) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	B	314	ILE	Peptide
1	B	315	PRO	Peptide
1	B	344	LEU	Peptide

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within

the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3257	0	3259	130	0
1	B	3255	0	3257	166	0
All	All	6512	0	6516	288	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 22.

The worst 5 of 288 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:71:PHE:HA	1:B:72:THR:HB	1.21	1.16
1:B:41:ARG:HB3	1:B:56:VAL:CG2	1.87	1.05
1:A:314:ILE:HB	1:A:315:PRO:HA	1.38	1.04
1:B:71:PHE:HA	1:B:72:THR:CB	1.87	1.03
1:A:99:GLY:H	1:A:100:PRO:HD3	1.20	1.02

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	416/469 (89%)	340 (82%)	54 (13%)	22 (5%)	2	5
1	B	416/469 (89%)	323 (78%)	66 (16%)	27 (6%)	1	3
All	All	832/938 (89%)	663 (80%)	120 (14%)	49 (6%)	1	4

5 of 49 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	94	GLU
1	A	313	PRO

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	A	314	ILE
1	A	339	PRO
1	B	14	PRO

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	346/384 (90%)	293 (85%)	53 (15%)	2	8
1	B	346/384 (90%)	297 (86%)	49 (14%)	3	10
All	All	692/768 (90%)	590 (85%)	102 (15%)	3	9

5 of 102 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	41	ARG
1	B	133	THR
1	B	422	ILE
1	B	50	ASP
1	B	77	LYS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 20 such sidechains are listed below:

Mol	Chain	Res	Type
1	B	211	ASN
1	B	417	ASN
1	B	450	GLN
1	B	439	ASN
1	A	325	GLN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 5.6 Ligand geometry [i](#)

There are no ligands in this entry.

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ > 2		OWAB(Å <sup>2</sup> )	Q < 0.9
1	A	422/469 (89%)	0.33	28 (6%)	18 11	62, 76, 103, 109	0
1	B	422/469 (89%)	0.36	22 (5%)	27 18	64, 76, 93, 104	0
All	All	844/938 (89%)	0.34	50 (5%)	22 14	62, 76, 102, 109	0

The worst 5 of 50 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	310	ILE	8.2
1	B	261	ALA	5.8
1	B	455	ALA	5.7
1	B	357	ALA	5.0
1	B	11	ILE	4.8

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

### 6.4 Ligands [i](#)

There are no ligands in this entry.

### 6.5 Other polymers [i](#)

There are no such residues in this entry.