



Full wwPDB X-ray Structure Validation Report ⓘ

Aug 9, 2020 – 04:34 PM BST

PDB ID : 4RK0
Title : Crystal structure of LacI family transcriptional regulator from *Enterococcus faecalis* V583, Target EFI-512923, with bound ribose
Authors : Patskovsky, Y.; Toro, R.; Bhosle, R.; Al Obaidi, N.; Chamala, S.; Attonito, J.D.; Scott Glenn, A.; Chowdhury, S.; Lafleur, J.; Siedel, R.D.; Hillerich, B.; Love, J.; Whalen, K.L.; Gerlt, J.A.; Almo, S.C.; Enzyme Function Initiative (EFI)
Deposited on : 2014-10-11
Resolution : 1.80 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.13.1
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.13.1


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.80 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	5950 (1.80-1.80)
Clashscore	141614	6793 (1.80-1.80)
Ramachandran outliers	138981	6697 (1.80-1.80)
Sidechain outliers	138945	6696 (1.80-1.80)
RSRZ outliers	127900	5850 (1.80-1.80)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	274	
1	B	274	
1	C	274	
1	D	274	

2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 9006 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called LacI family sugar-binding transcriptional regulator.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	268	Total	C	N	O	S	0	1	0
			2122	1356	353	408	5			
1	B	267	Total	C	N	O	S	0	4	0
			2137	1367	355	410	5			
1	C	268	Total	C	N	O	S	0	2	0
			2134	1364	357	408	5			
1	D	269	Total	C	N	O	S	0	1	0
			2128	1360	354	409	5			

There are 8 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	63	SER	-	expression tag	UNP Q82ZT5
A	103	ILE	MET	conflict	UNP Q82ZT5
B	63	SER	-	expression tag	UNP Q82ZT5
B	103	ILE	MET	conflict	UNP Q82ZT5
C	63	SER	-	expression tag	UNP Q82ZT5
C	103	ILE	MET	conflict	UNP Q82ZT5
D	63	SER	-	expression tag	UNP Q82ZT5
D	103	ILE	MET	conflict	UNP Q82ZT5

- Molecule 2 is alpha-D-ribofuranose (three-letter code: RIB) (formula: C₅H₁₀O₅).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	C	O	0	0
			10	5	5		
2	B	1	Total	C	O	0	0
			10	5	5		
2	C	1	Total	C	O	0	0
			10	5	5		
2	D	1	Total	C	O	0	0
			10	5	5		

- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	113	Total	O	0	0
			113	113		
3	B	95	Total	O	0	0
			95	95		
3	C	133	Total	O	0	0
			133	133		
3	D	104	Total	O	0	0
			104	104		

3 Residue-property plots [i](#)


These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: LacI family sugar-binding transcriptional regulator

Chain A: 



- Molecule 1: LacI family sugar-binding transcriptional regulator

Chain B: 



- Molecule 1: LacI family sugar-binding transcriptional regulator

Chain C: 



- Molecule 1: LacI family sugar-binding transcriptional regulator

Chain D: 



4 Data and refinement statistics

Property	Value	Source
Space group	P 1	Depositor
Cell constants a, b, c, α , β , γ	36.92Å 77.23Å 91.81Å 87.72° 82.57° 84.33°	Depositor
Resolution (Å)	50.00 – 1.80 45.50 – 1.79	Depositor EDS
% Data completeness (in resolution range)	97.7 (50.00-1.80) 97.5 (45.50-1.79)	Depositor EDS
R_{merge}	0.10	Depositor
R_{sym}	0.10	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.18 (at 1.79Å)	Xtriage
Refinement program	REFMAC 5.8.0073	Depositor
R, R_{free}	0.170 , 0.223 0.179 , 0.228	Depositor DCC
R_{free} test set	2769 reflections (3.03%)	wwPDB-VP
Wilson B-factor (Å ²)	27.5	Xtriage
Anisotropy	0.256	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.34 , 43.3	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.97	EDS
Total number of atoms	9006	wwPDB-VP
Average B, all atoms (Å ²)	39.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.84% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: RIB

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.58	0/2165	0.71	0/2941
1	B	0.57	0/2189	0.75	1/2972 (0.0%)
1	C	0.63	0/2180	0.76	1/2959 (0.0%)
1	D	0.60	0/2171	0.73	1/2949 (0.0%)
All	All	0.59	0/8705	0.74	3/11821 (0.0%)

There are no bond length outliers.

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	C	279	ASP	CB-CG-OD1	5.95	123.66	118.30
1	D	249	ASP	CB-CG-OD1	5.19	122.97	118.30
1	B	249	ASP	CB-CG-OD1	5.12	122.91	118.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2122	0	2137	17	0
1	B	2137	0	2163	14	0
1	C	2134	0	2161	15	0
1	D	2128	0	2147	22	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
2	A	10	0	0	3	0
2	B	10	0	0	2	0
2	C	10	0	0	2	0
2	D	10	0	0	2	0
3	A	113	0	0	0	0
3	B	95	0	0	0	0
3	C	133	0	0	1	0
3	D	104	0	0	2	0
All	All	9006	0	8608	58	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (58) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:D:191:ILE:HD13	1:D:218:GLN:HB3	1.65	0.79
1:B:81:ASN:HD21	2:B:401:RIB:C5	2.06	0.68
1:A:81:ASN:HD21	2:A:401:RIB:C5	2.08	0.67
1:B:81:ASN:ND2	2:B:401:RIB:C5	2.63	0.62
1:C:124[B]:ARG:HG3	1:D:101:ILE:HD11	1.82	0.61
1:D:315:LYS:HE3	3:D:522:HOH:O	2.04	0.58
1:A:194:MET:O	1:A:221:GLU:HA	2.05	0.55
1:D:81:ASN:HD21	2:D:401:RIB:C5	2.20	0.54
1:C:124[B]:ARG:NH2	3:C:577:HOH:O	2.40	0.54
1:A:75:ILE:HD13	1:A:105:CYS:HB3	1.89	0.54
1:A:103:ILE:HG23	1:B:101:ILE:HG23	1.90	0.53
1:B:194:MET:HG2	1:B:219:LEU:HD11	1.90	0.53
1:A:81:ASN:ND2	2:A:401:RIB:C5	2.71	0.53
1:D:70:LYS:HA	1:D:127:ASP:OD2	2.10	0.51
1:C:81:ASN:HD21	2:C:401:RIB:C5	2.23	0.51
1:D:86:GLN:OE1	1:D:89:ARG:NH1	2.44	0.51
1:D:142:ASN:OD1	1:D:146:ARG:NH1	2.38	0.51
1:C:307:LEU:HD23	1:C:310:ILE:HD12	1.94	0.49
1:B:71:THR:HG22	1:B:101:ILE:HB	1.92	0.49
1:D:94:VAL:HG23	1:D:300:GLN:HG3	1.94	0.49
1:A:191:ILE:HD13	1:A:218:GLN:HB3	1.95	0.48
1:A:103:ILE:HD12	1:B:101:ILE:CG2	2.42	0.48
1:A:73:GLY:HA2	1:A:103:ILE:HB	1.94	0.48
1:C:315:LYS:NZ	1:C:318:GLU:OE1	2.37	0.48
1:B:197:HIS:O	1:B:197:HIS:CG	2.67	0.47

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:D:78:ASP:HB3	1:D:108:ASP:OD2	2.15	0.47
1:A:276:ASP:OD2	2:A:401:RIB:O5	2.32	0.47
1:C:101:ILE:CG2	1:D:103:ILE:HG23	2.44	0.47
1:C:194:MET:O	1:C:221:GLU:HA	2.15	0.47
1:C:89:ARG:HD2	1:D:80:THR:HG23	1.95	0.46
1:B:312:GLN:HE21	1:B:312:GLN:HA	1.80	0.46
1:D:81:ASN:ND2	2:D:401:RIB:C5	2.78	0.46
1:D:146:ARG:NH2	1:D:163:PHE:O	2.44	0.46
1:C:101:ILE:HG22	1:D:103:ILE:HG23	1.98	0.46
1:D:94:VAL:CG2	1:D:300:GLN:HG3	2.47	0.45
1:A:164:SER:HA	1:A:317:TRP:CE3	2.52	0.45
1:A:194:MET:HG2	1:A:219:LEU:HD11	1.98	0.45
1:B:312:GLN:HA	1:B:312:GLN:NE2	2.32	0.44
1:C:195:PRO:HA	1:C:222:THR:O	2.17	0.44
1:D:106:ASN:ND2	3:D:536:HOH:O	2.49	0.44
1:A:195:PRO:HA	1:A:222:THR:O	2.18	0.44
1:D:168:LEU:N	1:D:168:LEU:HD12	2.32	0.44
1:A:312:GLN:CG	1:A:312:GLN:O	2.66	0.43
1:C:234:PRO:O	1:C:238:LYS:HD3	2.18	0.43
1:C:89:ARG:HD2	1:D:80:THR:CG2	2.49	0.43
1:C:81:ASN:ND2	2:C:401:RIB:C5	2.83	0.42
1:C:124[A]:ARG:HG3	1:D:101:ILE:HD11	2.01	0.42
1:C:103:ILE:HG23	1:D:101:ILE:HG23	2.01	0.42
1:B:289:THR:HG23	1:B:328:GLU:HA	2.02	0.42
1:A:71:THR:HG22	1:A:101:ILE:HD11	2.01	0.41
1:A:105:CYS:HB2	1:A:121:LEU:CD2	2.50	0.41
1:B:195:PRO:HA	1:B:222:THR:O	2.20	0.41
1:B:163:PHE:HA	1:B:317:TRP:CD1	2.54	0.41
1:D:233:VAL:HG21	1:D:258:GLY:HA3	2.01	0.41
1:A:194:MET:SD	1:A:219:LEU:HD11	2.61	0.41
1:D:267:PRO:HA	1:D:270:TYR:O	2.21	0.41
1:A:106:ASN:OD1	1:B:92:GLU:OE2	2.40	0.40
1:B:189[A]:GLN:HE22	1:B:240:GLU:HB2	1.87	0.40

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	267/274 (97%)	259 (97%)	7 (3%)	1 (0%)	34	21
1	B	269/274 (98%)	260 (97%)	7 (3%)	2 (1%)	22	10
1	C	268/274 (98%)	260 (97%)	7 (3%)	1 (0%)	34	21
1	D	268/274 (98%)	261 (97%)	6 (2%)	1 (0%)	34	21
All	All	1072/1096 (98%)	1040 (97%)	27 (2%)	5 (0%)	29	15

All (5) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	276	ASP
1	B	276	ASP
1	C	276	ASP
1	D	276	ASP
1	B	315	LYS

5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	235/241 (98%)	230 (98%)	5 (2%)	53	42
1	B	238/241 (99%)	233 (98%)	5 (2%)	53	42
1	C	237/241 (98%)	235 (99%)	2 (1%)	81	78
1	D	236/241 (98%)	234 (99%)	2 (1%)	81	78
All	All	946/964 (98%)	932 (98%)	14 (2%)	65	56

All (14) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	125	SER
1	A	315	LYS
1	A	316	ASP
1	A	319	GLU
1	A	335	LEU
1	B	111	VAL
1	B	113	ARG
1	B	122	ILE
1	B	312	GLN
1	B	315	LYS
1	C	69	SER
1	C	247	ILE
1	D	121	LEU
1	D	312	GLN

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (10) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	99	ASN
1	A	203	GLN
1	B	203	GLN
1	B	218	GLN
1	B	312	GLN
1	C	203	GLN
1	C	231	GLN
1	D	197	HIS
1	D	218	GLN
1	D	231	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

4 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z > 2$	Counts	RMSZ	# $ Z > 2$
2	RIB	C	401	-	10,10,10	1.58	2 (20%)	13,14,14	4.87	8 (61%)
2	RIB	B	401	-	10,10,10	0.95	0	13,14,14	2.90	7 (53%)
2	RIB	D	401	-	10,10,10	1.00	1 (10%)	13,14,14	3.61	8 (61%)
2	RIB	A	401	-	10,10,10	0.75	0	13,14,14	3.67	8 (61%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	RIB	C	401	-	-	1/2/18/18	0/1/1/1
2	RIB	B	401	-	-	0/2/18/18	0/1/1/1
2	RIB	D	401	-	-	0/2/18/18	0/1/1/1
2	RIB	A	401	-	-	0/2/18/18	0/1/1/1

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	C	401	RIB	C1-C2	3.86	1.57	1.52
2	C	401	RIB	C5-C4	-2.65	1.43	1.51
2	D	401	RIB	C5-C4	-2.17	1.44	1.51

All (31) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	C	401	RIB	C5-C4-C3	-9.95	91.11	115.09
2	C	401	RIB	O1-C1-O4	8.44	121.94	111.13
2	A	401	RIB	C5-C4-C3	-7.68	96.56	115.09
2	C	401	RIB	O4-C4-C3	6.73	118.43	105.11
2	D	401	RIB	C5-C4-C3	-6.69	98.96	115.09
2	C	401	RIB	O4-C1-C2	6.44	112.39	104.46
2	D	401	RIB	O4-C4-C3	5.94	116.87	105.11
2	D	401	RIB	O4-C1-C2	5.73	111.52	104.46
2	A	401	RIB	O4-C4-C3	5.41	115.82	105.11
2	A	401	RIB	O4-C1-C2	4.94	110.54	104.46
2	B	401	RIB	O4-C1-C2	4.74	110.30	104.46
2	B	401	RIB	C1-C2-C3	4.43	107.84	102.30
2	D	401	RIB	C1-C2-C3	4.43	107.84	102.30
2	A	401	RIB	C1-C2-C3	4.38	107.78	102.30
2	B	401	RIB	C5-C4-C3	-4.33	104.65	115.09
2	A	401	RIB	O1-C1-O4	4.19	116.50	111.13
2	B	401	RIB	O4-C4-C3	3.99	113.01	105.11
2	B	401	RIB	O5-C5-C4	-3.94	97.76	111.29
2	C	401	RIB	O3-C3-C4	-3.84	99.95	111.05
2	C	401	RIB	O2-C2-C1	3.48	121.43	111.82
2	C	401	RIB	C1-C2-C3	3.33	106.47	102.30
2	A	401	RIB	O3-C3-C4	-2.91	102.64	111.05
2	D	401	RIB	O3-C3-C2	2.90	121.21	111.82
2	B	401	RIB	O4-C4-C5	2.82	115.32	109.21
2	D	401	RIB	O2-C2-C1	2.62	119.06	111.82
2	D	401	RIB	O3-C3-C4	-2.42	104.06	111.05
2	A	401	RIB	O3-C3-C2	2.41	119.60	111.82
2	D	401	RIB	O1-C1-O4	2.32	114.10	111.13
2	A	401	RIB	O4-C4-C5	2.28	114.13	109.21
2	B	401	RIB	O1-C1-O4	2.08	113.79	111.13
2	C	401	RIB	O5-C5-C4	-2.05	104.25	111.29

There are no chirality outliers.

All (1) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	C	401	RIB	O4-C4-C5-O5

There are no ring outliers.

4 monomers are involved in 9 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	C	401	RIB	2	0
2	B	401	RIB	2	0
2	D	401	RIB	2	0
2	A	401	RIB	3	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	268/274 (97%)	-0.51	0 100 100	23, 36, 63, 98	0
1	B	267/274 (97%)	-0.37	0 100 100	21, 37, 68, 92	0
1	C	268/274 (97%)	-0.61	1 (0%) 92 90	21, 30, 57, 92	0
1	D	269/274 (98%)	-0.45	0 100 100	24, 35, 67, 99	0
All	All	1072/1096 (97%)	-0.48	1 (0%) 95 93	21, 35, 66, 99	0

All (1) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	C	336	LYS	2.4

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	RIB	B	401	10/10	0.91	0.12	26,31,44,69	0

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
2	RIB	D	401	10/10	0.92	0.11	24,29,43,60	0
2	RIB	C	401	10/10	0.94	0.11	20,23,29,51	0
2	RIB	A	401	10/10	0.94	0.11	22,29,43,60	0

6.5 Other polymers [i](#)

There are no such residues in this entry.