



wwPDB X-ray Structure Validation Summary Report ⓘ

Feb 3, 2022 – 12:29 PM EST

PDB ID : 7RM4
Title : Neoantigen p53R175H-specific TCR 6-11 binds to p53R175H-HLA-A2
Authors : Wu, D.; Mariuzza, R.A.
Deposited on : 2021-07-26
Resolution : 3.33 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Xtriage (Phenix)	:	1.13
EDS	:	2.26
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac	:	5.8.0158
CCP4	:	7.0.044 (Gargrove)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.26


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 3.33 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	1060 (3.38-3.30)
Clashscore	141614	1111 (3.38-3.30)
Ramachandran outliers	138981	1090 (3.38-3.30)
Sidechain outliers	138945	1089 (3.38-3.30)
RSRZ outliers	127900	1028 (3.38-3.30)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	275	<div> <div></div> <div>75%</div> <div>24%</div> <div>.</div> </div>
1	F	275	<div> <div></div> <div>81%</div> <div>19%</div> </div>
1	K	275	<div> <div></div> <div>82%</div> <div>17%</div> <div>.</div> </div>
1	P	275	<div> <div></div> <div>81%</div> <div>19%</div> <div>.</div> </div>
2	B	100	<div> <div></div> <div>79%</div> <div>21%</div> </div>

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
2	G	100	 78%20%.
2	L	100	 88%12%
2	Q	100	 81%17%.
3	C	9	 22%78%
3	H	9	 56%44%
3	M	9	 56%44%
3	R	9	 67%33%
4	D	246	 71%26%..
4	I	246	 72%27%.
4	N	246	 73%25%.
4	S	246	 69%29%..
5	E	206	 68%25%..
5	J	206	 76%20%..
5	O	206	 67%29%..
5	T	206	 70%26%..

2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 26474 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called HLA class I histocompatibility antigen, A alpha chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	275	Total	C	N	O	S	0	0	0
			2240	1400	406	425	9			
1	F	275	Total	C	N	O	S	0	0	0
			2236	1398	406	423	9			
1	K	275	Total	C	N	O	S	0	0	0
			2236	1398	406	423	9			
1	P	275	Total	C	N	O	S	0	0	0
			2236	1398	406	423	9			

There are 76 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	62	GLY	GLN	conflict	UNP P04439
A	66	LYS	ASN	conflict	UNP P04439
A	70	HIS	GLN	conflict	UNP P04439
A	74	HIS	ASP	conflict	UNP P04439
A	95	VAL	ILE	conflict	UNP P04439
A	97	ARG	ILE	conflict	UNP P04439
A	107	TRP	GLY	conflict	UNP P04439
A	114	HIS	ARG	conflict	UNP P04439
A	116	TYR	ASP	conflict	UNP P04439
A	127	LYS	ASN	conflict	UNP P04439
A	142	THR	ILE	conflict	UNP P04439
A	145	HIS	ARG	conflict	UNP P04439
A	152	VAL	GLU	conflict	UNP P04439
A	161	GLU	ASP	conflict	UNP P04439
A	184	ALA	PRO	conflict	UNP P04439
A	193	ALA	PRO	conflict	UNP P04439
A	194	VAL	ILE	conflict	UNP P04439
A	207	SER	GLY	conflict	UNP P04439
A	253	GLN	GLU	conflict	UNP P04439
F	62	GLY	GLN	conflict	UNP P04439
F	66	LYS	ASN	conflict	UNP P04439

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
F	70	HIS	GLN	conflict	UNP P04439
F	74	HIS	ASP	conflict	UNP P04439
F	95	VAL	ILE	conflict	UNP P04439
F	97	ARG	ILE	conflict	UNP P04439
F	107	TRP	GLY	conflict	UNP P04439
F	114	HIS	ARG	conflict	UNP P04439
F	116	TYR	ASP	conflict	UNP P04439
F	127	LYS	ASN	conflict	UNP P04439
F	142	THR	ILE	conflict	UNP P04439
F	145	HIS	ARG	conflict	UNP P04439
F	152	VAL	GLU	conflict	UNP P04439
F	161	GLU	ASP	conflict	UNP P04439
F	184	ALA	PRO	conflict	UNP P04439
F	193	ALA	PRO	conflict	UNP P04439
F	194	VAL	ILE	conflict	UNP P04439
F	207	SER	GLY	conflict	UNP P04439
F	253	GLN	GLU	conflict	UNP P04439
K	62	GLY	GLN	conflict	UNP P04439
K	66	LYS	ASN	conflict	UNP P04439
K	70	HIS	GLN	conflict	UNP P04439
K	74	HIS	ASP	conflict	UNP P04439
K	95	VAL	ILE	conflict	UNP P04439
K	97	ARG	ILE	conflict	UNP P04439
K	107	TRP	GLY	conflict	UNP P04439
K	114	HIS	ARG	conflict	UNP P04439
K	116	TYR	ASP	conflict	UNP P04439
K	127	LYS	ASN	conflict	UNP P04439
K	142	THR	ILE	conflict	UNP P04439
K	145	HIS	ARG	conflict	UNP P04439
K	152	VAL	GLU	conflict	UNP P04439
K	161	GLU	ASP	conflict	UNP P04439
K	184	ALA	PRO	conflict	UNP P04439
K	193	ALA	PRO	conflict	UNP P04439
K	194	VAL	ILE	conflict	UNP P04439
K	207	SER	GLY	conflict	UNP P04439
K	253	GLN	GLU	conflict	UNP P04439
P	62	GLY	GLN	conflict	UNP P04439
P	66	LYS	ASN	conflict	UNP P04439
P	70	HIS	GLN	conflict	UNP P04439
P	74	HIS	ASP	conflict	UNP P04439
P	95	VAL	ILE	conflict	UNP P04439
P	97	ARG	ILE	conflict	UNP P04439

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
P	107	TRP	GLY	conflict	UNP P04439
P	114	HIS	ARG	conflict	UNP P04439
P	116	TYR	ASP	conflict	UNP P04439
P	127	LYS	ASN	conflict	UNP P04439
P	142	THR	ILE	conflict	UNP P04439
P	145	HIS	ARG	conflict	UNP P04439
P	152	VAL	GLU	conflict	UNP P04439
P	161	GLU	ASP	conflict	UNP P04439
P	184	ALA	PRO	conflict	UNP P04439
P	193	ALA	PRO	conflict	UNP P04439
P	194	VAL	ILE	conflict	UNP P04439
P	207	SER	GLY	conflict	UNP P04439
P	253	GLN	GLU	conflict	UNP P04439

- Molecule 2 is a protein called Beta-2-microglobulin.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	100	Total	C	N	O	S	0	0	0
			824	525	138	157	4			
2	G	100	Total	C	N	O	S	0	0	0
			824	525	138	157	4			
2	L	100	Total	C	N	O	S	0	0	0
			824	525	138	157	4			
2	Q	100	Total	C	N	O	S	0	0	0
			820	523	138	155	4			

There are 4 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	1	MET	-	initiating methionine	UNP P61769
G	1	MET	-	initiating methionine	UNP P61769
L	1	MET	-	initiating methionine	UNP P61769
Q	1	MET	-	initiating methionine	UNP P61769

- Molecule 3 is a protein called Cellular tumor antigen p53 peptide.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	C	9	Total	C	N	O	S	0	0	0
			76	45	16	13	2			
3	H	9	Total	C	N	O	S	0	0	0
			76	45	16	13	2			

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	M	9	Total	C	N	O	S	0	0	0
			76	45	16	13	2			
3	R	9	Total	C	N	O	S	0	0	0
			76	45	16	13	2			

There are 4 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
C	8	HIS	ARG	engineered mutation	UNP P04637
H	8	HIS	ARG	engineered mutation	UNP P04637
M	8	HIS	ARG	engineered mutation	UNP P04637
R	8	HIS	ARG	engineered mutation	UNP P04637

- Molecule 4 is a protein called 6-11 T cell receptor beta chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
4	D	243	Total	C	N	O	S	3	0	0
			1915	1217	330	363	5			
4	I	243	Total	C	N	O	S	3	0	0
			1913	1217	328	363	5			
4	N	243	Total	C	N	O	S	3	0	0
			1913	1217	328	363	5			
4	S	243	Total	C	N	O	S	3	0	0
			1913	1217	328	363	5			


- Molecule 5 is a protein called 6-11 T cell receptor alpha chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
5	E	200	Total	C	N	O	S	0	0	0
			1569	986	261	314	8			
5	J	200	Total	C	N	O	S	0	0	0
			1569	986	261	314	8			
5	O	200	Total	C	N	O	S	0	0	0
			1569	986	261	314	8			
5	T	200	Total	C	N	O	S	0	0	0
			1569	986	261	314	8			


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: HLA class I histocompatibility antigen, A alpha chain




- Molecule 1: HLA class I histocompatibility antigen, A alpha chain




- Molecule 1: HLA class I histocompatibility antigen, A alpha chain





- Molecule 1: HLA class I histocompatibility antigen, A alpha chain







- Molecule 2: Beta-2-microglobulin





- Molecule 2: Beta-2-microglobulin



- Molecule 2: Beta-2-microglobulin



- Molecule 3: Cellular tumor antigen p53 peptide



- Molecule 3: Cellular tumor antigen p53 peptide




- Molecule 3: Cellular tumor antigen p53 peptide




- Molecule 3: Cellular tumor antigen p53 peptide

Chain M:  56% 44%




- Molecule 3: Cellular tumor antigen p53 peptide

Chain R:  67% 33%




- Molecule 4: 6-11 T cell receptor beta chain

Chain D:  71% 26% ..




- Molecule 4: 6-11 T cell receptor beta chain

Chain I:  72% 27% .



- Molecule 4: 6-11 T cell receptor beta chain


Chain N:  73% 25% .



- Molecule 4: 6-11 T cell receptor beta chain


Chain S:  69% 29% ..






- Molecule 5: 6-11 T cell receptor alpha chain

Chain E: 68% 25% ..




- Molecule 5: 6-11 T cell receptor alpha chain

Chain J: 76% 20% ..




- Molecule 5: 6-11 T cell receptor alpha chain

Chain O: 6% 67% 29% ..



- Molecule 5: 6-11 T cell receptor alpha chain

Chain T: 2% 70% 26% ..



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	127.23Å 55.14Å 304.88Å 90.00° 98.74° 90.00°	Depositor
Resolution (Å)	49.32 – 3.33 49.32 – 3.33	Depositor EDS
% Data completeness (in resolution range)	92.7 (49.32-3.33) 92.7 (49.32-3.33)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	4.00 (at 3.33Å)	Xtriage
Refinement program	PHENIX 1.16_3549	Depositor
R, R_{free}	0.237 , 0.295 0.237 , 0.293	Depositor DCC
R_{free} test set	2703 reflections (4.65%)	wwPDB-VP
Wilson B-factor (Å ²)	59.5	Xtriage
Anisotropy	0.087	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.27 , 27.9	EDS
L-test for twinning ²	$\langle L \rangle = 0.52$, $\langle L^2 \rangle = 0.35$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.91	EDS
Total number of atoms	26474	wwPDB-VP
Average B, all atoms (Å ²)	50.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 49.24 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 7.6773e-05. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.35	0/2305	0.57	2/3130 (0.1%)
1	F	0.26	0/2301	0.50	0/3125
1	K	0.47	4/2301 (0.2%)	0.54	2/3125 (0.1%)
1	P	0.25	0/2301	0.49	0/3125
2	B	0.39	0/847	0.56	0/1149
2	G	0.27	0/847	0.49	0/1149
2	L	0.32	0/847	0.53	0/1149
2	Q	0.32	0/843	0.56	0/1144
3	C	1.15	0/77	0.93	0/101
3	H	0.32	0/77	0.58	0/101
3	M	0.26	0/77	0.54	0/101
3	R	0.29	0/77	0.59	0/101
4	D	0.45	1/1966 (0.1%)	0.57	1/2678 (0.0%)
4	I	0.35	0/1964	0.54	0/2675
4	N	0.34	0/1964	0.58	1/2675 (0.0%)
4	S	0.31	0/1964	0.56	0/2675
5	E	0.50	0/1604	0.65	1/2181 (0.0%)
5	J	0.29	0/1604	0.55	0/2181
5	O	0.27	0/1604	0.59	1/2181 (0.0%)
5	T	0.37	0/1604	0.68	3/2181 (0.1%)
All	All	0.36	5/27174 (0.0%)	0.56	11/36927 (0.0%)

All (5) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	D	98	PRO	N-CD	-12.87	1.29	1.47
1	K	195	SER	N-CA	12.62	1.71	1.46
1	K	196	ASP	CA-CB	-9.98	1.31	1.53
1	K	196	ASP	CA-C	7.35	1.72	1.52
1	K	195	SER	C-N	6.37	1.48	1.34

The worst 5 of 11 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
5	T	30	PRO	N-CA-CB	-10.30	90.94	103.30
1	K	195	SER	N-CA-CB	-9.28	96.58	110.50
1	A	137	ASP	CB-CA-C	-8.60	93.19	110.40
1	A	49	ALA	N-CA-CB	6.60	119.33	110.10
5	T	30	PRO	N-CA-C	6.49	128.96	112.10

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2240	0	2083	50	0
1	F	2236	0	2081	35	0
1	K	2236	0	2081	35	0
1	P	2236	0	2080	34	0
2	B	824	0	775	19	0
2	G	824	0	775	15	0
2	L	824	0	775	7	0
2	Q	820	0	769	11	0
3	C	76	0	74	8	0
3	H	76	0	74	5	0
3	M	76	0	74	6	0
3	R	76	0	74	6	0
4	D	1915	0	1839	42	0
4	I	1913	0	1839	41	0
4	N	1913	0	1839	44	0
4	S	1913	0	1839	48	0
5	E	1569	0	1482	48	0
5	J	1569	0	1482	33	0
5	O	1569	0	1482	39	0
5	T	1569	0	1482	38	0
All	All	26474	0	24999	494	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 10.

The worst 5 of 494 close contacts within the same asymmetric unit are listed below, sorted by

their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:K:195:SER:CA	1:K:195:SER:N	1.71	1.49
1:K:146:LYS:NZ	4:N:97:ASP:OD1	1.71	1.19
4:S:98:PRO:HD3	5:T:97:HIS:CE1	1.77	1.16
5:E:186:ASN:HB2	5:E:189:ASN:ND2	1.66	1.11
1:A:131:ARG:HG2	1:A:157:ARG:HH21	1.20	1.02

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	273/275 (99%)	264 (97%)	8 (3%)	1 (0%)	34	68
1	F	273/275 (99%)	262 (96%)	11 (4%)	0	100	100
1	K	273/275 (99%)	263 (96%)	10 (4%)	0	100	100
1	P	273/275 (99%)	261 (96%)	12 (4%)	0	100	100
2	B	98/100 (98%)	90 (92%)	8 (8%)	0	100	100
2	G	98/100 (98%)	92 (94%)	5 (5%)	1 (1%)	15	49
2	L	98/100 (98%)	92 (94%)	5 (5%)	1 (1%)	15	49
2	Q	98/100 (98%)	91 (93%)	6 (6%)	1 (1%)	15	49
3	C	7/9 (78%)	7 (100%)	0	0	100	100
3	H	7/9 (78%)	6 (86%)	1 (14%)	0	100	100
3	M	7/9 (78%)	6 (86%)	1 (14%)	0	100	100
3	R	7/9 (78%)	6 (86%)	1 (14%)	0	100	100
4	D	241/246 (98%)	221 (92%)	18 (8%)	2 (1%)	19	53
4	I	241/246 (98%)	218 (90%)	21 (9%)	2 (1%)	19	53
4	N	241/246 (98%)	218 (90%)	19 (8%)	4 (2%)	9	37

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
4	S	241/246 (98%)	220 (91%)	18 (8%)	3 (1%)	13	45
5	E	198/206 (96%)	177 (89%)	19 (10%)	2 (1%)	15	49
5	J	198/206 (96%)	173 (87%)	23 (12%)	2 (1%)	15	49
5	O	198/206 (96%)	175 (88%)	21 (11%)	2 (1%)	15	49
5	T	198/206 (96%)	172 (87%)	23 (12%)	3 (2%)	10	40
All	All	3268/3344 (98%)	3014 (92%)	230 (7%)	24 (1%)	22	57

5 of 24 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
4	D	39	LEU
4	I	39	LEU
4	I	70	LEU
4	N	14	GLU
4	N	39	LEU

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	230/231 (100%)	223 (97%)	7 (3%)	41	70
1	F	229/231 (99%)	223 (97%)	6 (3%)	46	73
1	K	229/231 (99%)	221 (96%)	8 (4%)	36	66
1	P	229/231 (99%)	221 (96%)	8 (4%)	36	66
2	B	92/95 (97%)	91 (99%)	1 (1%)	73	86
2	G	92/95 (97%)	91 (99%)	1 (1%)	73	86
2	L	92/95 (97%)	91 (99%)	1 (1%)	73	86
2	Q	91/95 (96%)	88 (97%)	3 (3%)	38	68
3	C	9/9 (100%)	7 (78%)	2 (22%)	1	4
3	H	9/9 (100%)	9 (100%)	0	100	100
3	M	9/9 (100%)	9 (100%)	0	100	100

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
3	R	9/9 (100%)	9 (100%)	0	100	100
4	D	206/212 (97%)	194 (94%)	12 (6%)	20	53
4	I	206/212 (97%)	198 (96%)	8 (4%)	32	63
4	N	206/212 (97%)	199 (97%)	7 (3%)	37	67
4	S	206/212 (97%)	197 (96%)	9 (4%)	28	61
5	E	174/187 (93%)	167 (96%)	7 (4%)	31	63
5	J	174/187 (93%)	172 (99%)	2 (1%)	73	86
5	O	174/187 (93%)	171 (98%)	3 (2%)	60	80
5	T	174/187 (93%)	169 (97%)	5 (3%)	42	71
All	All	2840/2936 (97%)	2750 (97%)	90 (3%)	39	69

5 of 90 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
4	N	96	LEU
1	P	122	ASP
4	N	120	ASN
1	P	44	ARG
2	Q	99	ASP

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 13 such sidechains are listed below:

Mol	Chain	Res	Type
4	N	80	GLN
2	Q	18	ASN
5	T	97	HIS
4	S	120	ASN
4	S	204	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	275/275 (100%)	-0.37	0 100 100	21, 36, 57, 80	0
1	F	275/275 (100%)	-0.34	0 100 100	22, 37, 59, 72	0
1	K	275/275 (100%)	-0.11	0 100 100	30, 52, 76, 94	0
1	P	275/275 (100%)	-0.19	0 100 100	29, 52, 73, 84	0
2	B	100/100 (100%)	-0.27	0 100 100	20, 35, 52, 73	0
2	G	100/100 (100%)	-0.33	0 100 100	20, 35, 54, 70	0
2	L	100/100 (100%)	-0.18	0 100 100	30, 50, 70, 86	0
2	Q	100/100 (100%)	-0.01	1 (1%) 82 83	30, 60, 81, 94	0
3	C	9/9 (100%)	-0.42	0 100 100	19, 23, 28, 28	0
3	H	9/9 (100%)	-0.34	0 100 100	22, 28, 34, 37	0
3	M	9/9 (100%)	0.12	0 100 100	35, 51, 55, 70	0
3	R	9/9 (100%)	0.08	0 100 100	34, 44, 47, 57	0
4	D	243/246 (98%)	-0.25	0 100 100	18, 36, 77, 97	3 (1%)
4	I	243/246 (98%)	-0.21	0 100 100	22, 36, 66, 96	3 (1%)
4	N	243/246 (98%)	0.18	3 (1%) 79 79	36, 68, 105, 113	3 (1%)
4	S	243/246 (98%)	-0.06	1 (0%) 92 94	30, 58, 94, 115	3 (1%)
5	E	200/206 (97%)	-0.26	0 100 100	16, 38, 64, 88	0
5	J	200/206 (97%)	-0.27	0 100 100	17, 39, 64, 91	0
5	O	200/206 (97%)	0.54	12 (6%) 21 23	44, 82, 108, 118	0
5	T	200/206 (97%)	0.20	4 (2%) 65 64	35, 70, 99, 130	0
All	All	3308/3344 (98%)	-0.12	21 (0%) 89 90	16, 47, 89, 130	12 (0%)

The worst 5 of 21 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
5	O	135	LEU	3.9
5	T	202	PRO	3.3
5	O	192	ILE	3.0
5	O	165	SER	2.9
4	S	116	GLU	2.9

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.