



Full wwPDB X-ray Structure Validation Report ⓘ

May 15, 2020 – 05:33 pm BST

PDB ID : 3ROO
Title : Murine class I major histocompatibility complex H-2Kb in complex with immunodominant LCMV-derived gp34-41 peptide
Authors : Madhurantakam, C.; Duru, A.D.; Leong, C.; Sandalova, T.; Webb, J.R.; Achour, A.
Deposited on : 2011-04-26
Resolution : 2.00 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix)	:	1.13
EDS	:	2.11
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac	:	5.8.0158
CCP4	:	7.0.044 (Gargrove)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.11


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	8085 (2.00-2.00)
Clashscore	141614	9178 (2.00-2.00)
Ramachandran outliers	138981	9054 (2.00-2.00)
Sidechain outliers	138945	9053 (2.00-2.00)
RSRZ outliers	127900	7900 (2.00-2.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	275	<div> <div>11%</div> <div>95%</div> <div>..</div> </div>
1	C	275	<div> <div>16%</div> <div>93%</div> <div>6% .</div> </div>
2	B	99	<div> <div>2%</div> <div>93%</div> <div>7%</div> </div>
2	D	99	<div> <div>8%</div> <div>89%</div> <div>10% .</div> </div>
3	E	8	<div> <div>100%</div> </div>
3	F	8	<div> <div>100%</div> </div>

2 Entry composition

There are 7 unique types of molecules in this entry. The entry contains 6694 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Murine class I major histocompatibility complex H-2Kb.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	275	Total	C	N	O	S	0	4	0
			2258	1425	397	427	9			
1	C	271	Total	C	N	O	S	0	5	0
			2222	1404	388	420	10			

- Molecule 2 is a protein called Beta-2-microglobulin.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	99	Total	C	N	O	S	0	3	0
			830	529	138	155	8			
2	D	99	Total	C	N	O	S	0	3	0
			824	526	135	155	8			

- Molecule 3 is a protein called Pre-glycoprotein polyprotein GP complex.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	E	8	Total	C	N	O	S	0	0	0
			64	42	9	12	1			
3	F	8	Total	C	N	O	S	0	0	0
			64	42	9	12	1			

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
E	8	MET	CYS	ENGINEERED MUTATION	UNP Q9QDK7
F	8	MET	CYS	ENGINEERED MUTATION	UNP Q9QDK7

- Molecule 4 is GLYCEROL (three-letter code: GOL) (formula: C₃H₈O₃).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	C	O	0	0
			6	3	3		
4	D	1	Total	C	O	0	0
			6	3	3		

- Molecule 5 is (4S)-2-METHYL-2,4-PENTANEDIOL (three-letter code: MPD) (formula: $C_6H_{14}O_2$).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	B	1	Total	C	O	0	0
			8	6	2		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	C	1	Total	C	O	0	0
			8	6	2		

- Molecule 6 is PHOSPHATE ION (three-letter code: PO4) (formula: O₄P).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
6	B	1	Total	O	P	0	0
			5	4	1		
6	C	1	Total	O	P	0	0
			5	4	1		


- Molecule 7 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
7	A	146	Total	O	0	0
			146	146		
7	B	74	Total	O	0	0
			74	74		
7	C	126	Total	O	0	0
			126	126		
7	D	34	Total	O	0	0
			34	34		
7	E	7	Total	O	0	0
			7	7		
7	F	7	Total	O	0	0
			7	7		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Murine class I major histocompatibility complex H-2Kb




- Molecule 1: Murine class I major histocompatibility complex H-2Kb



- Molecule 2: Beta-2-microglobulin



- Molecule 2: Beta-2-microglobulin



- Molecule 3: Pre-glycoprotein polyprotein GP complex

Chain E:  100%

There are no outlier residues recorded for this chain.

- Molecule 3: Pre-glycoprotein polyprotein GP complex

Chain F:  100%

There are no outlier residues recorded for this chain.

4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	88.45Å 92.59Å 128.83Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	57.26 – 2.00 57.28 – 2.00	Depositor EDS
% Data completeness (in resolution range)	99.9 (57.26-2.00) 99.9 (57.28-2.00)	Depositor EDS
R_{merge}	0.08	Depositor
R_{sym}	0.08	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.23 (at 2.00Å)	Xtriage
Refinement program	REFMAC 5.5.0109	Depositor
R, R_{free}	0.225 , 0.267 0.230 , 0.271	Depositor DCC
R_{free} test set	3633 reflections (5.04%)	wwPDB-VP
Wilson B-factor (Å ²)	30.7	Xtriage
Anisotropy	0.112	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.38 , 46.7	EDS
L-test for twinning ²	$\langle L \rangle = 0.51$, $\langle L^2 \rangle = 0.34$	Xtriage
Estimated twinning fraction	0.000 for k,h,-l	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	6694	wwPDB-VP
Average B, all atoms (Å ²)	41.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 19.60% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: GOL, PO4, MPD

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.45	0/2336	0.60	0/3171
1	C	0.44	0/2302	0.58	0/3121
2	B	0.43	0/870	0.57	0/1178
2	D	0.43	0/864	0.56	0/1171
3	E	0.45	0/65	0.58	0/86
3	F	0.58	0/65	0.55	0/86
All	All	0.45	0/6502	0.58	0/8813

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2258	0	2146	9	0
1	C	2222	0	2113	9	0
2	B	830	0	800	6	0
2	D	824	0	789	10	0
3	E	64	0	61	0	0
3	F	64	0	61	0	0
4	A	6	0	8	1	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
4	D	6	0	8	0	0
5	B	8	0	14	5	0
5	C	8	0	14	5	0
6	B	5	0	0	0	0
6	C	5	0	0	0	0
7	A	146	0	0	1	0
7	B	74	0	0	0	0
7	C	126	0	0	0	0
7	D	34	0	0	1	0
7	E	7	0	0	0	0
7	F	7	0	0	0	0
All	All	6694	0	6014	33	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (33) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
5:C:5276:MPD:H11	2:D:33:PRO:O	1.83	0.79
2:D:12:ARG:CB	2:D:22:ILE:HG23	2.12	0.78
2:B:54[B]:MET:CE	5:B:5276:MPD:HM2	2.24	0.66
2:B:54[B]:MET:HE2	5:B:5276:MPD:H11	1.83	0.61
1:C:23[B]:MET:SD	2:D:54[B]:MET:SD	2.99	0.60
1:C:117:ALA:HB2	2:D:60:TRP:CE2	2.38	0.59
1:A:23:MET:HE3	5:B:5276:MPD:H53	1.85	0.58
1:C:94[B]:THR:HG21	2:D:33:PRO:CD	2.34	0.58
2:B:54[B]:MET:HE2	5:B:5276:MPD:HM2	1.85	0.57
2:D:31:HIS:ND1	7:D:338:HOH:O	2.33	0.56
1:C:181:ARG:NH2	1:C:239:GLY:O	2.35	0.56
5:C:5276:MPD:H12	5:C:5276:MPD:O4	2.06	0.56
1:A:215:LEU:HD22	1:A:261:VAL:HG22	1.88	0.54
1:A:117:ALA:HB2	2:B:60:TRP:CE2	2.43	0.54
2:B:27:VAL:HG21	2:B:37:ILE:HD13	1.89	0.53
5:C:5276:MPD:H4	2:D:33:PRO:HB2	1.93	0.49
2:D:1:ILE:HG23	2:D:1:ILE:O	2.13	0.48
2:D:22:ILE:HD11	2:D:67:HIS:HB2	1.98	0.46
1:A:182:THR:HG21	1:A:265:GLY:HA2	1.97	0.46
1:C:216:THR:OG1	1:C:260:HIS:HB2	2.18	0.44
1:C:251:LEU:HD12	1:C:252:GLY:H	1.82	0.44
2:D:21:ASN:OD1	2:D:22:ILE:N	2.45	0.44

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:182:THR:HG23	1:A:265:GLY:HA3	2.00	0.43
1:C:23[A]:MET:SD	5:C:5276:MPD:H53	2.59	0.43
2:B:54[B]:MET:HE1	5:B:5276:MPD:HM2	2.01	0.42
1:C:251:LEU:HD12	1:C:252:GLY:N	2.35	0.42
1:A:228:MET:HG2	1:A:247:VAL:HG12	2.02	0.42
1:A:215:LEU:CD2	1:A:261:VAL:HG22	2.50	0.41
1:A:54:GLN:NE2	7:A:289:HOH:O	2.53	0.41
1:C:203:CYS:HB2	1:C:217:TRP:CZ2	2.56	0.41
5:C:5276:MPD:C1	5:C:5276:MPD:O4	2.69	0.40
1:A:141:LEU:HD12	1:A:141:LEU:HA	2.00	0.40

There are no symmetry-related clashes.

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	277/275 (101%)	268 (97%)	9 (3%)	0	100	100
1	C	270/275 (98%)	263 (97%)	7 (3%)	0	100	100
2	B	100/99 (101%)	98 (98%)	2 (2%)	0	100	100
2	D	100/99 (101%)	100 (100%)	0	0	100	100
3	E	6/8 (75%)	6 (100%)	0	0	100	100
3	F	6/8 (75%)	6 (100%)	0	0	100	100
All	All	759/764 (99%)	741 (98%)	18 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	237/233 (102%)	232 (98%)	5 (2%)	53	57
1	C	234/233 (100%)	229 (98%)	5 (2%)	53	57
2	B	97/94 (103%)	94 (97%)	3 (3%)	40	40
2	D	96/94 (102%)	93 (97%)	3 (3%)	40	40
3	E	6/6 (100%)	6 (100%)	0	100	100
3	F	6/6 (100%)	6 (100%)	0	100	100
All	All	676/666 (102%)	660 (98%)	16 (2%)	49	51

All (16) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	45	TYR
1	A	114	GLN
1	A	182	THR
1	A	196	GLU
1	A	228	MET
2	B	56	PHE
2	B	58	LYS
2	B	70	PHE
1	C	45	TYR
1	C	114	GLN
1	C	141	LEU
1	C	230	LEU
1	C	255	GLN
2	D	22	ILE
2	D	56	PHE
2	D	58	LYS

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (13) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	87	GLN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	114	GLN
1	A	115	GLN
1	A	174	ASN
1	A	218	GLN
2	B	38	GLN
1	C	42	ASN
1	C	87	GLN
1	C	114	GLN
1	C	115	GLN
1	C	174	ASN
2	D	29	GLN
2	D	38	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

6 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
5	MPD	B	5276	-	7,7,7	0.23	0	9,10,10	0.67	0
4	GOL	D	3968	-	5,5,5	0.43	0	5,5,5	0.40	0

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
6	PO4	C	276	-	4,4,4	0.97	0	6,6,6	0.43	0
4	GOL	A	3968	-	5,5,5	0.32	0	5,5,5	0.56	0
6	PO4	B	100	-	4,4,4	0.89	0	6,6,6	0.37	0
5	MPD	C	5276	-	7,7,7	0.46	0	9,10,10	0.61	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	GOL	A	3968	-	-	4/4/4/4	-
5	MPD	B	5276	-	-	4/5/5/5	-
5	MPD	C	5276	-	-	3/5/5/5	-
4	GOL	D	3968	-	-	2/4/4/4	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (13) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	D	3968	GOL	C1-C2-C3-O3
4	A	3968	GOL	O1-C1-C2-C3
5	C	5276	MPD	C1-C2-C3-C4
5	C	5276	MPD	O2-C2-C3-C4
4	A	3968	GOL	C1-C2-C3-O3
4	D	3968	GOL	O2-C2-C3-O3
4	A	3968	GOL	O2-C2-C3-O3
4	A	3968	GOL	O1-C1-C2-O2
5	B	5276	MPD	O2-C2-C3-C4
5	B	5276	MPD	C1-C2-C3-C4
5	B	5276	MPD	CM-C2-C3-C4
5	B	5276	MPD	C2-C3-C4-O4
5	C	5276	MPD	C2-C3-C4-O4

There are no ring outliers.

3 monomers are involved in 11 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
5	B	5276	MPD	5	0
4	A	3968	GOL	1	0
5	C	5276	MPD	5	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	275/275 (100%)	0.60	29 (10%) 6 5	24, 37, 64, 69	1 (0%)
1	C	271/275 (98%)	0.74	43 (15%) 1 1	25, 40, 67, 75	1 (0%)
2	B	99/99 (100%)	0.26	2 (2%) 65 63	26, 34, 48, 54	0
2	D	99/99 (100%)	0.64	8 (8%) 12 11	29, 43, 59, 62	0
3	E	8/8 (100%)	0.22	0 100 100	25, 28, 33, 33	0
3	F	8/8 (100%)	0.10	0 100 100	27, 29, 32, 33	0
All	All	760/764 (99%)	0.60	82 (10%) 5 5	24, 38, 64, 75	2 (0%)

All (82) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	C	256	TYR	11.3
1	A	256	TYR	7.5
1	A	220	ASN	6.5
1	C	274	TRP	5.9
1	A	226	GLN	5.7
1	C	250	PRO	5.5
1	C	251	LEU	5.4
1	A	219	LEU	5.2
1	C	273	ARG	5.0
1	C	196	GLU	4.9
1	C	253	LYS	4.9
1	C	252	GLY	4.9
1	C	199	VAL	4.7
1	C	222	GLU	4.6
1	C	248	VAL	4.2
1	C	221	GLY	4.1
2	D	22	ILE	4.0
1	C	190	THR	4.0
1	A	274	TRP	3.8

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	225	ILE	3.8
1	C	224	LEU	3.8
1	C	257	TYR	3.6
1	C	254	GLU	3.6
1	C	258	THR	3.6
1	A	273	ARG	3.5
1	A	198	LYS	3.4
1	C	275	GLU	3.4
1	C	201	LEU	3.3
1	A	227	ASP	3.3
1	A	224	LEU	3.3
2	D	19	LYS	3.2
1	A	272	LEU	3.2
1	C	255	GLN	3.2
1	A	222	GLU	3.2
1	C	219	LEU	3.2
1	C	198	LYS	3.1
1	C	195	PRO	3.1
1	C	217	TRP	3.1
1	A	196	GLU	3.1
1	C	191	HIS	3.1
2	B	48	LYS	3.1
1	C	249	VAL	3.0
1	A	41	GLU	3.0
2	D	14	PRO	3.0
1	A	262	TYR	3.0
1	A	199	VAL	2.9
1	C	189	VAL	2.9
1	C	192	HIS	2.9
1	C	268	GLU	2.8
1	A	253	LYS	2.8
1	C	218	GLN	2.7
1	C	200	THR	2.7
1	C	271	THR	2.7
1	C	228	MET	2.7
1	A	195	PRO	2.6
2	D	20	PRO	2.6
1	A	254	GLU	2.6
1	A	257	TYR	2.5
1	A	260	HIS	2.5
2	D	71	THR	2.4
1	C	267	PRO	2.4

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	C	194	ARG	2.4
1	C	272	LEU	2.4
1	C	41	GLU	2.3
1	C	193	SER	2.3
1	A	191	HIS	2.2
1	A	255	GLN	2.2
1	A	249	VAL	2.2
1	A	221	GLY	2.2
1	A	197	ASP	2.2
1	C	259	CYS	2.2
2	D	1	ILE	2.1
2	D	16	GLU	2.1
2	B	88	ALA	2.1
1	C	197	ASP	2.1
1	C	188	HIS	2.1
2	D	18	GLY	2.1
1	A	259	CYS	2.1
1	C	230	LEU	2.0
1	C	262	TYR	2.0
1	A	89	LYS	2.0
1	A	247	VAL	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
4	GOL	D	3968	6/6	0.82	0.19	58,60,60,61	0
4	GOL	A	3968	6/6	0.86	0.19	29,38,40,40	0

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
6	PO4	C	276	5/5	0.91	0.19	65,66,66,66	0
5	MPD	C	5276	8/8	0.91	0.19	34,37,39,41	0
6	PO4	B	100	5/5	0.92	0.18	71,72,73,73	0
5	MPD	B	5276	8/8	0.95	0.14	33,35,38,40	0

6.5 Other polymers [i](#)

There are no such residues in this entry.