



# Full wwPDB X-ray Structure Validation Report ⓘ

May 15, 2020 – 12:59 pm BST

PDB ID : 3RQJ  
Title : Structure of the neuronal nitric oxide synthase heme domain in complex with 6-(((3R,4R)-4-(2-((1S,2R)-2-(3-Fluorophenyl)cyclopropylamino)ethoxy)pyrrolidin-3-yl)methyl)-4-methylpyridin-2-amine  
Authors : Li, H.; Delker, S.L.; Poulos, T.L.  
Deposited on : 2011-04-28  
Resolution : 1.84 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.11  
buster-report : 1.1.7 (2018)  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.11


# 1 Overall quality at a glance ⓘ

The following experimental techniques were used to determine the structure:

## *X-RAY DIFFRACTION*

The reported resolution of this entry is 1.84 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	4003 (1.86-1.82)
Clashscore	141614	4233 (1.86-1.82)
Ramachandran outliers	138981	4185 (1.86-1.82)
Sidechain outliers	138945	4186 (1.86-1.82)
RSRZ outliers	127900	3957 (1.86-1.82)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	422	<div> <div>18%</div> <div>86%</div> <div>10%</div> <div>••</div> </div>
1	B	422	<div> <div>8%</div> <div>91%</div> <div>7%</div> <div>•</div> </div>

## 2 Entry composition [i](#)

There are 7 unique types of molecules in this entry. The entry contains 7189 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Nitric oxide synthase, brain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	407	Total	C	N	O	S	0	2	0
			3322	2127	566	608	21			
1	B	411	Total	C	N	O	S	0	2	0
			3354	2146	574	612	22			

- Molecule 2 is PROTOPORPHYRIN IX CONTAINING FE (three-letter code: HEM) (formula:  $C_{34}H_{32}FeN_4O_4$ ).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total	C	Fe	N	O	0	0
			43	34	1	4	4		
2	B	1	Total	C	Fe	N	O	0	0
			43	34	1	4	4		

- Molecule 3 is 5,6,7,8-TETRAHYDROBIOPTERIN (three-letter code: H4B) (formula:  $C_9H_{15}N_5O_3$ ).




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
3	A	1	Total	C	N	O	0	0
			17	9	5	3		
3	B	1	Total	C	N	O	0	0
			17	9	5	3		

- Molecule 4 is ACETATE ION (three-letter code: ACT) (formula:  $C_2H_3O_2$ ).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	C	O	0	0
			4	2	2		
4	B	1	Total	C	O	0	0
			4	2	2		

- Molecule 5 is 6-([(3R,4R)-4-(2-([(1S,2R)-2-(3-fluorophenyl)cyclopropyl]amino)ethoxy)pyrrolidin-3-yl)methyl]-4-methylpyridin-2-amine (three-letter code: Y2B) (formula: C<sub>22</sub>H<sub>29</sub>FN<sub>4</sub>O).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
5	A	1	Total 28	C 22	F 1	N 4	O 1	0	0
5	B	1	Total 28	C 22	F 1	N 4	O 1	0	0

- Molecule 6 is ZINC ION (three-letter code: ZN) (formula:  $\text{Zn}$ ).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
6	A	1	Total Zn 1 1	0	0


- Molecule 7 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
7	A	138	Total O 138 138	0	0
7	B	190	Total O 190 190	0	0


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Nitric oxide synthase, brain



- Molecule 1: Nitric oxide synthase, brain



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	51.84Å 110.74Å 164.30Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	38.11 – 1.84 37.84 – 1.84	Depositor EDS
% Data completeness (in resolution range)	98.3 (38.11-1.84) 98.3 (37.84-1.84)	Depositor EDS
$R_{merge}$	0.04	Depositor
$R_{sym}$	0.04	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.46 (at 1.84Å)	Xtriage
Refinement program	REFMAC	Depositor
R, $R_{free}$	0.189 , 0.221 0.208 , 0.234	Depositor DCC
$R_{free}$ test set	4059 reflections (4.94%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	36.5	Xtriage
Anisotropy	0.302	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.36 , 39.6	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.96	EDS
Total number of atoms	7189	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	49.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.33% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: Y2B, ZN, H4B, HEM, ACT

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.69	0/3421	0.69	1/4641 (0.0%)
1	B	0.77	0/3453	0.73	0/4681
All	All	0.73	0/6874	0.71	1/9322 (0.0%)

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	410	ARG	NE-CZ-NH2	5.17	122.88	120.30

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3322	0	3232	31	0
1	B	3354	0	3270	21	0
2	A	43	0	30	2	0
2	B	43	0	30	3	0
3	A	17	0	15	0	0
3	B	17	0	15	0	0
4	A	4	0	3	0	0
4	B	4	0	3	0	0

*Continued on next page...*



*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
5	A	28	0	29	6	0
5	B	28	0	29	4	0
6	A	1	0	0	0	0
7	A	138	0	0	0	0
7	B	190	0	0	3	0
All	All	7189	0	6656	52	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (52) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:592[A]:GLU:OE1	5:A:800:Y2B:H26	1.72	0.89
1:B:592[A]:GLU:OE1	5:B:800:Y2B:H26	1.81	0.79
2:A:750:HEM:HMC2	2:A:750:HEM:HBC2	1.70	0.73
2:B:750:HEM:HHC	2:B:750:HEM:HBB2	1.71	0.72
1:A:371:ARG:HG3	1:A:371:ARG:HH21	1.55	0.72
1:A:307:GLU:HG3	7:B:1003:HOH:O	1.94	0.68
1:A:371:ARG:CG	1:A:371:ARG:HH21	2.09	0.65
1:A:380:ARG:HD3	1:A:400:GLU:OE1	2.01	0.60
1:A:592[A]:GLU:OE1	5:A:800:Y2B:H14	2.03	0.58
1:B:350:THR:HG22	1:B:352:ASP:H	1.70	0.57
1:A:371:ARG:CG	1:A:371:ARG:NH2	2.66	0.56
1:A:455:LEU:HD12	1:A:587:TRP:HB3	1.88	0.55
1:B:659:ILE:O	1:B:663:GLU:HG3	2.07	0.55
1:A:332:MET:HE1	1:B:301:LEU:HD22	1.92	0.52
1:A:592[A]:GLU:CD	5:A:800:Y2B:H26	2.28	0.52
1:A:645:LYS:HB2	1:A:645:LYS:HZ1	1.75	0.52
1:A:332:MET:CE	1:B:301:LEU:HD22	2.40	0.51
2:B:750:HEM:HBC2	2:B:750:HEM:CMC	2.41	0.51
1:A:330:ILE:HD11	1:B:696:LEU:HB3	1.95	0.49
1:B:499:VAL:O	1:B:503:GLU:HG3	2.13	0.49
1:A:569:ASN:H	1:A:569:ASN:HD22	1.58	0.48
1:B:614:MET:CE	1:B:632:GLU:HG3	2.44	0.48
1:A:478:GLN:HB2	1:A:481:ARG:HG3	1.96	0.48
2:A:750:HEM:HBA1	5:A:800:Y2B:H14	1.97	0.47
1:A:694:GLU:HB3	1:B:335:ILE:HD13	1.97	0.47
1:A:480:ILE:HD13	1:A:541:VAL:HG13	1.98	0.46
1:A:645:LYS:NZ	1:A:645:LYS:HB2	2.29	0.45
1:A:455:LEU:HD12	1:A:587:TRP:CB	2.46	0.45

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:592[A]:GLU:CD	5:B:800:Y2B:H26	2.38	0.45
1:B:678:TRP:CH2	5:B:800:Y2B:H2'A	2.52	0.45
1:B:605:ASN:ND2	7:B:1019:HOH:O	2.39	0.44
1:A:686:SER:HA	1:A:691:PHE:CG	2.52	0.44
1:B:535:GLN:HG3	7:B:1188:HOH:O	2.18	0.44
1:B:355:PHE:N	1:B:356:PRO:HD2	2.33	0.43
1:B:510:TRP:CE2	1:B:521:PRO:HD3	2.53	0.43
2:B:750:HEM:HBC2	2:B:750:HEM:HMC1	2.01	0.43
1:B:351:LYS:H	1:B:351:LYS:HG3	1.69	0.43
1:A:709:ASP:HA	1:A:710:PRO:HD3	1.85	0.42
1:B:353:GLN:HB3	1:B:353:GLN:HE21	1.66	0.42
1:A:663:GLU:O	1:A:667:ARG:HG2	2.19	0.42
1:A:592[B]:GLU:OE2	5:A:800:Y2B:N12	2.49	0.42
1:A:569:ASN:O	1:A:707:GLN:HG2	2.20	0.42
1:B:403:TYR:CE1	1:B:407:HIS:CE1	3.07	0.42
1:A:409:TRP:CE3	1:A:421:TRP:HA	2.55	0.42
1:B:478:GLN:HB2	1:B:481:ARG:HG3	2.01	0.42
1:A:350:THR:N	1:A:353:GLN:HE21	2.18	0.41
1:A:572:LEU:HB3	1:A:579:PHE:HB2	2.03	0.41
1:A:628:GLN:HG2	1:B:631:VAL:HG11	2.03	0.41
1:A:676:TRP:CE2	1:A:680:VAL:HG21	2.56	0.41
1:A:352:ASP:N	1:A:352:ASP:OD2	2.54	0.41
1:A:567:VAL:CG2	5:A:800:Y2B:H15A	2.50	0.41
1:B:592[A]:GLU:OE1	5:B:800:Y2B:H14	2.22	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	405/422 (96%)	393 (97%)	12 (3%)	0	<a href="#">100</a> <a href="#">100</a>

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	B	409/422 (97%)	403 (98%)	6 (2%)	0	100	100
All	All	814/844 (96%)	796 (98%)	18 (2%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	365/377 (97%)	350 (96%)	15 (4%)	30	13
1	B	368/377 (98%)	364 (99%)	4 (1%)	73	64
All	All	733/754 (97%)	714 (97%)	19 (3%)	46	29

All (19) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	320	SER
1	A	321	THR
1	A	350	THR
1	A	352	ASP
1	A	370	LYS
1	A	390	SER
1	A	507	GLN
1	A	514	ARG
1	A	527	ASN
1	A	547	ARG
1	A	555	LYS
1	A	569	ASN
1	A	620	LYS
1	A	645	LYS
1	A	715	VAL
1	B	348	VAL
1	B	353	GLN
1	B	390	SER

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	B	547	ARG

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (12) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	454	ASN
1	A	527	ASN
1	A	569	ASN
1	A	605	ASN
1	A	697	ASN
1	B	425	GLN
1	B	440	ASN
1	B	454	ASN
1	B	507	GLN
1	B	601	ASN
1	B	605	ASN
1	B	697	ASN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 5.6 Ligand geometry [i](#)

Of 9 ligands modelled in this entry, 1 is monoatomic - leaving 8 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the

expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
2	HEM	B	750	1	27,50,50	2.09	10 (37%)	17,82,82	2.45	3 (17%)
5	Y2B	B	800	-	29,31,31	0.99	1 (3%)	33,43,43	2.54	10 (30%)
3	H4B	A	760	-	16,18,18	0.82	0	11,26,26	2.82	5 (45%)
4	ACT	A	860	-	1,3,3	1.43	0	0,3,3	0.00	-
4	ACT	B	860	-	1,3,3	1.91	0	0,3,3	0.00	-
2	HEM	A	750	1	27,50,50	2.18	6 (22%)	17,82,82	2.32	5 (29%)
5	Y2B	A	800	-	29,31,31	1.01	1 (3%)	33,43,43	2.55	10 (30%)
3	H4B	B	760	-	16,18,18	1.28	1 (6%)	11,26,26	2.57	5 (45%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	HEM	B	750	1	-	0/6/54/54	-
5	Y2B	B	800	-	-	2/13/30/30	0/4/4/4
3	H4B	A	760	-	-	0/8/17/17	0/2/2/2
2	HEM	A	750	1	-	0/6/54/54	-
5	Y2B	A	800	-	-	2/13/30/30	0/4/4/4
3	H4B	B	760	-	-	0/8/17/17	0/2/2/2

All (19) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	750	HEM	C3B-C2B	-5.31	1.33	1.40
2	A	750	HEM	C3D-C2D	5.12	1.52	1.37
2	B	750	HEM	C3D-C2D	4.36	1.50	1.37
2	B	750	HEM	C3C-C2C	-4.16	1.34	1.40
2	B	750	HEM	C3B-C2B	-3.89	1.35	1.40
2	B	750	HEM	CAD-C3D	3.63	1.58	1.52
2	A	750	HEM	C3C-CAC	3.54	1.55	1.47
5	B	800	Y2B	C15-C13	3.12	1.53	1.49
2	A	750	HEM	C3B-CAB	3.02	1.54	1.47
2	A	750	HEM	C3C-C2C	-2.79	1.36	1.40
5	A	800	Y2B	C15-C13	2.76	1.53	1.49
3	B	760	H4B	C7-N8	2.74	1.49	1.44
2	B	750	HEM	C3C-CAC	2.67	1.53	1.47

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	750	HEM	CAA-C2A	2.61	1.55	1.52
2	A	750	HEM	CMA-C3A	2.58	1.57	1.51
2	B	750	HEM	C3B-CAB	2.55	1.53	1.47
2	B	750	HEM	CMA-C3A	2.33	1.56	1.51
2	B	750	HEM	C4A-NA	2.16	1.40	1.36
2	B	750	HEM	C4B-NB	2.13	1.40	1.36

All (38) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
5	B	800	Y2B	C15-C13-N12	-8.00	101.90	117.50
5	A	800	Y2B	C15-C13-N12	-7.43	103.02	117.50
2	B	750	HEM	CBA-CAA-C2A	-7.04	99.51	112.49
5	B	800	Y2B	C15-C14-C21	-6.50	109.73	122.24
5	A	800	Y2B	C15-C14-C21	-6.41	109.91	122.24
2	A	750	HEM	CBD-CAD-C3D	-6.23	100.99	112.48
5	A	800	Y2B	C02-N01-C06	6.08	122.71	118.10
5	B	800	Y2B	C02-N01-C06	5.52	122.28	118.10
3	B	760	H4B	C4-C4A-C8A	5.43	119.39	114.57
3	A	760	H4B	C4-C4A-N5	5.40	123.65	119.12
2	B	750	HEM	CBD-CAD-C3D	-5.35	102.63	112.48
2	A	750	HEM	CBA-CAA-C2A	-4.64	103.92	112.49
3	A	760	H4B	C4-C4A-C8A	4.37	118.45	114.57
5	A	800	Y2B	C26-C25-C24	-3.77	114.90	120.25
3	B	760	H4B	C4-N3-C2	3.55	121.57	115.93
3	B	760	H4B	N3-C2-N1	-3.30	120.24	125.42
3	A	760	H4B	N3-C2-N1	-3.26	120.31	125.42
3	A	760	H4B	C2-N1-C8A	3.23	121.79	114.54
5	A	800	Y2B	C5'-N1'-C2'	3.05	112.63	105.42
5	A	800	Y2B	C05-C06-N01	-2.91	119.82	122.90
5	B	800	Y2B	C11-N12-C13	2.90	118.30	114.20
3	B	760	H4B	N2-C2-N1	2.76	121.54	117.25
5	B	800	Y2B	N02-C02-N01	2.75	120.84	116.49
3	A	760	H4B	C4-N3-C2	2.73	120.27	115.93
2	A	750	HEM	CMC-C2C-C3C	2.73	129.79	124.68
5	B	800	Y2B	C04-C05-C06	-2.47	118.70	120.32
5	B	800	Y2B	C5'-N1'-C2'	2.42	111.14	105.42
5	A	800	Y2B	C25-C24-C23	2.41	123.11	118.23
2	A	750	HEM	C4C-C3C-C2C	2.34	108.53	106.90
3	B	760	H4B	C2-N1-C8A	2.27	119.62	114.54
5	B	800	Y2B	C05-C06-N01	-2.20	120.57	122.90
5	A	800	Y2B	C26-C21-C22	2.19	121.28	118.76

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
5	B	800	Y2B	C24-C23-C22	-2.14	120.51	123.29
2	A	750	HEM	C3C-C4C-NC	-2.11	106.95	110.94
2	B	750	HEM	CMA-C3A-C4A	-2.10	125.24	128.46
5	B	800	Y2B	C10-C11-N12	2.09	116.77	111.54
5	A	800	Y2B	C24-C23-C22	-2.07	120.60	123.29
5	A	800	Y2B	N02-C02-N01	2.00	119.65	116.49

There are no chirality outliers.

All (4) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
5	A	800	Y2B	N01-C06-C08-C3'
5	A	800	Y2B	O09-C10-C11-N12
5	B	800	Y2B	N01-C06-C08-C3'
5	B	800	Y2B	O09-C10-C11-N12

There are no ring outliers.


4 monomers are involved in 14 short contacts:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	B	750	HEM	3	0
5	B	800	Y2B	4	0
2	A	750	HEM	2	0
5	A	800	Y2B	6	0


The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.











## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	407/422 (96%)	0.97	78 (19%) <b>1</b> <b>1</b>	28, 53, 92, 121	0
1	B	411/422 (97%)	0.47	34 (8%) <b>11</b> <b>10</b>	28, 41, 66, 80	0
All	All	818/844 (96%)	0.72	112 (13%) <b>3</b> <b>2</b>	28, 47, 86, 121	0

All (112) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	300	PHE	8.6
1	A	488	PRO	8.2
1	A	716	TRP	8.1
1	A	351	LYS	5.9
1	A	713	THR	5.9
1	A	355	PHE	5.8
1	B	619	ARG	5.5
1	A	352	ASP	5.5
1	A	486	LYS	5.3
1	A	388	ILE	4.7
1	A	506	ILE	4.5
1	A	300	PHE	4.3
1	B	348	VAL	4.3
1	A	715	VAL	4.2
1	B	350	THR	4.1
1	A	514	ARG	4.0
1	A	491	SER	3.7
1	A	714	HIS	3.7
1	A	487	GLN	3.7
1	A	507	GLN	3.7
1	A	567	VAL	3.7
1	A	678	TRP	3.6
1	B	667	ARG	3.6
1	A	385	ASN	3.5

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	A	389	GLU	3.5
1	B	616	LEU	3.5
1	A	386	LYS	3.4
1	A	392	SER	3.4
1	A	490	GLY	3.4
1	A	592[A]	GLU	3.3
1	A	551	PHE	3.3
1	A	479	LEU	3.3
1	A	390	SER	3.3
1	A	552	ASP	3.2
1	A	503	GLU	3.2
1	A	593	ILE	3.2
1	A	469	LYS	3.2
1	A	299	ARG	3.1
1	B	620	LYS	3.1
1	A	619	ARG	3.1
1	B	677	VAL	3.1
1	B	567	VAL	3.1
1	A	511	LYS	3.1
1	A	391	THR	3.0
1	B	680	VAL	3.0
1	A	480	ILE	3.0
1	B	691	PHE	2.9
1	A	680	VAL	2.9
1	B	479	LEU	2.9
1	A	584	PHE	2.9
1	A	591	THR	2.8
1	A	489	ASP	2.8
1	B	561	TRP	2.8
1	B	310	VAL	2.8
1	A	382	GLU	2.8
1	A	321	THR	2.8
1	A	350	THR	2.8
1	A	682	PRO	2.8
1	B	566	ALA	2.8
1	A	676	TRP	2.8
1	B	592[A]	GLU	2.8
1	B	302	LYS	2.7
1	A	561	TRP	2.7
1	B	591	THR	2.7
1	A	416	VAL	2.7
1	A	415	CYS	2.7

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	A	492	THR	2.7
1	B	713	THR	2.7
1	A	677	VAL	2.6
1	B	715	VAL	2.6
1	A	588	TYR	2.6
1	A	712	ASN	2.6
1	B	480	ILE	2.6
1	A	508	GLN	2.6
1	B	617	ASP	2.6
1	B	682	PRO	2.6
1	A	505	CYS	2.6
1	B	299	ARG	2.5
1	A	681	PRO	2.5
1	A	595	VAL	2.5
1	B	615	ASP	2.5
1	A	566	ALA	2.4
1	B	352	ASP	2.4
1	A	499	VAL	2.4
1	A	565	PRO	2.4
1	A	667	ARG	2.4
1	A	467	ASP	2.3
1	B	389	GLU	2.3
1	B	321	THR	2.3
1	A	683	MET	2.3
1	A	470	HIS	2.3
1	A	550	LYS	2.3
1	B	679	ILE	2.3
1	B	681	PRO	2.3
1	A	370	LYS	2.2
1	A	322	LEU	2.2
1	A	686	SER	2.2
1	B	375	LYS	2.2
1	A	679	ILE	2.2
1	A	446	VAL	2.2
1	A	572	LEU	2.2
1	A	328	GLU	2.2
1	B	678	TRP	2.1
1	A	485	TYR	2.1
1	B	676	TRP	2.1
1	A	704	PHE	2.1
1	A	568	SER	2.1
1	A	617	ASP	2.1

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	B	593	ILE	2.1
1	A	618	MET	2.0
1	A	710	PRO	2.0
1	A	711	TRP	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 6.4 Ligands [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
5	Y2B	A	800	28/28	0.81	0.20	33,42,44,44	0
5	Y2B	B	800	28/28	0.83	0.17	31,38,42,47	0
4	ACT	A	860	4/4	0.93	0.22	51,53,53,55	0
3	H4B	A	760	17/17	0.95	0.20	32,35,39,40	0
3	H4B	B	760	17/17	0.96	0.18	32,34,38,39	0
4	ACT	B	860	4/4	0.97	0.10	47,47,49,49	0
2	HEM	A	750	43/43	0.97	0.21	26,34,41,44	0
2	HEM	B	750	43/43	0.98	0.17	24,31,41,46	0
6	ZN	A	900	1/1	0.99	0.07	38,38,38,38	0




The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

**Electron density around Y2B A 800:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)

**Electron density around Y2B B 800:**




$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)






**Electron density around HEM A 750:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)





## 6.5 Other polymers [i](#)

There are no such residues in this entry.