



Full wwPDB X-ray Structure Validation Report ⓘ

Apr 19, 2022 – 10:03 AM EDT

PDB ID : 7RQH
Title : Crystal Structure of carboxyl-terminal processing protease A mutant S302A, CtpA_S302A, of *Pseudomonas aeruginosa*
Authors : Hsu, H.C.; Li, H.
Deposited on : 2021-08-06
Resolution : 3.20 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Xtriage (Phenix)	:	1.13
EDS	:	2.27
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac	:	5.8.0158
CCP4	:	7.0.044 (Gargrove)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.27


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION



The reported resolution of this entry is 3.20 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	1133 (3.20-3.20)
Clashscore	141614	1253 (3.20-3.20)
Ramachandran outliers	138981	1234 (3.20-3.20)
Sidechain outliers	138945	1233 (3.20-3.20)
RSRZ outliers	127900	1095 (3.20-3.20)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	402	
1	B	402	

2 Entry composition

There is only 1 type of molecule in this entry. The entry contains 3931 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Probable carboxyl-terminal protease.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	258	Total	C	N	O	S	0	0	0
			1962	1234	345	381	2			
1	B	258	Total	C	N	O	S	0	0	0
			1969	1238	348	381	2			

There are 10 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	34	GLY	-	expression tag	UNP Q9HU50
A	35	SER	-	expression tag	UNP Q9HU50
A	36	HIS	-	expression tag	UNP Q9HU50
A	37	MET	-	expression tag	UNP Q9HU50
A	302	ALA	SER	engineered mutation	UNP Q9HU50
B	34	GLY	-	expression tag	UNP Q9HU50
B	35	SER	-	expression tag	UNP Q9HU50
B	36	HIS	-	expression tag	UNP Q9HU50
B	37	MET	-	expression tag	UNP Q9HU50
B	302	ALA	SER	engineered mutation	UNP Q9HU50

GLY	ASN	GLY	GLY	ALA	ASP	ARG	PRO	THR	GLY	LYS	ARG	ALA	ALA	PRO	SER	E411	D417	A423	L424	R434	GLY
-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	------	------	------	------	------	-----

4 Data and refinement statistics

Property	Value	Source
Space group	H 3	Depositor
Cell constants a, b, c, α , β , γ	189.65Å 189.65Å 131.10Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	43.03 – 3.20 43.03 – 3.20	Depositor EDS
% Data completeness (in resolution range)	99.5 (43.03-3.20) 99.5 (43.03-3.20)	Depositor EDS
R_{merge}	0.09	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.55 (at 3.19Å)	Xtriage
Refinement program	PHENIX (1.19_4092)	Depositor
R, R_{free}	0.241 , 0.265 0.238 , 0.263	Depositor DCC
R_{free} test set	1415 reflections (4.90%)	wwPDB-VP
Wilson B-factor (Å ²)	127.4	Xtriage
Anisotropy	0.054	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.34 , 108.3	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	0.026 for h,-h-k,-l	Xtriage
F_o, F_c correlation	0.93	EDS
Total number of atoms	3931	wwPDB-VP
Average B, all atoms (Å ²)	126.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.30% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.27	0/1985	0.57	0/2687
1	B	0.27	0/1992	0.55	0/2696
All	All	0.27	0/3977	0.56	0/5383

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1962	0	2021	35	0
1	B	1969	0	2031	40	0
All	All	3931	0	4052	75	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 9.

All (75) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:44:ASP:OD1	1:A:45:GLU:N	2.19	0.76
1:A:78:LEU:HD12	1:A:85:SER:HB3	1.69	0.75

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:242:ARG:NH2	1:B:417:ASP:OD2	2.26	0.69
1:B:208:ILE:O	1:B:244:ASN:ND2	2.28	0.66
1:B:252:ALA:HB2	1:B:303:ALA:HB1	1.77	0.66
1:A:260:LEU:HD21	1:A:314:GLN:HG3	1.76	0.66
1:B:258:ALA:HA	1:B:286:PRO:HD2	1.81	0.63
1:A:252:ALA:HB2	1:A:303:ALA:HB1	1.82	0.62
1:B:199:LEU:HD23	1:B:205:TYR:HB2	1.83	0.61
1:A:277:LEU:HD13	1:A:279:PHE:HE2	1.66	0.60
1:A:204:ALA:HB1	1:A:226:LEU:HD13	1.87	0.56
1:A:243:ASN:OD1	1:A:297:ASN:ND2	2.34	0.56
1:A:272:ILE:HG22	1:A:275:SER:HB3	1.88	0.55
1:A:239:LEU:HB2	1:A:294:VAL:HG22	1.91	0.53
1:A:258:ALA:HA	1:A:286:PRO:HD2	1.92	0.52
1:B:242:ARG:HB2	1:B:297:ASN:HB2	1.91	0.52
1:B:214:ASN:O	1:B:217:GLU:HG2	2.10	0.52
1:B:259:PHE:O	1:B:316:ARG:NH1	2.42	0.51
1:B:293:VAL:HG11	1:B:423:ALA:HB1	1.91	0.51
1:A:295:LEU:HD23	1:A:320:MET:HB3	1.93	0.51
1:B:216:GLY:HA3	1:B:254:GLU:HG3	1.93	0.51
1:A:202:GLY:N	1:A:230:ASN:OD1	2.42	0.50
1:B:351:THR:OG1	1:B:355:ARG:HG2	2.12	0.50
1:B:204:ALA:HB1	1:B:226:LEU:HD13	1.93	0.50
1:A:366:ILE:HG22	1:A:368:VAL:HG23	1.95	0.49
1:A:250:GLN:O	1:A:253:VAL:HG22	2.13	0.49
1:A:305:GLU:OE2	1:A:326:GLY:N	2.30	0.49
1:A:253:VAL:O	1:A:257:ASP:HB2	2.13	0.49
1:B:41:LEU:HD21	1:B:43:LEU:HG	1.95	0.48
1:B:55:ARG:HD3	1:B:55:ARG:HA	1.71	0.48
1:B:290:VAL:O	1:B:316:ARG:NH2	2.39	0.47
1:A:194:VAL:HG12	1:A:218:GLU:HB3	1.97	0.47
1:B:268:THR:O	1:B:276:GLU:HA	2.14	0.47
1:B:215:THR:HB	1:B:251:SER:OG	2.15	0.46
1:B:216:GLY:O	1:B:219:VAL:HG22	2.15	0.46
1:B:331:GLN:HG3	1:B:345:THR:HG22	1.98	0.46
1:A:411:GLU:HG2	1:A:412:ARG:H	1.80	0.46
1:B:366:ILE:HG22	1:B:368:VAL:HG23	1.96	0.46
1:A:43:LEU:HA	1:A:46:LEU:HB3	1.98	0.46
1:A:339:ASP:N	1:A:339:ASP:OD2	2.48	0.46
1:B:242:ARG:CB	1:B:297:ASN:HB2	2.46	0.45
1:B:305:GLU:HG2	1:B:362:ILE:HG12	1.99	0.45
1:A:272:ILE:CG2	1:A:275:SER:HB3	2.46	0.45

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:434:ARG:HD3	1:B:434:ARG:HA	1.60	0.45
1:B:251:SER:O	1:B:254:GLU:HG2	2.17	0.45
1:A:199:LEU:HD11	1:A:205:TYR:HB2	1.98	0.45
1:A:429:GLY:O	1:A:433:THR:HG23	2.18	0.44
1:B:199:LEU:HD11	1:B:424:LEU:HD13	1.99	0.44
1:B:222:ALA:O	1:B:226:LEU:HG	2.17	0.44
1:B:328:GLY:O	1:B:358:GLN:HB2	2.18	0.44
1:A:211:PHE:CE1	1:A:252:ALA:HA	2.53	0.44
1:A:268:THR:HG22	1:A:349:TYR:HD1	1.82	0.44
1:B:223:LEU:HD23	1:B:286:PRO:HB2	1.99	0.44
1:A:294:VAL:HB	1:A:319:LEU:CD2	2.48	0.43
1:A:368:VAL:HG12	1:A:419:GLN:HG3	1.99	0.43
1:A:266:VAL:HG21	1:A:306:ILE:HG12	2.00	0.43
1:B:193:SER:OG	1:B:212:GLN:OE1	2.35	0.43
1:A:196:SER:HB2	1:A:226:LEU:HD11	2.00	0.42
1:A:269:LYS:HB2	1:A:350:TYR:CE1	2.54	0.42
1:B:219:VAL:O	1:B:223:LEU:HD13	2.18	0.42
1:B:315:LYS:HB2	1:B:315:LYS:HE2	1.79	0.42
1:B:326:GLY:HA2	1:B:361:GLY:HA2	2.01	0.42
1:B:295:LEU:HD23	1:B:320:MET:HG3	2.01	0.42
1:B:312:GLN:HB2	1:B:319:LEU:HD23	2.02	0.42
1:A:268:THR:HG22	1:A:349:TYR:CD1	2.54	0.42
1:A:261:THR:O	1:A:283:PRO:HA	2.20	0.42
1:B:306:ILE:HD11	1:B:349:TYR:CD2	2.55	0.42
1:B:251:SER:HA	1:B:254:GLU:HG2	2.02	0.41
1:B:267:TYR:HA	1:B:277:LEU:O	2.19	0.41
1:A:191:VAL:HG23	1:A:212:GLN:HE22	1.85	0.41
1:A:82:ASP:HB3	1:A:85:SER:OG	2.20	0.41
1:B:227:ARG:NH2	1:B:286:PRO:O	2.54	0.41
1:B:319:LEU:HB3	1:B:364:PRO:HA	2.03	0.41
1:A:194:VAL:CG1	1:A:218:GLU:HB3	2.51	0.41
1:B:208:ILE:HD12	1:B:241:LEU:HD21	2.02	0.40

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries

of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	252/402 (63%)	244 (97%)	8 (3%)	0	100	100
1	B	252/402 (63%)	246 (98%)	6 (2%)	0	100	100
All	All	504/804 (63%)	490 (97%)	14 (3%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	213/322 (66%)	212 (100%)	1 (0%)	88	95
1	B	214/322 (66%)	209 (98%)	5 (2%)	50	78
All	All	427/644 (66%)	421 (99%)	6 (1%)	67	86

All (6) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	325	PHE
1	B	55	ARG
1	B	323	ASP
1	B	327	LYS
1	B	330	VAL
1	B	377	ARG

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
1	B	244	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ > 2			OWAB(Å ²)	Q < 0.9
1	A	258/402 (64%)	0.15	8 (3%)	49	32	84, 121, 173, 206	0
1	B	258/402 (64%)	0.18	9 (3%)	44	28	84, 121, 180, 218	0
All	All	516/804 (64%)	0.17	17 (3%)	46	30	84, 121, 176, 218	0

All (17) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	377	ARG	3.3
1	A	191	VAL	2.9
1	B	45	GLU	2.7
1	A	49	PHE	2.4
1	B	43	LEU	2.4
1	B	49	PHE	2.4
1	B	44	ASP	2.3
1	A	56	VAL	2.3
1	A	70	LEU	2.3
1	A	360	GLN	2.3
1	B	411	GLU	2.2
1	A	336	LEU	2.2
1	B	336	LEU	2.2
1	A	337	ASN	2.1
1	B	335	PRO	2.0
1	B	64	VAL	2.0
1	A	270	GLY	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.