



# Full wwPDB NMR Structure Validation Report ⓘ

Dec 3, 2020 – 09:07 AM GMT

PDB ID : 6RSF  
Title : NMR structure of pleurocidin KR in SDS micelles  
Authors : Manzo, G.; Mason, A.J.  
Deposited on : 2019-05-21

This is a Full wwPDB NMR Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/NMRValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

Cyrange : Kirchner and Güntert (2011)  
NmrClust : Kelley et al. (1996)  
MolProbity : 4.02b-467  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
RCI : v\_1n\_11\_5\_13\_A (Berjanski et al., 2005)  
PANAV : Wang et al. (2010)  
ShiftChecker : 2.14.6  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.14.6


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*SOLUTION NMR*


The overall completeness of chemical shifts assignment is 24%.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	NMR archive (#Entries)
Clashscore	158937	12864
Ramachandran outliers	154571	11451
Sidechain outliers	154315	11428

The table below summarises the geometric issues observed across the polymeric chains and their fit to the experimental data. The red, orange, yellow and green segments indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A cyan segment indicates the fraction of residues that are not part of the well-defined cores, and a grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$

Mol	Chain	Length	Quality of chain
1	A	25	

## 2 Ensemble composition and analysis ⓘ

This entry contains 50 models.

Cyrange was unable to find well-defined residues.

Error message: The number of core atoms ( 3) was below the domain threshold value ( 8).

NmrClust was unable to cluster the ensemble.

Error message: Wrapper check: not enough residues in core to run NmrClust

### 3 Entry composition

There is only 1 type of molecule in this entry. The entry contains 397 atoms, of which 196 are hydrogens and 0 are deuteriums.

- Molecule 1 is a protein called Pleurocidin.

Mol	Chain	Residues	Atoms					Trace
1	A	25	Total	C	H	N	O	0
			397	129	196	44	28	

There are 4 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	7	ARG	LYS	engineered mutation	UNP Q90ZY0
A	8	ARG	LYS	engineered mutation	UNP Q90ZY0
A	14	ARG	LYS	engineered mutation	UNP Q90ZY0
A	18	ARG	LYS	engineered mutation	UNP Q90ZY0

## 4 Residue-property plots

### 4.1 Average score per residue in the NMR ensemble

These plots are provided for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic is the same as shown in the summary in section 1 of this report. The second graphic shows the sequence where residues are colour-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outliers are shown as green connectors. Residues which are classified as ill-defined in the NMR ensemble, are shown in cyan with an underline colour-coded according to the previous scheme. Residues which were present in the experimental sample, but not modelled in the final structure are shown in grey.

- Molecule 1: Pleurocidin




### 4.2 Scores per residue for each member of the ensemble

Colouring as in section 4.1 above.


#### 4.2.1 Score per residue for model 1

- Molecule 1: Pleurocidin




#### 4.2.2 Score per residue for model 2

- Molecule 1: Pleurocidin



### 4.2.3 Score per residue for model 3

- Molecule 1: Pleurocidin




### 4.2.4 Score per residue for model 4

- Molecule 1: Pleurocidin



### 4.2.5 Score per residue for model 5

- Molecule 1: Pleurocidin




### 4.2.6 Score per residue for model 6

- Molecule 1: Pleurocidin




### 4.2.7 Score per residue for model 7

- Molecule 1: Pleurocidin



#### 4.2.8 Score per residue for model 8

- Molecule 1: Pleurocidin



#### 4.2.9 Score per residue for model 9

- Molecule 1: Pleurocidin



#### 4.2.10 Score per residue for model 10

- Molecule 1: Pleurocidin




#### 4.2.11 Score per residue for model 11

- Molecule 1: Pleurocidin



#### 4.2.12 Score per residue for model 12

- Molecule 1: Pleurocidin




#### 4.2.13 Score per residue for model 13

- Molecule 1: Pleurocidin




#### 4.2.14 Score per residue for model 14

- Molecule 1: Pleurocidin




#### 4.2.15 Score per residue for model 15

- Molecule 1: Pleurocidin



#### 4.2.16 Score per residue for model 16

- Molecule 1: Pleurocidin



#### 4.2.17 Score per residue for model 17

- Molecule 1: Pleurocidin





#### 4.2.18 Score per residue for model 18

- Molecule 1: Pleurocidin




#### 4.2.19 Score per residue for model 19

- Molecule 1: Pleurocidin




#### 4.2.20 Score per residue for model 20

- Molecule 1: Pleurocidin




#### 4.2.21 Score per residue for model 21

- Molecule 1: Pleurocidin




#### 4.2.22 Score per residue for model 22

- Molecule 1: Pleurocidin



#### 4.2.23 Score per residue for model 23

- Molecule 1: Pleurocidin




#### 4.2.24 Score per residue for model 24

- Molecule 1: Pleurocidin




#### 4.2.25 Score per residue for model 25

- Molecule 1: Pleurocidin




#### 4.2.26 Score per residue for model 26

- Molecule 1: Pleurocidin




#### 4.2.27 Score per residue for model 27

- Molecule 1: Pleurocidin




#### 4.2.28 Score per residue for model 28

- Molecule 1: Pleurocidin




#### 4.2.29 Score per residue for model 29

- Molecule 1: Pleurocidin



#### 4.2.30 Score per residue for model 30

- Molecule 1: Pleurocidin




#### 4.2.31 Score per residue for model 31

- Molecule 1: Pleurocidin




#### 4.2.32 Score per residue for model 32

- Molecule 1: Pleurocidin




#### 4.2.33 Score per residue for model 33

- Molecule 1: Pleurocidin




#### 4.2.34 Score per residue for model 34

- Molecule 1: Pleurocidin



#### 4.2.35 Score per residue for model 35

- Molecule 1: Pleurocidin



#### 4.2.36 Score per residue for model 36

- Molecule 1: Pleurocidin



#### 4.2.37 Score per residue for model 37


- Molecule 1: Pleurocidin



#### 4.2.38 Score per residue for model 38

- Molecule 1: Pleurocidin

Chain A:  44% 40% 12% .



#### 4.2.39 Score per residue for model 39

- Molecule 1: Pleurocidin

Chain A:  44% 48% . .



#### 4.2.40 Score per residue for model 40


- Molecule 1: Pleurocidin

Chain A:  44% 44% 12%



#### 4.2.41 Score per residue for model 41

- Molecule 1: Pleurocidin


Chain A:  36% 60% .



#### 4.2.42 Score per residue for model 42

- Molecule 1: Pleurocidin

Chain A:  56% 24% 20%




#### 4.2.43 Score per residue for model 43

- Molecule 1: Pleurocidin




#### 4.2.44 Score per residue for model 44

- Molecule 1: Pleurocidin



#### 4.2.45 Score per residue for model 45

- Molecule 1: Pleurocidin




#### 4.2.46 Score per residue for model 46

- Molecule 1: Pleurocidin



#### 4.2.47 Score per residue for model 47

- Molecule 1: Pleurocidin



#### 4.2.48 Score per residue for model 48

- Molecule 1: Pleurocidin

Chain A:  44% 52% .



#### 4.2.49 Score per residue for model 49

- Molecule 1: Pleurocidin


Chain A:  40% 36% 20% .



#### 4.2.50 Score per residue for model 50

- Molecule 1: Pleurocidin

Chain A:  52% 36% 12% .



## 5 Refinement protocol and experimental data overview

The models were refined using the following method: *simulated annealing*.

Of the 500 calculated structures, 50 were deposited, based on the following criterion: *structures with the lowest energy*.

The following table shows the software used for structure solution, optimisation and refinement.

Software name	Classification	Version
DYNAMO	structure calculation	

The following table shows chemical shift validation statistics as aggregates over all chemical shift files. Detailed validation can be found in section 7 of this report.

Chemical shift file(s)	working_cs.cif
Number of chemical shift lists	1
Total number of shifts	103
Number of shifts mapped to atoms	103
Number of unparsed shifts	0
Number of shifts with mapping errors	0
Number of shifts with mapping warnings	0
Assignment completeness (well-defined parts)	24%

No validations of the models with respect to experimental NMR restraints is performed at this time.



## 6 Model quality [i](#)

### 6.1 Standard geometry [i](#)

There are no covalent bond-length or bond-angle outliers.

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	Chirality	Planarity
1	A	0.0±0.0	3.6±0.6
All	All	0	181

There are no bond-length outliers.

There are no bond-angle outliers.

There are no chirality outliers.

All unique planar outliers are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Group	Models (Total)
1	A	8	ARG	Sidechain	48
1	A	18	ARG	Sidechain	47
1	A	14	ARG	Sidechain	44
1	A	7	ARG	Sidechain	42

### 6.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in each chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes averaged over the ensemble.

Mol	Chain	Non-H	H(model)	H(added)	Clashes
1	A	201	196	196	5±2
All	All	10050	9800	9800	230

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 12.

All unique clashes are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Clash(Å)	Distance(Å)	Models	
				Worst	Total
1:A:2:TRP:CG	1:A:3:GLY:N	0.58	2.71	23	3
1:A:2:TRP:CD1	1:A:3:GLY:N	0.58	2.71	23	1
1:A:11:HIS:N	1:A:11:HIS:HD1	0.55	2.00	44	1
1:A:21:LEU:N	1:A:21:LEU:HD12	0.54	2.17	5	4
1:A:21:LEU:HD12	1:A:21:LEU:N	0.54	2.17	48	6
1:A:25:LEU:HD12	1:A:25:LEU:N	0.53	2.19	41	3
1:A:2:TRP:CD2	1:A:3:GLY:N	0.53	2.77	25	2
1:A:25:LEU:N	1:A:25:LEU:HD12	0.53	2.19	4	3
1:A:15:HIS:CG	1:A:16:VAL:N	0.52	2.75	40	3
1:A:25:LEU:N	1:A:25:LEU:CD2	0.51	2.73	35	1
1:A:21:LEU:CD1	1:A:21:LEU:N	0.51	2.73	31	5
1:A:11:HIS:O	1:A:13:GLY:N	0.51	2.44	31	9
1:A:11:HIS:ND1	1:A:11:HIS:N	0.51	2.56	44	2
1:A:25:LEU:CD1	1:A:25:LEU:N	0.50	2.74	4	4
1:A:25:LEU:N	1:A:25:LEU:HD23	0.50	2.21	35	1
1:A:24:TYR:CD2	1:A:25:LEU:N	0.50	2.79	22	1
1:A:15:HIS:CG	1:A:16:VAL:H	0.50	2.24	31	1
1:A:23:HIS:O	1:A:25:LEU:N	0.50	2.44	24	3
1:A:5:PHE:O	1:A:6:PHE:CG	0.50	2.65	46	17
1:A:21:LEU:O	1:A:23:HIS:N	0.50	2.45	3	1
1:A:5:PHE:O	1:A:5:PHE:CD1	0.50	2.65	12	1
1:A:11:HIS:H	1:A:11:HIS:HD1	0.50	1.50	48	1
1:A:1:GLY:O	1:A:2:TRP:CG	0.50	2.65	21	7
1:A:23:HIS:CD2	1:A:23:HIS:H	0.50	2.24	42	2
1:A:22:THR:O	1:A:23:HIS:ND1	0.50	2.45	8	2
1:A:23:HIS:CG	1:A:24:TYR:H	0.50	2.25	43	1
1:A:22:THR:O	1:A:23:HIS:CD2	0.49	2.65	12	1
1:A:9:ALA:O	1:A:11:HIS:N	0.49	2.46	2	1
1:A:15:HIS:O	1:A:15:HIS:CG	0.49	2.65	16	1
1:A:10:ALA:O	1:A:11:HIS:CG	0.49	2.65	23	5
1:A:1:GLY:O	1:A:2:TRP:CD1	0.49	2.65	11	3
1:A:21:LEU:N	1:A:21:LEU:CD1	0.49	2.75	48	5
1:A:13:GLY:O	1:A:15:HIS:N	0.49	2.46	32	1
1:A:14:ARG:NH1	1:A:17:GLY:N	0.49	2.60	39	1
1:A:4:SER:O	1:A:5:PHE:CG	0.49	2.65	50	3
1:A:4:SER:O	1:A:6:PHE:CE1	0.49	2.66	39	3
1:A:6:PHE:O	1:A:6:PHE:CD1	0.49	2.65	25	2
1:A:22:THR:O	1:A:24:TYR:CD1	0.49	2.66	36	2
1:A:14:ARG:O	1:A:16:VAL:N	0.49	2.46	49	8
1:A:22:THR:O	1:A:23:HIS:CG	0.49	2.65	34	2
1:A:4:SER:O	1:A:6:PHE:CE2	0.49	2.66	36	1
1:A:23:HIS:O	1:A:24:TYR:CG	0.49	2.65	37	1

Continued on next page...

*Continued from previous page...*

Atom-1	Atom-2	Clash(Å)	Distance(Å)	Models	
				Worst	Total
1:A:23:HIS:N	1:A:23:HIS:ND1	0.48	2.59	17	3
1:A:6:PHE:CD1	1:A:6:PHE:O	0.48	2.66	31	2
1:A:23:HIS:CG	1:A:23:HIS:O	0.48	2.66	49	2
1:A:23:HIS:CE1	1:A:25:LEU:O	0.48	2.66	9	1
1:A:11:HIS:N	1:A:11:HIS:ND1	0.48	2.61	9	5
1:A:18:ARG:N	1:A:18:ARG:CD	0.48	2.76	4	1
1:A:6:PHE:CD2	1:A:6:PHE:O	0.48	2.66	42	2
1:A:5:PHE:O	1:A:6:PHE:CD1	0.48	2.67	14	2
1:A:22:THR:O	1:A:24:TYR:N	0.48	2.46	23	4
1:A:24:TYR:CG	1:A:25:LEU:N	0.48	2.81	11	1
1:A:22:THR:O	1:A:24:TYR:CD2	0.48	2.66	17	1
1:A:6:PHE:O	1:A:6:PHE:CG	0.48	2.66	25	1
1:A:25:LEU:HD23	1:A:25:LEU:N	0.48	2.24	22	1
1:A:23:HIS:CG	1:A:24:TYR:N	0.47	2.81	43	2
1:A:8:ARG:HE	1:A:8:ARG:N	0.47	2.07	6	1
1:A:15:HIS:N	1:A:15:HIS:ND1	0.47	2.61	50	1
1:A:18:ARG:O	1:A:20:ALA:N	0.47	2.48	12	1
1:A:23:HIS:ND1	1:A:23:HIS:N	0.47	2.62	28	2
1:A:10:ALA:O	1:A:11:HIS:ND1	0.47	2.48	13	1
1:A:2:TRP:CG	1:A:2:TRP:O	0.47	2.68	18	4
1:A:25:LEU:CD2	1:A:25:LEU:N	0.47	2.77	22	1
1:A:23:HIS:ND1	1:A:23:HIS:O	0.47	2.48	7	2
1:A:22:THR:OG1	1:A:23:HIS:N	0.46	2.48	5	1
1:A:9:ALA:O	1:A:11:HIS:ND1	0.46	2.48	44	1
1:A:24:TYR:CD1	1:A:25:LEU:N	0.45	2.83	11	1
1:A:11:HIS:O	1:A:11:HIS:ND1	0.45	2.49	27	1
1:A:21:LEU:O	1:A:23:HIS:CE1	0.45	2.69	40	1
1:A:23:HIS:O	1:A:23:HIS:CG	0.45	2.69	3	2
1:A:21:LEU:O	1:A:23:HIS:ND1	0.45	2.49	40	1
1:A:5:PHE:C	1:A:6:PHE:CG	0.45	2.90	15	4
1:A:15:HIS:CD2	1:A:16:VAL:N	0.45	2.85	31	1
1:A:1:GLY:C	1:A:2:TRP:CG	0.45	2.91	32	2
1:A:10:ALA:C	1:A:11:HIS:CG	0.44	2.91	13	1
1:A:4:SER:O	1:A:6:PHE:N	0.44	2.50	44	1
1:A:10:ALA:C	1:A:11:HIS:CD2	0.44	2.91	27	1
1:A:4:SER:C	1:A:5:PHE:CG	0.44	2.91	31	2
1:A:23:HIS:N	1:A:23:HIS:CD2	0.44	2.84	10	1
1:A:5:PHE:C	1:A:6:PHE:CD1	0.44	2.91	14	3
1:A:5:PHE:C	1:A:6:PHE:CD2	0.44	2.92	21	2
1:A:2:TRP:O	1:A:2:TRP:CG	0.44	2.70	42	2
1:A:22:THR:C	1:A:24:TYR:N	0.43	2.71	3	1

*Continued on next page...*

Continued from previous page...

Atom-1	Atom-2	Clash(Å)	Distance(Å)	Models	
				Worst	Total
1:A:8:ARG:O	1:A:10:ALA:N	0.43	2.50	11	1
1:A:15:HIS:ND1	1:A:15:HIS:C	0.43	2.72	28	2
1:A:8:ARG:C	1:A:10:ALA:N	0.43	2.71	43	4
1:A:4:SER:C	1:A:5:PHE:CD1	0.42	2.92	50	1
1:A:23:HIS:CD2	1:A:23:HIS:O	0.42	2.72	29	1
1:A:16:VAL:HG13	1:A:17:GLY:N	0.42	2.30	9	2
1:A:2:TRP:CD1	1:A:2:TRP:O	0.42	2.73	9	1
1:A:6:PHE:CG	1:A:6:PHE:O	0.42	2.72	44	1
1:A:25:LEU:N	1:A:25:LEU:CD1	0.41	2.83	8	2
1:A:2:TRP:N	1:A:2:TRP:CD1	0.41	2.88	3	1
1:A:10:ALA:O	1:A:11:HIS:CD2	0.41	2.73	23	1
1:A:9:ALA:O	1:A:10:ALA:O	0.41	2.39	3	1
1:A:3:GLY:O	1:A:5:PHE:N	0.41	2.54	28	1
1:A:12:VAL:HG22	1:A:13:GLY:N	0.41	2.29	5	2
1:A:21:LEU:O	1:A:22:THR:O	0.41	2.39	19	1
1:A:10:ALA:C	1:A:12:VAL:N	0.41	2.72	26	1
1:A:9:ALA:C	1:A:11:HIS:N	0.41	2.74	47	1
1:A:2:TRP:O	1:A:2:TRP:CD1	0.41	2.74	22	1
1:A:11:HIS:CG	1:A:11:HIS:O	0.41	2.74	27	1
1:A:14:ARG:O	1:A:19:ALA:N	0.41	2.55	21	1
1:A:22:THR:O	1:A:23:HIS:O	0.40	2.39	50	1
1:A:15:HIS:ND1	1:A:15:HIS:N	0.40	2.67	17	1
1:A:18:ARG:O	1:A:19:ALA:O	0.40	2.40	34	1
1:A:11:HIS:CD2	1:A:11:HIS:H	0.40	2.35	43	1
1:A:11:HIS:CD2	1:A:11:HIS:N	0.40	2.90	24	1
1:A:6:PHE:C	1:A:8:ARG:N	0.40	2.75	29	1
1:A:16:VAL:HG23	1:A:17:GLY:N	0.40	2.32	7	1
1:A:4:SER:O	1:A:5:PHE:CD2	0.40	2.75	17	1

## 6.3 Torsion angles ⓘ

### 6.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the backbone conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	23/25 (92%)	8±3 (36±11%)	7±3 (31±11%)	8±2 (33±10%)	0 0

Continued on next page...

*Continued from previous page...*

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
All	All	1150/1250 (92%)	418 (36%)	351 (31%)	381 (33%)	0 0

All 23 unique Ramachandran outliers are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Models (Total)
1	A	2	TRP	27
1	A	13	GLY	25
1	A	24	TYR	24
1	A	3	GLY	23
1	A	23	HIS	23
1	A	20	ALA	23
1	A	6	PHE	22
1	A	10	ALA	22
1	A	15	HIS	22
1	A	19	ALA	20
1	A	9	ALA	19
1	A	11	HIS	18
1	A	8	ARG	17
1	A	12	VAL	14
1	A	5	PHE	13
1	A	22	THR	13
1	A	14	ARG	11
1	A	4	SER	11
1	A	16	VAL	8
1	A	7	ARG	8
1	A	17	GLY	7
1	A	21	LEU	6
1	A	18	ARG	5

### 6.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the sidechain conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	17/17 (100%)	16±1 (96±4%)	1±1 (4±4%)	34 82
All	All	850/850 (100%)	815 (96%)	35 (4%)	34 82

All 7 unique residues with a non-rotameric sidechain are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Models (Total)
1	A	11	HIS	11
1	A	23	HIS	8
1	A	15	HIS	7
1	A	25	LEU	6
1	A	21	LEU	1
1	A	18	ARG	1
1	A	22	THR	1

### 6.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 6.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 6.6 Ligand geometry [i](#)

There are no ligands in this entry.

## 6.7 Other polymers [i](#)

There are no such molecules in this entry.

## 6.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 7 Chemical shift validation

The completeness of assignment taking into account all chemical shift lists is 24% for the well-defined parts and 24% for the entire structure.

### 7.1 Chemical shift list 1

File name: `working_cs.cif`

Chemical shift list name: *starch\_output*

#### 7.1.1 Bookkeeping

The following table shows the results of parsing the chemical shift list and reports the number of nuclei with statistically unusual chemical shifts.

Total number of shifts	103
Number of shifts mapped to atoms	103
Number of unparsed shifts	0
Number of shifts with mapping errors	0
Number of shifts with mapping warnings	0
Number of shift outliers (ShiftChecker)	0

#### 7.1.2 Chemical shift referencing

No chemical shift referencing corrections were calculated (not enough data).

#### 7.1.3 Completeness of resonance assignments

The following table shows the completeness of the chemical shift assignments for the well-defined regions of the structure. The overall completeness is 24%, i.e. 77 atoms were assigned a chemical shift out of a possible 322. 0 out of 4 assigned methyl groups (LEU and VAL) were assigned stereospecifically.

	Total	<sup>1</sup> H	<sup>13</sup> C	<sup>15</sup> N
Backbone	42/125 (34%)	42/50 (84%)	0/50 (0%)	0/25 (0%)
Sidechain	35/138 (25%)	35/82 (43%)	0/44 (0%)	0/12 (0%)
Aromatic	0/59 (0%)	0/32 (0%)	0/23 (0%)	0/4 (0%)
Overall	77/322 (24%)	77/164 (47%)	0/117 (0%)	0/41 (0%)

The following table shows the completeness of the chemical shift assignments for the full structure. The overall completeness is 24%, i.e. 77 atoms were assigned a chemical shift out of a possible 322. 0 out of 4 assigned methyl groups (LEU and VAL) were assigned stereospecifically.

	<b>Total</b>	<b><sup>1</sup>H</b>	<b><sup>13</sup>C</b>	<b><sup>15</sup>N</b>
Backbone	42/125 (34%)	42/50 (84%)	0/50 (0%)	0/25 (0%)
Sidechain	35/138 (25%)	35/82 (43%)	0/44 (0%)	0/12 (0%)
Aromatic	0/59 (0%)	0/32 (0%)	0/23 (0%)	0/4 (0%)
Overall	77/322 (24%)	77/164 (47%)	0/117 (0%)	0/41 (0%)

#### 7.1.4 Statistically unusual chemical shifts [i](#)

There are no statistically unusual chemical shifts.

#### 7.1.5 Random Coil Index (RCI) plots [i](#)

The image below reports *random coil index* values for the protein chains in the structure. The height of each bar gives a probability of a given residue to be disordered, as predicted from the available chemical shifts and the amino acid sequence. A value above 0.2 is an indication of significant predicted disorder. The colour of the bar shows whether the residue is in the well-defined core (black) or in the ill-defined residue ranges (cyan), as described in section 2 on ensemble composition.

Random coil index (RCI) for chain A:

