



Full wwPDB X-ray Structure Validation Report ⓘ

Aug 15, 2022 – 02:13 PM EDT

PDB ID : 7RST
Title : The Crystal Structure of Recombinant Chloroperoxidase Expressed in *Aspergillus niger*
Authors : Tang, X.; Venkadesh, S.; Zhou, J.; Rosen, B.; Wang, X.
Deposited on : 2021-08-11
Resolution : 1.69 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at <http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix)	:	1.13
EDS	:	2.29
buster-report	:	1.1.7 (2018)
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac	:	5.8.0158
CCP4	:	7.0.044 (Gargrove)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.29


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.69 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	4298 (1.70-1.70)
Clashscore	141614	4695 (1.70-1.70)
Ramachandran outliers	138981	4610 (1.70-1.70)
Sidechain outliers	138945	4610 (1.70-1.70)
RSRZ outliers	127900	4222 (1.70-1.70)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	299	
2	B	2	
2	C	2	
3	D	2	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard

residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
7	MAN	A	310	-	-	-	X

2 Entry composition [i](#)


There are 9 unique types of molecules in this entry. The entry contains 3113 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Chloroperoxidase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	299	Total	C	N	O	S	0	4	0
			2345	1485	389	465	6			

- Molecule 2 is an oligosaccharide called 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
2	B	2	Total	C	N	O	0	0	0
			28	16	2	10			
2	C	2	Total	C	N	O	0	0	0
			28	16	2	10			

- Molecule 3 is an oligosaccharide called alpha-D-mannopyranose-(1-2)-alpha-D-mannopyranose.




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf	Trace
3	D	2	Total	C	O	0	0	0
			22	12	10			

- Molecule 4 is MANGANESE (II) ION (three-letter code: MN) (formula: Mn).


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	1	Total	Mn	0	0
			1	1		

- Molecule 5 is PROTOPORPHYRIN IX CONTAINING FE (three-letter code: HEM) (formula: $C_{34}H_{32}FeN_4O_4$) (labeled as "Ligand of Interest" by depositor).




Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
5	A	1	Total 43	C 34	Fe 1	N 4	O 4	0	0

- Molecule 6 is 2-acetamido-2-deoxy-beta-D-glucopyranose (three-letter code: NAG) (formula: $\text{C}_8\text{H}_{15}\text{NO}_6$).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
6	A	1	Total	C	N	O	0	0
			14	8	1	5		

- Molecule 7 is alpha-D-mannopyranose (three-letter code: MAN) (formula: C₆H₁₂O₆).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
7	A	1	Total	C	O	0	0
			11	6	5		
7	A	1	Total	C	O	0	0
			11	6	5		
7	A	1	Total	C	O	0	0
			11	6	5		
7	A	1	Total	C	O	0	1
			22	12	10		
7	A	1	Total	C	O	0	0
			11	6	5		
7	A	1	Total	C	O	0	0
			11	6	5		
7	A	1	Total	C	O	0	0
			11	6	5		
7	A	1	Total	C	O	0	0
			11	6	5		
7	A	1	Total	C	O	0	0
			11	6	5		
7	A	1	Total	C	O	0	1
			22	12	10		

- Molecule 8 is IODIDE ION (three-letter code: IOD) (formula: I).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
8	A	10	Total	I	0	0
			10	10		

- Molecule 9 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
9	A	468	Total	O	0	0
			468	468		

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Chloroperoxidase

Chain A:  96% ..



- Molecule 2: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain B:  50% 50%



- Molecule 2: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain C:  100%



- Molecule 3: alpha-D-mannopyranose-(1-2)-alpha-D-mannopyranose

Chain D:  100%



4 Data and refinement statistics

Property	Value	Source
Space group	C 2 2 21	Depositor
Cell constants a, b, c, α , β , γ	58.05Å 150.92Å 100.85Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	30.36 – 1.69 37.73 – 1.69	Depositor EDS
% Data completeness (in resolution range)	90.6 (30.36-1.69) 90.6 (37.73-1.69)	Depositor EDS
R_{merge}	0.11	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.27 (at 1.69Å)	Xtriage
Refinement program	PHENIX 1.19.2_4158	Depositor
R, R_{free}	0.170 , 0.210 0.170 , 0.209	Depositor DCC
R_{free} test set	2292 reflections (5.06%)	wwPDB-VP
Wilson B-factor (Å ²)	18.3	Xtriage
Anisotropy	0.053	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.33 , 40.5	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	3113	wwPDB-VP
Average B, all atoms (Å ²)	22.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.17% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: NAG, PCA, MAN, HEM, MN, IOD

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.39	0/2414	0.62	0/3311

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2345	0	2154	9	0
2	B	28	0	25	1	0
2	C	28	0	25	0	0
3	D	22	0	19	0	0
4	A	1	0	0	0	0
5	A	43	0	30	3	0
6	A	14	0	13	0	0
7	A	154	0	140	4	0
8	A	10	0	0	1	0
9	A	468	0	0	2	0
All	All	3113	0	2406	12	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (12) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:242:SER:HB3	7:A:305[B]:MAN:H5	1.53	0.90
1:A:242:SER:HB3	7:A:305[B]:MAN:C5	2.06	0.85
1:A:166:GLU:OE2	1:A:255:SER:OG	1.99	0.79
8:A:315:IOD:I	9:A:779:HOH:O	2.72	0.77
1:A:258:SER:HB3	7:A:311:MAN:H5	1.70	0.74
1:A:258:SER:HB3	7:A:311:MAN:C5	2.22	0.69
1:A:201:GLU:OE1	9:A:401:HOH:O	2.15	0.64
5:A:300:HEM:HMB2	5:A:300:HEM:HBB2	1.88	0.56
5:A:300:HEM:HBC2	5:A:300:HEM:HMC1	1.91	0.52
1:A:186:PHE:HB3	5:A:300:HEM:CAC	2.45	0.46
1:A:212:TYR:O	1:A:216:ASN:HB2	2.19	0.42
1:A:82:VAL:HG13	2:B:2:NAG:H82	2.02	0.41

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	301/299 (101%)	293 (97%)	8 (3%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	255/251 (102%)	249 (98%)	6 (2%)	49 31

All (6) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	103	PHE
1	A	201	GLU
1	A	206	ARG
1	A	242	SER
1	A	258	SER
1	A	296	PRO

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	120	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

1 non-standard protein/DNA/RNA residue is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
1	PCA	A	0	1	7,8,9	1.59	1 (14%)	9,10,12	1.77	4 (44%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	PCA	A	0	1	-	0/0/11/13	0/1/1/1

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	0	PCA	CD-N	4.03	1.45	1.34

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	0	PCA	CA-N-CD	-2.62	104.60	113.58
1	A	0	PCA	OE-CD-CG	-2.55	122.31	126.76
1	A	0	PCA	CG-CD-N	2.22	114.14	108.39
1	A	0	PCA	CB-CA-N	2.10	109.33	103.30

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.5 Carbohydrates [i](#)

6 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	NAG	B	1	2,1	14,14,15	0.46	0	17,19,21	0.58	0
2	NAG	B	2	2	14,14,15	0.21	0	17,19,21	0.58	0
2	NAG	C	1	2,1	14,14,15	0.72	1 (7%)	17,19,21	1.86	3 (17%)
2	NAG	C	2	2	14,14,15	1.50	1 (7%)	17,19,21	0.79	1 (5%)
3	MAN	D	1	1,3	11,11,12	0.79	0	15,15,17	1.47	3 (20%)
3	MAN	D	2	3	11,11,12	1.18	1 (9%)	15,15,17	0.86	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	NAG	B	1	2,1	-	0/6/23/26	0/1/1/1
2	NAG	B	2	2	-	0/6/23/26	0/1/1/1
2	NAG	C	1	2,1	-	0/6/23/26	0/1/1/1
2	NAG	C	2	2	-	3/6/23/26	0/1/1/1
3	MAN	D	1	1,3	-	0/2/19/22	0/1/1/1
3	MAN	D	2	3	-	2/2/19/22	0/1/1/1

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	C	2	NAG	O5-C1	-5.37	1.35	1.43
3	D	2	MAN	C2-C3	2.55	1.56	1.52
2	C	1	NAG	O5-C1	2.46	1.47	1.43

All (7) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	C	1	NAG	O4-C4-C5	5.36	122.61	109.30
3	D	1	MAN	O2-C2-C3	-3.32	103.49	110.14
2	C	1	NAG	C1-O5-C5	3.24	116.58	112.19
3	D	1	MAN	C1-O5-C5	3.07	116.35	112.19
2	C	1	NAG	C4-C3-C2	-2.46	107.41	111.02
3	D	1	MAN	O5-C1-C2	2.44	114.54	110.77
2	C	2	NAG	C3-C4-C5	2.27	114.29	110.24

There are no chirality outliers.

All (5) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
3	D	2	MAN	C4-C5-C6-O6
2	C	2	NAG	C8-C7-N2-C2
2	C	2	NAG	O7-C7-N2-C2
3	D	2	MAN	O5-C5-C6-O6
2	C	2	NAG	O5-C5-C6-O6


There are no ring outliers.


1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	B	2	NAG	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.







5.6 Ligand geometry [i](#)

Of 27 ligands modelled in this entry, 11 are monoatomic - leaving 16 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
7	MAN	A	312	1	11,11,12	0.74	0	15,15,17	1.33	2 (13%)
7	MAN	A	307	1	11,11,12	1.31	2 (18%)	15,15,17	0.87	1 (6%)
7	MAN	A	303	1	11,11,12	0.74	0	15,15,17	1.21	2 (13%)
7	MAN	A	302	1	11,11,12	0.90	1 (9%)	15,15,17	0.93	1 (6%)
7	MAN	A	305[A]	1	11,11,12	1.23	2 (18%)	15,15,17	1.01	1 (6%)
7	MAN	A	313[A]	1	11,11,12	1.23	1 (9%)	15,15,17	1.10	0
7	MAN	A	313[B]	-	11,11,12	1.06	1 (9%)	15,15,17	1.25	2 (13%)

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
7	MAN	A	305[B]	1	11,11,12	2.83	4 (36%)	15,15,17	2.28	6 (40%)
7	MAN	A	306	1	11,11,12	1.30	2 (18%)	15,15,17	2.09	4 (26%)
7	MAN	A	311	1	11,11,12	1.28	1 (9%)	15,15,17	1.58	1 (6%)
6	NAG	A	301	1	14,14,15	0.33	0	17,19,21	0.56	0
7	MAN	A	310	1	11,11,12	1.13	2 (18%)	15,15,17	1.55	3 (20%)
7	MAN	A	308	1	11,11,12	0.77	0	15,15,17	1.26	2 (13%)
5	HEM	A	300	4,1	41,50,50	1.52	6 (14%)	45,82,82	1.53	8 (17%)
7	MAN	A	304	1	11,11,12	0.65	0	15,15,17	0.82	0
7	MAN	A	309	1	11,11,12	0.71	0	15,15,17	1.16	2 (13%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
7	MAN	A	312	1	-	2/2/19/22	0/1/1/1
7	MAN	A	307	1	-	0/2/19/22	0/1/1/1
7	MAN	A	303	1	-	0/2/19/22	0/1/1/1
7	MAN	A	302	1	-	0/2/19/22	0/1/1/1
7	MAN	A	305[A]	1	-	2/2/19/22	0/1/1/1
7	MAN	A	313[A]	1	-	1/2/19/22	0/1/1/1
7	MAN	A	313[B]	-	-	0/2/19/22	0/1/1/1
7	MAN	A	305[B]	1	-	1/2/19/22	0/1/1/1
7	MAN	A	306	1	-	0/2/19/22	0/1/1/1
7	MAN	A	311	1	-	2/2/19/22	0/1/1/1
6	NAG	A	301	1	-	0/6/23/26	0/1/1/1
7	MAN	A	310	1	-	0/2/19/22	0/1/1/1
7	MAN	A	308	1	-	2/2/19/22	0/1/1/1
5	HEM	A	300	4,1	-	3/12/54/54	-
7	MAN	A	304	1	-	0/2/19/22	0/1/1/1
7	MAN	A	309	1	-	0/2/19/22	0/1/1/1

All (22) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
7	A	305[B]	MAN	O5-C5	5.63	1.54	1.43
7	A	305[B]	MAN	C2-C3	4.96	1.59	1.52
5	A	300	HEM	C3C-C2C	-3.85	1.35	1.40

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
7	A	305[B]	MAN	C1-C2	3.65	1.60	1.52
7	A	311	MAN	O5-C5	3.62	1.50	1.43
5	A	300	HEM	C3C-CAC	3.54	1.55	1.47
7	A	305[B]	MAN	O5-C1	3.47	1.49	1.43
7	A	306	MAN	C1-C2	3.35	1.59	1.52
7	A	305[A]	MAN	O5-C1	-3.23	1.38	1.43
7	A	313[A]	MAN	O5-C1	-3.01	1.38	1.43
7	A	307	MAN	O5-C1	-2.89	1.39	1.43
5	A	300	HEM	CAB-C3B	2.80	1.55	1.47
5	A	300	HEM	FE-NB	2.79	2.10	1.96
5	A	300	HEM	CAA-C2A	2.67	1.56	1.52
7	A	306	MAN	C2-C3	2.64	1.56	1.52
7	A	302	MAN	O5-C1	-2.46	1.39	1.43
5	A	300	HEM	CMB-C2B	2.24	1.55	1.50
7	A	313[B]	MAN	C2-C3	2.23	1.55	1.52
7	A	305[A]	MAN	C1-C2	2.23	1.57	1.52
7	A	310	MAN	O5-C1	-2.09	1.40	1.43
7	A	307	MAN	C2-C3	2.07	1.55	1.52
7	A	310	MAN	C4-C5	2.05	1.57	1.53

All (35) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
7	A	311	MAN	C1-O5-C5	4.61	118.43	112.19
7	A	305[B]	MAN	C1-C2-C3	4.49	115.19	109.67
7	A	306	MAN	O5-C1-C2	4.38	117.54	110.77
7	A	306	MAN	C1-C2-C3	4.09	114.69	109.67
5	A	300	HEM	C4B-CHC-C1C	3.85	127.63	122.56
7	A	305[B]	MAN	O5-C5-C6	3.76	113.09	107.20
7	A	310	MAN	C1-O5-C5	3.70	117.20	112.19
7	A	306	MAN	C1-O5-C5	3.61	117.08	112.19
7	A	312	MAN	C1-O5-C5	3.49	116.92	112.19
7	A	305[B]	MAN	C2-C3-C4	3.23	116.48	110.89
7	A	305[B]	MAN	C1-O5-C5	3.16	116.47	112.19
7	A	309	MAN	C1-O5-C5	3.16	116.47	112.19
7	A	303	MAN	C1-O5-C5	3.10	116.39	112.19
7	A	310	MAN	C3-C4-C5	3.05	115.69	110.24
5	A	300	HEM	CMA-C3A-C4A	-3.04	123.80	128.46
5	A	300	HEM	C4D-ND-C1D	3.03	108.20	105.07
5	A	300	HEM	C1B-NB-C4B	3.02	108.19	105.07
7	A	308	MAN	C1-O5-C5	2.78	115.95	112.19
7	A	305[A]	MAN	O2-C2-C3	-2.58	104.97	110.14

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
7	A	305[B]	MAN	C6-C5-C4	2.53	118.92	113.00
5	A	300	HEM	C2D-C1D-ND	-2.37	107.04	109.88
7	A	305[B]	MAN	O2-C2-C3	-2.35	105.43	110.14
7	A	302	MAN	O5-C1-C2	2.31	114.34	110.77
7	A	303	MAN	O2-C2-C3	-2.29	105.55	110.14
7	A	310	MAN	O2-C2-C3	-2.28	105.56	110.14
7	A	306	MAN	C3-C4-C5	-2.25	106.23	110.24
7	A	309	MAN	O2-C2-C3	-2.24	105.64	110.14
5	A	300	HEM	CAA-CBA-CGA	-2.17	107.68	113.76
7	A	308	MAN	O2-C2-C3	-2.17	105.80	110.14
7	A	312	MAN	O3-C3-C2	2.09	114.00	109.99
7	A	313[B]	MAN	C1-O5-C5	2.07	115.00	112.19
7	A	307	MAN	O2-C2-C3	-2.03	106.06	110.14
7	A	313[B]	MAN	C3-C4-C5	-2.03	106.61	110.24
5	A	300	HEM	C1D-C2D-C3D	2.02	109.08	106.96
5	A	300	HEM	O2D-CGD-CBD	2.01	120.50	114.03

There are no chirality outliers.

All (13) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
7	A	305[A]	MAN	O5-C5-C6-O6
7	A	311	MAN	O5-C5-C6-O6
7	A	305[A]	MAN	C4-C5-C6-O6
7	A	311	MAN	C4-C5-C6-O6
7	A	312	MAN	C4-C5-C6-O6
7	A	308	MAN	O5-C5-C6-O6
7	A	313[A]	MAN	O5-C5-C6-O6
7	A	308	MAN	C4-C5-C6-O6
7	A	312	MAN	O5-C5-C6-O6
5	A	300	HEM	C2A-CAA-CBA-CGA
5	A	300	HEM	CAA-CBA-CGA-O1A
5	A	300	HEM	CAA-CBA-CGA-O2A
7	A	305[B]	MAN	O5-C5-C6-O6

There are no ring outliers.

3 monomers are involved in 7 short contacts:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
7	A	305[B]	MAN	2	0
7	A	311	MAN	2	0

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Clashes	Symm-Clashes
5	A	300	HEM	3	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	298/299 (99%)	-0.07	5 (1%) 70 74	11, 17, 33, 57	0

All (5) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	275	THR	3.3
1	A	274	SER	2.9
1	A	273	ALA	2.5
1	A	120	ASN	2.2
1	A	122	ASN	2.2

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
1	PCA	A	0	8/9	0.96	0.07	15,17,18,20	0

6.3 Carbohydrates [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.


Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	NAG	C	2	14/15	0.68	0.32	45,59,65,70	0

Continued on next page...

Continued from previous page...




Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
3	MAN	D	2	11/12	0.83	0.22	38,47,54,56	0
3	MAN	D	1	11/12	0.92	0.16	34,37,48,52	0
2	NAG	C	1	14/15	0.93	0.10	12,17,24,50	0
2	NAG	B	2	14/15	0.93	0.15	18,23,37,49	0
2	NAG	B	1	14/15	0.95	0.07	14,19,23,25	0


The following is a graphical depiction of the model fit to experimental electron density for oligosaccharide. Each fit is shown from different orientation to approximate a three-dimensional view.



Electron density around Chain C:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)





6.4 Ligands ⓘ

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.


Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
7	MAN	A	311	11/12	0.63	0.32	64,72,76,80	0
7	MAN	A	310	11/12	0.67	0.48	75,85,89,97	0
7	MAN	A	313[A]	11/12	0.75	0.21	26,34,37,37	11
7	MAN	A	313[B]	11/12	0.75	0.21	29,31,39,42	11
7	MAN	A	305[A]	11/12	0.76	0.18	17,21,28,29	11
7	MAN	A	305[B]	11/12	0.76	0.18	19,25,29,30	11
7	MAN	A	306	11/12	0.81	0.27	56,58,63,67	0
7	MAN	A	308	11/12	0.82	0.21	30,36,46,51	0
7	MAN	A	307	11/12	0.87	0.12	36,43,54,56	0

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
7	MAN	A	312	11/12	0.88	0.22	37,47,54,56	0
6	NAG	A	301	14/15	0.90	0.17	23,29,34,35	0
7	MAN	A	309	11/12	0.93	0.16	26,33,39,41	0
7	MAN	A	302	11/12	0.94	0.10	22,27,31,34	0
7	MAN	A	303	11/12	0.96	0.06	16,19,24,24	0
8	IOD	A	318	1/1	0.96	0.05	35,35,35,35	1
5	HEM	A	300	43/43	0.98	0.12	8,11,13,15	0
8	IOD	A	314	1/1	0.98	0.04	35,35,35,35	1
8	IOD	A	315	1/1	0.98	0.04	27,27,27,27	1
7	MAN	A	304	11/12	0.98	0.06	13,15,16,18	0
8	IOD	A	319	1/1	0.98	0.05	33,33,33,33	1
8	IOD	A	320	1/1	0.98	0.05	37,37,37,37	1
8	IOD	A	317	1/1	0.99	0.03	28,28,28,28	1
8	IOD	A	321	1/1	0.99	0.06	20,20,20,20	0
8	IOD	A	323	1/1	0.99	0.03	36,36,36,36	1
4	MN	A	299	1/1	1.00	0.05	21,21,21,21	0
8	IOD	A	322	1/1	1.00	0.02	27,27,27,27	1
8	IOD	A	316	1/1	1.00	0.02	32,32,32,32	1

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.



6.5 Other polymers [i](#)

There are no such residues in this entry.