



# Full wwPDB X-ray Structure Validation Report ⓘ

May 13, 2020 – 05:30 am BST

PDB ID : 6RTS  
Title : Piperideine-6-carboxylate dehydrogenase from *Streptomyces clavuligerus* complexed with NAD<sup>+</sup>  
Authors : Hasse, D.; Huelsemann, J.; Carlsson, G.; Andersson, I.  
Deposited on : 2019-05-26  
Resolution : 2.25 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.11  
buster-report : 1.1.7 (2018)  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.11


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.25 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	1377 (2.26-2.26)
Clashscore	141614	1487 (2.26-2.26)
Ramachandran outliers	138981	1449 (2.26-2.26)
Sidechain outliers	138945	1450 (2.26-2.26)
RSRZ outliers	127900	1356 (2.26-2.26)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	512	 95%
1	B	512	 2% 96%

## 2 Entry composition [i](#)

There are 6 unique types of molecules in this entry. The entry contains 8260 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Semialdehyde dehydrogenase Pcd.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	509	Total	C	N	O	S	0	1	0
			3777	2356	681	725	15			
1	B	508	Total	C	N	O	S	0	1	0
			3770	2352	680	723	15			

There are 4 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	140	THR	SER	engineered mutation	UNP O85725
A	503	THR	ALA	engineered mutation	UNP O85725
B	140	THR	SER	engineered mutation	UNP O85725
B	503	THR	ALA	engineered mutation	UNP O85725

- Molecule 2 is NICOTINAMIDE-ADENINE-DINUCLEOTIDE (three-letter code: NAD) (formula: C<sub>21</sub>H<sub>27</sub>N<sub>7</sub>O<sub>14</sub>P<sub>2</sub>) (labeled as "Ligand of Interest" by author).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total	C	N	O	P	0	0
			44	21	7	14	2		
2	B	1	Total	C	N	O	P	0	0
			44	21	7	14	2		

- Molecule 3 is SULFATE ION (three-letter code: SO4) (formula: O<sub>4</sub>S).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	O	S	0	0
			5	4	1		
3	A	1	Total	O	S	0	0
			5	4	1		
3	A	1	Total	O	S	0	0
			5	4	1		
3	A	1	Total	O	S	0	0
			5	4	1		
3	A	1	Total	O	S	0	0
			5	4	1		
3	B	1	Total	O	S	0	0
			5	4	1		
3	B	1	Total	O	S	0	0
			5	4	1		
3	B	1	Total	O	S	0	0
			5	4	1		

- Molecule 4 is GLYCEROL (three-letter code: GOL) (formula: C<sub>3</sub>H<sub>8</sub>O<sub>3</sub>).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	C	O	0	0
			6	3	3		
4	A	1	Total	C	O	0	0
			6	3	3		
4	A	1	Total	C	O	0	0
			6	3	3		
4	B	1	Total	C	O	0	0
			6	3	3		
4	B	1	Total	C	O	0	0
			6	3	3		

- Molecule 5 is ACETATE ION (three-letter code: ACT) (formula:  $C_2H_3O_2$ ).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	A	1	Total C O 4 2 2	0	0
5	B	1	Total C O 4 2 2	0	0
5	B	1	Total C O 4 2 2	0	0

- Molecule 6 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
6	A	268	Total O 268 268	0	0
6	B	275	Total O 275 275	0	0

### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $\text{RSRZ} > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Semialdehyde dehydrogenase Pcd

Chain A:  95%



- Molecule 1: Semialdehyde dehydrogenase Pcd

Chain B:  96%



## 4 Data and refinement statistics

Property	Value	Source
Space group	C 2 2 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	111.04Å 131.24Å 157.71Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	45.40 – 2.25 45.40 – 2.25	Depositor EDS
% Data completeness (in resolution range)	99.4 (45.40-2.25) 99.4 (45.40-2.25)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.09 (at 2.24Å)	Xtriage
Refinement program	REFMAC 5.8.0107	Depositor
R, $R_{free}$	0.166 , 0.205 0.173 , 0.205	Depositor DCC
$R_{free}$ test set	2727 reflections (5.00%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	21.8	Xtriage
Anisotropy	0.281	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.38 , 37.3	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.47$ , $\langle L^2 \rangle = 0.30$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.95	EDS
Total number of atoms	8260	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	23.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.20% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



## 5 Model quality

### 5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: GOL, SO4, NAD, ACT

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.57	0/3845	0.77	0/5240
1	B	0.55	0/3838	0.73	0/5230
All	All	0.56	0/7683	0.75	0/10470

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3777	0	3759	7	0
1	B	3770	0	3752	5	0
2	A	44	0	26	0	0
2	B	44	0	26	0	0
3	A	25	0	0	0	0
3	B	15	0	0	0	0
4	A	18	0	24	1	0
4	B	12	0	16	0	0
5	A	4	0	3	1	0
5	B	8	0	6	0	0
6	A	268	0	0	1	0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
6	B	275	0	0	1	0
All	All	8260	0	7612	13	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 1.

All (13) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:218:GLN:HB2	6:B:705:HOH:O	2.05	0.56
1:A:99:ASP:OD1	1:A:109:ARG:NH1	2.39	0.54
1:B:299[B]:CYS:SG	1:B:427:LEU:HD22	2.48	0.54
1:A:107:LYS:NZ	1:A:115:GLU:OE1	2.36	0.53
1:A:243:GLY:O	1:A:266:LEU:HA	2.15	0.47
1:A:300:THR:OG1	5:A:610:ACT:H1	2.15	0.45
1:B:243:GLY:O	1:B:266:LEU:HA	2.17	0.45
1:B:132:GLN:O	1:B:133:LEU:C	2.55	0.44
1:A:271:ALA:HA	1:A:304:ARG:O	2.19	0.43
1:A:299[A]:CYS:SG	1:A:427:LEU:HD22	2.58	0.43
1:A:132:GLN:O	1:A:133:LEU:C	2.58	0.42
4:A:607:GOL:H2	6:A:737:HOH:O	2.19	0.41
1:B:35:LEU:O	1:B:49:LEU:N	2.52	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles ⓘ

### 5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	508/512 (99%)	490 (96%)	17 (3%)	1 (0%)	47	55
1	B	507/512 (99%)	491 (97%)	15 (3%)	1 (0%)	47	55
All	All	1015/1024 (99%)	981 (97%)	32 (3%)	2 (0%)	47	55

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	480	SER
1	B	480	SER

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	381/383 (100%)	374 (98%)	7 (2%)	59	68
1	B	380/383 (99%)	375 (99%)	5 (1%)	69	79
All	All	761/766 (99%)	749 (98%)	12 (2%)	62	73

All (12) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	7	SER
1	A	118	GLU
1	A	138	MET
1	A	187	LYS
1	A	238	LEU
1	A	294	THR
1	A	334	ASP
1	B	138	MET
1	B	140	THR
1	B	218	GLN
1	B	294	THR
1	B	443	LEU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 5.6 Ligand geometry [i](#)

18 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
3	SO4	A	603	-	4,4,4	0.43	0	6,6,6	0.09	0
3	SO4	A	604	-	4,4,4	0.33	0	6,6,6	0.24	0
4	GOL	B	605	-	5,5,5	0.35	0	5,5,5	0.61	0
3	SO4	B	604	-	4,4,4	0.33	0	6,6,6	0.19	0
4	GOL	A	609	-	5,5,5	0.38	0	5,5,5	0.31	0
3	SO4	A	606	-	4,4,4	0.34	0	6,6,6	0.35	0
2	NAD	B	601	-	42,48,48	0.99	2 (4%)	50,73,73	1.20	3 (6%)
4	GOL	B	606	-	5,5,5	0.35	0	5,5,5	0.62	0
3	SO4	A	602	-	4,4,4	0.25	0	6,6,6	0.24	0
3	SO4	A	605	-	4,4,4	0.36	0	6,6,6	0.51	0
4	GOL	A	607	-	5,5,5	0.81	0	5,5,5	1.83	1 (20%)
4	GOL	A	608	-	5,5,5	0.42	0	5,5,5	0.76	0
5	ACT	B	608	-	1,3,3	1.77	0	0,3,3	0.00	-
5	ACT	A	610	-	1,3,3	3.02	1 (100%)	0,3,3	0.00	-
2	NAD	A	601	-	42,48,48	0.90	1 (2%)	50,73,73	1.38	7 (14%)
3	SO4	B	602	-	4,4,4	0.23	0	6,6,6	0.38	0
5	ACT	B	607	-	1,3,3	2.87	1 (100%)	0,3,3	0.00	-
3	SO4	B	603	-	4,4,4	0.39	0	6,6,6	0.63	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the

Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns.  
'-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	GOL	B	605	-	-	0/4/4/4	-
4	GOL	A	609	-	-	0/4/4/4	-
2	NAD	B	601	-	-	1/26/62/62	0/5/5/5
4	GOL	B	606	-	-	2/4/4/4	-
4	GOL	A	607	-	-	3/4/4/4	-
4	GOL	A	608	-	-	0/4/4/4	-
2	NAD	A	601	-	-	1/26/62/62	0/5/5/5

All (5) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
5	A	610	ACT	CH3-C	3.02	1.52	1.48
5	B	607	ACT	CH3-C	2.87	1.52	1.48
2	B	601	NAD	O4D-C1D	2.79	1.45	1.41
2	B	601	NAD	C5A-C4A	2.51	1.47	1.40
2	A	601	NAD	C5A-C4A	2.44	1.47	1.40

All (11) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	601	NAD	C3N-C7N-N7N	4.26	122.87	117.75
2	B	601	NAD	C3N-C7N-N7N	4.03	122.58	117.75
2	A	601	NAD	N3A-C2A-N1A	-3.84	122.67	128.68
2	A	601	NAD	C1B-N9A-C4A	-3.23	120.96	126.64
2	B	601	NAD	N3A-C2A-N1A	-3.14	123.76	128.68
4	A	607	GOL	O2-C2-C1	3.06	122.62	109.12
2	A	601	NAD	O7N-C7N-C3N	-2.95	116.10	119.63
2	A	601	NAD	C4A-C5A-N7A	-2.38	106.92	109.40
2	A	601	NAD	PN-O3-PA	-2.37	124.69	132.83
2	A	601	NAD	C2A-N1A-C6A	2.36	122.78	118.75
2	B	601	NAD	O7N-C7N-C3N	-2.03	117.20	119.63

There are no chirality outliers.

All (7) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	B	606	GOL	O1-C1-C2-C3
4	A	607	GOL	C1-C2-C3-O3
4	A	607	GOL	O1-C1-C2-O2

*Continued on next page...*

*Continued from previous page...*


Mol	Chain	Res	Type	Atoms
4	B	606	GOL	O1-C1-C2-O2
4	A	607	GOL	O2-C2-C3-O3
2	B	601	NAD	C4D-C5D-O5D-PN
2	A	601	NAD	C4D-C5D-O5D-PN


There are no ring outliers.

2 monomers are involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	A	607	GOL	1	0
5	A	610	ACT	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.



## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	509/512 (99%)	-0.44	2 (0%) 92 93	12, 20, 38, 65	0
1	B	508/512 (99%)	-0.35	9 (1%) 68 71	14, 20, 40, 90	0
All	All	1017/1024 (99%)	-0.39	11 (1%) 80 82	12, 20, 39, 90	0

All (11) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	508	VAL	4.3
1	B	505	ALA	3.9
1	B	507	GLY	3.9
1	B	506	GLN	3.6
1	A	246	ARG	3.3
1	B	511	SER	2.8
1	A	249	ARG	2.6
1	B	33	ASP	2.5
1	B	4	ALA	2.4
1	B	249	ARG	2.4
1	B	5	ALA	2.1

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 6.4 Ligands ⓘ




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
5	ACT	A	610	4/4	0.79	0.20	32,36,37,41	0
4	GOL	A	609	6/6	0.80	0.25	35,38,39,40	6
5	ACT	B	607	4/4	0.81	0.21	37,40,40,42	0
4	GOL	A	607	6/6	0.82	0.19	28,32,35,37	0
5	ACT	B	608	4/4	0.83	0.17	50,53,54,54	0
4	GOL	A	608	6/6	0.86	0.19	22,26,26,27	6
3	SO4	A	606	5/5	0.88	0.22	38,42,43,43	5
4	GOL	B	605	6/6	0.91	0.15	27,29,31,32	6
3	SO4	B	603	5/5	0.91	0.15	41,42,45,47	5
4	GOL	B	606	6/6	0.93	0.16	33,35,37,37	0
2	NAD	A	601	44/44	0.95	0.12	26,31,37,39	0
2	NAD	B	601	44/44	0.95	0.13	22,28,34,35	0
3	SO4	B	604	5/5	0.95	0.14	47,50,53,53	5
3	SO4	A	605	5/5	0.96	0.12	35,40,41,42	0
3	SO4	A	603	5/5	0.97	0.19	37,38,39,40	0
3	SO4	A	604	5/5	0.97	0.11	53,54,55,57	0
3	SO4	A	602	5/5	0.99	0.09	30,30,30,31	0
3	SO4	B	602	5/5	0.99	0.13	32,33,34,34	0




The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

**Electron density around NAD A 601:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)

**Electron density around NAD B 601:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)



## 6.5 Other polymers

There are no such residues in this entry.