



Full wwPDB X-ray Structure Validation Report ⓘ

Aug 10, 2020 – 03:12 AM BST

PDB ID : 6RV9
Title : Crystal Structure of Glucuronoyl Esterase from *Cerrena unicolor* inactive S270A variant in complex with the aldouronic acid XUXXR
Authors : Ernst, H.A.; Mosbech, C.; Langkilde, A.; Westh, P.; Meyer, A.; Agger, J.W.; Larsen, S.
Deposited on : 2019-05-31
Resolution : 1.64 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.13.1
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.13.1


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.64 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	3122 (1.66-1.62)
Clashscore	141614	3268 (1.66-1.62)
Ramachandran outliers	138981	3215 (1.66-1.62)
Sidechain outliers	138945	3215 (1.66-1.62)
RSRZ outliers	127900	3079 (1.66-1.62)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	401	<div> <div>2%</div> <div> <div></div> <div>92%</div> <div>5%</div> </div> </div>
1	B	401	<div> <div>12%</div> <div> <div></div> <div>85%</div> <div>10%</div> <div>5%</div> </div> </div>
2	C	5	<div> <div></div> <div> <div></div> <div>80%</div> <div>20%</div> </div> </div>

2 Entry composition [i](#)

There are 5 unique types of molecules in this entry. The entry contains 6398 atoms, of which 51 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called 4-O-methyl-glucuronoyl methylesterase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	380	Total	C	N	O	S	0	2	0
			2872	1814	492	551	15			
1	B	380	Total	C	N	O	S	0	1	0
			2866	1810	491	550	15			

There are 44 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	73	GLU	-	expression tag	UNP A0A0A7EQR3
A	74	ALA	-	expression tag	UNP A0A0A7EQR3
A	75	GLU	-	expression tag	UNP A0A0A7EQR3
A	76	ALA	-	expression tag	UNP A0A0A7EQR3
A	77	GLU	-	expression tag	UNP A0A0A7EQR3
A	78	PHE	-	expression tag	UNP A0A0A7EQR3
A	270	ALA	SER	engineered mutation	UNP A0A0A7EQR3
A	459	GLU	-	expression tag	UNP A0A0A7EQR3
A	460	ASN	-	expression tag	UNP A0A0A7EQR3
A	461	LEU	-	expression tag	UNP A0A0A7EQR3
A	462	TYR	-	expression tag	UNP A0A0A7EQR3
A	463	PHE	-	expression tag	UNP A0A0A7EQR3
A	464	GLN	-	expression tag	UNP A0A0A7EQR3
A	465	GLY	-	expression tag	UNP A0A0A7EQR3
A	466	VAL	-	expression tag	UNP A0A0A7EQR3
A	467	ASP	-	expression tag	UNP A0A0A7EQR3
A	468	HIS	-	expression tag	UNP A0A0A7EQR3
A	469	HIS	-	expression tag	UNP A0A0A7EQR3
A	470	HIS	-	expression tag	UNP A0A0A7EQR3
A	471	HIS	-	expression tag	UNP A0A0A7EQR3
A	472	HIS	-	expression tag	UNP A0A0A7EQR3
A	473	HIS	-	expression tag	UNP A0A0A7EQR3
B	73	GLU	-	expression tag	UNP A0A0A7EQR3
B	74	ALA	-	expression tag	UNP A0A0A7EQR3
B	75	GLU	-	expression tag	UNP A0A0A7EQR3

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
B	76	ALA	-	expression tag	UNP A0A0A7EQR3
B	77	GLU	-	expression tag	UNP A0A0A7EQR3
B	78	PHE	-	expression tag	UNP A0A0A7EQR3
B	270	ALA	SER	engineered mutation	UNP A0A0A7EQR3
B	459	GLU	-	expression tag	UNP A0A0A7EQR3
B	460	ASN	-	expression tag	UNP A0A0A7EQR3
B	461	LEU	-	expression tag	UNP A0A0A7EQR3
B	462	TYR	-	expression tag	UNP A0A0A7EQR3
B	463	PHE	-	expression tag	UNP A0A0A7EQR3
B	464	GLN	-	expression tag	UNP A0A0A7EQR3
B	465	GLY	-	expression tag	UNP A0A0A7EQR3
B	466	VAL	-	expression tag	UNP A0A0A7EQR3
B	467	ASP	-	expression tag	UNP A0A0A7EQR3
B	468	HIS	-	expression tag	UNP A0A0A7EQR3
B	469	HIS	-	expression tag	UNP A0A0A7EQR3
B	470	HIS	-	expression tag	UNP A0A0A7EQR3
B	471	HIS	-	expression tag	UNP A0A0A7EQR3
B	472	HIS	-	expression tag	UNP A0A0A7EQR3
B	473	HIS	-	expression tag	UNP A0A0A7EQR3

- Molecule 2 is an oligosaccharide called 4-O-methyl-alpha-D-glucopyranuronic acid-(1-2)-[beta-D-xylopyranose-(1-4)]beta-D-xylopyranose-(1-4)-beta-D-xylopyranose-(1-4)-Xylitol.



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
2	C	5	Total	C	H	O	0	0	0
			95	27	45	23			

- Molecule 3 is 2-acetamido-2-deoxy-beta-D-glucopyranose (three-letter code: NAG) (formula: C₈H₁₅NO₆).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
3	A	1	Total	C	N	O	0	0
			14	8	1	5		
3	B	1	Total	C	N	O	0	0
			14	8	1	5		

- Molecule 4 is 1,2-ETHANEDIOL (three-letter code: EDO) (formula: $C_2H_6O_2$).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
4	A	1	Total	C	H	O	0	0
			10	2	6	2		

- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	302	Total 302	O 302	0	0
5	B	225	Total 225	O 225	0	0

4 Data and refinement statistics

Property	Value	Source
Space group	P 41 21 2	Depositor
Cell constants a, b, c, α , β , γ	84.48 Å 84.48 Å 261.73 Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	44.50 – 1.64 44.50 – 1.64	Depositor EDS
% Data completeness (in resolution range)	99.5 (44.50-1.64) 99.6 (44.50-1.64)	Depositor EDS
R_{merge}	0.07	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.89 (at 1.64 Å)	Xtriage
Refinement program	PHENIX 1.13 _2998	Depositor
R, R_{free}	0.186 , 0.210 0.185 , 0.209	Depositor DCC
R_{free} test set	5754 reflections (4.95%)	wwPDB-VP
Wilson B-factor (Å ²)	23.3	Xtriage
Anisotropy	0.023	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.35 , 42.7	EDS
L-test for twinning ²	$\langle L \rangle = 0.46$, $\langle L^2 \rangle = 0.29$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	6398	wwPDB-VP
Average B, all atoms (Å ²)	31.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.14% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: XYP, GCV, XYL, NAG, EDO

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.36	0/2956	0.56	0/4045
1	B	0.33	0/2947	0.54	0/4033
All	All	0.35	0/5903	0.55	0/8078

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2872	0	2760	7	0
1	B	2866	0	2752	23	0
2	C	50	45	20	1	0
3	A	14	0	13	0	0
3	B	14	0	13	0	0
4	A	4	6	6	0	0
5	A	302	0	0	1	0
5	B	225	0	0	1	0
All	All	6347	51	5564	28	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (28) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:174:PRO:HG2	1:B:183:TRP:CE3	2.36	0.60
1:B:252:GLU:HB3	5:B:627:HOH:O	2.01	0.60
1:B:270:ALA:HB1	2:C:4:GCV:C6	2.32	0.59
1:B:142:ALA:HA	1:B:154:ALA:O	2.10	0.52
1:A:429:THR:HG21	1:B:338:LYS:NZ	2.24	0.51
1:B:138:PRO:HD2	1:B:159:LEU:HA	1.92	0.51
1:B:179:PRO:HG2	1:B:182:GLY:O	2.10	0.51
1:B:186:ILE:HD12	1:B:196:ILE:HD12	1.93	0.50
1:B:190:GLU:HG3	1:B:207:ASN:OD1	2.13	0.48
1:B:153:LEU:O	1:B:168:SER:HA	2.13	0.48
1:B:144:PHE:CZ	1:B:151:GLY:HA3	2.49	0.47
1:B:248:ILE:HG21	1:B:285:ARG:HD2	1.95	0.47
1:B:153:LEU:O	1:B:169:PRO:HD2	2.14	0.47
1:A:385:TRP:CD2	1:A:394:HIS:HB2	2.50	0.47
1:A:314:GLN:NE2	5:A:603:HOH:O	2.48	0.46
1:B:165:ILE:HD12	1:B:223:LEU:HB3	1.97	0.46
1:B:174:PRO:HB2	1:B:183:TRP:CH2	2.51	0.45
1:A:178:PRO:HB3	1:A:183:TRP:NE1	2.32	0.44
1:B:242:TRP:O	1:B:246:ARG:HG2	2.17	0.44
1:B:140:VAL:HA	1:B:156:THR:O	2.19	0.42
1:A:177:THR:HB	1:B:229:GLY:HA2	2.01	0.42
1:B:174:PRO:HB2	1:B:183:TRP:CZ3	2.54	0.42
1:B:146:LYS:HE2	1:B:173:TYR:CD1	2.55	0.42
1:A:373:SER:O	1:A:377:CYS:HB2	2.20	0.42
1:B:137:PRO:HG3	1:B:242:TRP:HZ2	1.85	0.42
1:A:304:LEU:CD2	1:A:449:TRP:CD2	3.04	0.40
1:B:187:ILE:O	1:B:266:VAL:HA	2.21	0.40
1:B:385:TRP:CD2	1:B:394:HIS:HB2	2.56	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	380/401 (95%)	365 (96%)	15 (4%)	0	100	100
1	B	379/401 (94%)	366 (97%)	13 (3%)	0	100	100
All	All	759/802 (95%)	731 (96%)	28 (4%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	304/320 (95%)	303 (100%)	1 (0%)	92	87
1	B	303/320 (95%)	300 (99%)	3 (1%)	76	59
All	All	607/640 (95%)	603 (99%)	4 (1%)	84	71

All (4) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	370	SER
1	B	256	THR
1	B	370	SER
1	B	433	PHE

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

5 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z > 2$	Counts	RMSZ	# $ Z > 2$
2	XYL	C	1	2	9,9,9	0.83	0	11,11,11	1.02	1 (9%)
2	XYP	C	2	2	9,9,10	1.90	3 (33%)	10,12,14	1.51	3 (30%)
2	XYP	C	3	2	9,9,10	1.86	3 (33%)	10,12,14	1.48	1 (10%)
2	GCV	C	4	2	10,13,14	2.16	3 (30%)	11,18,20	1.80	4 (36%)
2	XYP	C	5	2	9,9,10	1.88	2 (22%)	10,12,14	1.24	1 (10%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	XYL	C	1	2	-	2/12/12/12	-
2	XYP	C	2	2	-	-	0/1/1/1
2	XYP	C	3	2	-	-	0/1/1/1
2	GCV	C	4	2	-	2/2/23/26	0/1/1/1
2	XYP	C	5	2	-	-	0/1/1/1

All (11) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	C	4	GCV	O5-C5	5.07	1.49	1.43
2	C	5	XYP	O5-C1	3.99	1.50	1.42
2	C	3	XYP	O5-C1	3.95	1.50	1.42
2	C	2	XYP	O5-C1	3.93	1.50	1.42

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	C	4	GCV	O5-C1	3.66	1.49	1.43
2	C	5	XYP	O5-C5	2.72	1.48	1.42
2	C	3	XYP	O5-C5	2.68	1.48	1.42
2	C	2	XYP	O5-C5	2.53	1.47	1.42
2	C	4	GCV	C2-C3	-2.46	1.48	1.52
2	C	2	XYP	C2-C3	-2.34	1.49	1.52
2	C	3	XYP	C2-C3	-2.07	1.49	1.52

All (10) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	C	3	XYP	C5-C4-C3	4.22	114.86	109.67
2	C	4	GCV	O2-C2-C1	-3.10	102.81	109.15
2	C	4	GCV	O3-C3-C2	-3.07	104.11	109.99
2	C	2	XYP	C1-C2-C3	3.04	113.40	109.67
2	C	5	XYP	C5-C4-C3	2.98	113.33	109.67
2	C	4	GCV	C3-C4-C5	-2.90	106.56	110.28
2	C	1	XYL	C5-C4-C3	-2.42	107.17	112.41
2	C	2	XYP	C4-C3-C2	2.33	113.68	110.92
2	C	4	GCV	O2-C2-C3	2.10	114.35	110.14
2	C	2	XYP	O4-C4-C3	-2.05	106.03	110.14

There are no chirality outliers.

All (4) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
2	C	4	GCV	C3-C4-O4-C7
2	C	4	GCV	C5-C4-O4-C7
2	C	1	XYL	C1-C2-C3-C4
2	C	1	XYL	O2-C2-C3-C4

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	C	4	GCV	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.



5.6 Ligand geometry [i](#)

3 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	NAG	B	501	1	14,14,15	0.27	0	17,19,21	0.53	0
3	NAG	A	501	1	14,14,15	0.43	0	17,19,21	0.55	0
4	EDO	A	502	-	3,3,3	0.47	0	2,2,2	0.37	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns.

'-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	NAG	B	501	1	-	0/6/23/26	0/1/1/1
3	NAG	A	501	1	-	0/6/23/26	0/1/1/1
4	EDO	A	502	-	-	0/1/1/1	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	380/401 (94%)	-0.14	8 (2%) 63 64	18, 24, 39, 62	0
1	B	380/401 (94%)	0.57	50 (13%) 3 2	18, 32, 67, 84	0
All	All	760/802 (94%)	0.22	58 (7%) 13 12	18, 27, 61, 84	0

All (58) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	148	GLY	8.3
1	B	175	SER	7.6
1	A	79	GLY	5.9
1	B	147	SER	5.7
1	B	149	ASN	5.5
1	B	176	GLY	5.5
1	B	163	GLN	5.4
1	B	144	PHE	5.4
1	B	150	THR	4.5
1	A	80	ALA	4.3
1	B	177	THR	4.3
1	B	183	TRP	4.2
1	B	258	GLN	4.1
1	B	79	GLY	4.1
1	B	178	PRO	4.1
1	B	259	ILE	4.0
1	B	250	ALA	3.8
1	A	176	GLY	3.7
1	B	139	VAL	3.7
1	B	256	THR	3.6
1	B	154	ALA	3.6
1	B	146	LYS	3.6
1	B	174	PRO	3.5
1	B	80	ALA	3.5

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	165	ILE	3.4
1	B	143	SER	3.4
1	B	253	MET	3.4
1	B	145	SER	3.3
1	B	180	ALA	3.1
1	B	173	TYR	3.1
1	B	151	GLY	2.9
1	B	141	THR	2.8
1	B	152	THR	2.8
1	B	171	ILE	2.7
1	B	153	LEU	2.7
1	B	198	ALA	2.7
1	B	261	THR	2.7
1	B	458	THR	2.7
1	B	166	LYS	2.6
1	B	456	THR	2.6
1	B	162	SER	2.5
1	B	257	ALA	2.4
1	B	167	PHE	2.4
1	A	175	SER	2.4
1	B	140	VAL	2.4
1	A	144	PHE	2.3
1	B	170	THR	2.3
1	B	112	ALA	2.3
1	A	256	THR	2.3
1	B	164	THR	2.3
1	B	251	LEU	2.2
1	A	149	ASN	2.2
1	B	217	SER	2.2
1	B	254	THR	2.2
1	A	81	CYS	2.1
1	B	172	SER	2.1
1	B	155	ILE	2.1
1	B	181	ASN	2.0

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ


There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	XYL	C	1	10/10	0.89	0.27	36,48,60,62	0
2	XYP	C	5	9/10	0.91	0.18	35,42,48,50	0
2	XYP	C	2	9/10	0.94	0.14	25,33,43,49	0
2	XYP	C	3	9/10	0.95	0.10	24,30,41,41	0
2	GCV	C	4	13/14	0.95	0.07	23,28,33,33	0

The following is a graphical depiction of the model fit to experimental electron density for oligosaccharide. Each fit is shown from different orientation to approximate a three-dimensional view.



6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
3	NAG	B	501	14/15	0.93	0.09	24,29,34,36	0
4	EDO	A	502	4/4	0.93	0.12	21,35,44,44	0
3	NAG	A	501	14/15	0.95	0.07	22,27,31,31	0

6.5 Other polymers [i](#)

There are no such residues in this entry.