



# Full wwPDB X-ray Structure Validation Report ⓘ

Aug 22, 2020 – 07:17 AM BST

PDB ID : 1RWC  
Title : Crystal structure of Arthrobacter aurescens chondroitin AC lyase  
Authors : Lunin, V.V.; Li, Y.; Miyazono, H.; Kyogashima, M.; Bell, A.W.; Cygler, M.  
Deposited on : 2003-12-16  
Resolution : 1.90 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.13.1  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.13.1


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

## *X-RAY DIFFRACTION*

The reported resolution of this entry is 1.90 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	6207 (1.90-1.90)
Clashscore	141614	6847 (1.90-1.90)
Ramachandran outliers	138981	6760 (1.90-1.90)
Sidechain outliers	138945	6760 (1.90-1.90)
RSRZ outliers	127900	6082 (1.90-1.90)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	757	<div> <div>2%</div> <div> <div></div> <div>89%</div> <div>10%</div> </div> </div>
2	B	2	<div> <div></div> <div>100%</div> </div>

## 2 Entry composition [i](#)


There are 6 unique types of molecules in this entry. The entry contains 6510 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called chondroitin AC lyase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	754	Total	C	N	O	S	0	11	0
			5629	3535	987	1089	18			

- Molecule 2 is an oligosaccharide called 2,6-anhydro-3-deoxy-L-threo-hex-2-enonic acid-(1-3)-2-acetamido-2-deoxy-beta-D-glucopyranose.



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
2	B	2	Total	C	N	O	0	0	0
			26	14	1	11			

- Molecule 3 is SODIUM ION (three-letter code: NA) (formula: Na).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	1	Total	Na	0	0
			1	1		

- Molecule 4 is PHOSPHATE ION (three-letter code: PO4) (formula: O<sub>4</sub>P).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	O	P	0	0
			5	4	1		

- Molecule 5 is GLYCEROL (three-letter code: GOL) (formula: C<sub>3</sub>H<sub>8</sub>O<sub>3</sub>).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	A	1	Total	C	O	0	1
			12	6	6		


- Molecule 6 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	837	Total 837	O 837	0	0


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: chondroitin AC lyase



- Molecule 2: 2,6-anhydro-3-deoxy-L-threo-hex-2-enonic acid-(1-3)-2-acetamido-2-deoxy-beta-D-glucopyranose



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	57.68Å 86.44Å 81.29Å 90.00° 106.88° 90.00°	Depositor
Resolution (Å)	29.36 – 1.90 29.31 – 1.90	Depositor EDS
% Data completeness (in resolution range)	100.0 (29.36-1.90) 100.0 (29.31-1.90)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	0.75	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.53 (at 1.91Å)	Xtriage
Refinement program	REFMAC 5.1.08	Depositor
R, $R_{free}$	0.190 , 0.252 0.191 , 0.249	Depositor DCC
$R_{free}$ test set	1031 reflections (1.71%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	18.5	Xtriage
Anisotropy	0.586	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.33 , 47.5	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.95	EDS
Total number of atoms	6510	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	23.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.37% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality

### 5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: GOL, PO4, GAD, NAG, NA

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.85	0/5809	0.90	16/7918 (0.2%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1

There are no bond length outliers.

All (16) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed( $^{\circ}$ )	Ideal( $^{\circ}$ )
1	A	46	ASP	CB-CG-OD2	6.85	124.47	118.30
1	A	517	ASP	CB-CG-OD2	6.35	124.01	118.30
1	A	222	ASP	CB-CG-OD2	5.85	123.56	118.30
1	A	732	ASP	CB-CG-OD2	5.83	123.54	118.30
1	A	635[A]	ASP	CB-CG-OD2	5.62	123.35	118.30
1	A	635[B]	ASP	CB-CG-OD2	5.62	123.35	118.30
1	A	436	ASP	CB-CG-OD2	5.57	123.31	118.30
1	A	531	ASP	CB-CG-OD2	5.56	123.31	118.30
1	A	459	ASP	CB-CG-OD2	5.55	123.30	118.30
1	A	430	ASP	CB-CG-OD2	5.38	123.15	118.30
1	A	117	ASP	CB-CG-OD2	5.26	123.04	118.30
1	A	307	ASP	CB-CG-OD2	5.17	122.95	118.30
1	A	96	ASP	CB-CG-OD2	5.05	122.85	118.30
1	A	32	ASP	CB-CG-OD2	5.05	122.84	118.30
1	A	734	ASP	CB-CG-OD2	5.04	122.83	118.30
1	A	472	ASP	CB-CG-OD2	5.03	122.82	118.30



There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	560	GLY	Peptide

## 5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	5629	0	5520	58	0
2	B	26	0	20	0	0
3	A	1	0	0	0	0
4	A	5	0	0	0	0
5	A	12	0	16	2	0
6	A	837	0	0	16	0
All	All	6510	0	5556	58	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (58) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:510[A]:VAL:HG11	1:A:627[A]:MET:CE	1.48	1.41
1:A:510[A]:VAL:HG11	1:A:627[A]:MET:HE3	1.19	1.17
1:A:510[A]:VAL:CG1	1:A:627[A]:MET:CE	2.33	1.05
1:A:4:PRO:HB2	6:A:1732:HOH:O	1.63	0.98
1:A:510[A]:VAL:HG11	1:A:627[A]:MET:HE2	1.48	0.93
1:A:510[A]:VAL:CG1	1:A:627[A]:MET:HE2	1.97	0.92
1:A:510[A]:VAL:CG1	1:A:627[A]:MET:HE3	2.04	0.76
1:A:509[B]:ASP:OD1	6:A:1346:HOH:O	2.08	0.72
1:A:46:ASP:HB2	6:A:1815:HOH:O	1.89	0.71
1:A:498[B]:LEU:HD13	1:A:499:THR:N	2.06	0.71
1:A:746:LEU:HD21	1:A:751:LYS:HE2	1.75	0.69
1:A:692:HIS:HD2	6:A:1719:HOH:O	1.78	0.66
1:A:510[A]:VAL:HG13	1:A:627[A]:MET:HE2	1.77	0.65
1:A:661:LYS:HB2	6:A:1718:HOH:O	2.00	0.62

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:430:ASP:OD1	5:A:801[B]:GOL:H32	2.00	0.61
1:A:430:ASP:OD1	5:A:801[A]:GOL:H11	2.00	0.61
1:A:528:THR:HB	1:A:613[A]:VAL:HG22	1.85	0.59
1:A:415:ARG:NH1	6:A:1391:HOH:O	2.23	0.57
1:A:120:GLU:HG2	1:A:173:LYS:NZ	2.18	0.57
1:A:597:ILE:HD12	6:A:1595:HOH:O	2.05	0.57
1:A:528:THR:HB	1:A:613[A]:VAL:CG2	2.35	0.56
1:A:510[B]:VAL:HG21	1:A:646:TYR:CE2	2.41	0.55
1:A:749:LYS:HE2	6:A:1164:HOH:O	2.05	0.55
1:A:660:PHE:HB3	6:A:1430:HOH:O	2.06	0.54
1:A:515:GLY:HA3	1:A:624:TYR:CE2	2.43	0.53
1:A:736:ARG:HD2	6:A:1511:HOH:O	2.09	0.52
1:A:650:ARG:O	1:A:656:GLN:HA	2.10	0.52
1:A:650:ARG:HD3	1:A:655:ALA:HB3	1.91	0.52
1:A:512:VAL:HG11	1:A:658[A]:VAL:HG11	1.91	0.52
1:A:746:LEU:CD2	1:A:751:LYS:HE2	2.41	0.51
1:A:749:LYS:HG3	6:A:1198:HOH:O	2.12	0.50
1:A:576:LEU:HD21	1:A:627[B]:MET:HE1	1.94	0.50
1:A:746:LEU:O	1:A:749:LYS:HD2	2.12	0.49
1:A:246:LEU:HD23	1:A:246:LEU:C	2.32	0.49
1:A:51:LEU:HD11	1:A:102:ALA:HB1	1.95	0.48
1:A:616:GLY:O	1:A:619:PRO:HD3	2.13	0.48
1:A:226:ARG:NH1	6:A:1519:HOH:O	2.47	0.47
1:A:121:GLU:HG2	1:A:128:TRP:CG	2.50	0.47
1:A:169:GLN:O	1:A:170:PHE:HB2	2.16	0.46
1:A:109:ASP:HA	1:A:112[A]:THR:OG1	2.17	0.45
1:A:211:LEU:O	1:A:214:VAL:HG22	2.17	0.44
1:A:650:ARG:HH21	1:A:654:THR:HB	1.82	0.44
1:A:301:GLU:HB3	1:A:592:TRP:CD2	2.52	0.44
1:A:57:ARG:HD2	1:A:59:SER:O	2.17	0.44
1:A:120:GLU:CG	1:A:173:LYS:NZ	2.81	0.44
1:A:736:ARG:HE	1:A:736:ARG:HB2	1.53	0.44
1:A:723:LEU:HG	1:A:756:LYS:HB2	2.00	0.43
1:A:19:GLN:HE22	1:A:252:LYS:HB3	1.83	0.43
1:A:381:ALA:HB2	6:A:1605:HOH:O	2.18	0.43
1:A:301:GLU:HB3	1:A:592:TRP:CG	2.54	0.43
1:A:12:LEU:HD21	1:A:260:THR:HG21	2.00	0.42
1:A:120:GLU:CD	1:A:173:LYS:HZ2	2.23	0.42
1:A:757:ARG:HH11	1:A:757:ARG:HG3	1.85	0.41
1:A:692:HIS:HE1	6:A:1620:HOH:O	2.04	0.41
1:A:415:ARG:CD	6:A:1391:HOH:O	2.69	0.41

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:751:LYS:HG3	6:A:1681:HOH:O	2.22	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	763/757 (101%)	731 (96%)	28 (4%)	4 (0%)	29 18

All (4) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	559	LEU
1	A	170	PHE
1	A	673	PRO
1	A	579	SER

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	579/570 (102%)	568 (98%)	11 (2%)	57 53

All (11) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	194	ARG
1	A	238	TYR
1	A	255	SER
1	A	406	TYR
1	A	417	TYR
1	A	561	ASP
1	A	658[A]	VAL
1	A	658[B]	VAL
1	A	673	PRO
1	A	702	GLU
1	A	718	THR

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (5) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	19	GLN
1	A	488	GLN
1	A	598	ASN
1	A	605	GLN
1	A	692	HIS

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

2 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
2	NAG	B	1	2	15,15,15	0.86	0	21,21,21	1.25	2 (9%)
2	GAD	B	2	2	7,11,11	2.71	2 (28%)	8,15,15	4.71	3 (37%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	NAG	B	1	2	-	0/6/26/26	0/1/1/1
2	GAD	B	2	2	-	0/0/17/17	0/1/1/1

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	2	GAD	O5-C5	5.01	1.44	1.37
2	B	2	GAD	C3-C4	4.77	1.56	1.50

All (5) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	2	GAD	O5-C5-C4	-12.52	114.24	124.81
2	B	2	GAD	O3-C3-C4	3.40	116.95	109.31
2	B	1	NAG	C6-C5-C4	-2.99	106.01	113.00
2	B	1	NAG	C4-C3-C2	-2.68	106.41	110.34
2	B	2	GAD	C1-O5-C5	-2.16	111.05	115.58


There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.



## 5.6 Ligand geometry [i](#)

Of 4 ligands modelled in this entry, 1 is monoatomic - leaving 3 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
5	GOL	A	801[A]	-	5,5,5	0.42	0	5,5,5	0.48	0
5	GOL	A	801[B]	-	5,5,5	0.43	0	5,5,5	0.78	0
4	PO4	A	850	-	4,4,4	0.52	0	6,6,6	0.84	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	GOL	A	801[B]	-	-	2/4/4/4	-
5	GOL	A	801[A]	-	-	0/4/4/4	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (2) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
5	A	801[B]	GOL	O1-C1-C2-C3
5	A	801[B]	GOL	O1-C1-C2-O2

There are no ring outliers.

2 monomers are involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
5	A	801[A]	GOL	1	0
5	A	801[B]	GOL	1	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	754/757 (99%)	-0.12	15 (1%) 65 68	7, 21, 40, 59	0

All (15) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	4	PRO	7.5
1	A	561	ASP	5.0
1	A	733	ALA	4.8
1	A	28	ALA	3.9
1	A	560	GLY	3.7
1	A	692	HIS	2.9
1	A	551	GLY	2.5
1	A	693	GLY	2.4
1	A	547	GLY	2.3
1	A	726	ALA	2.2
1	A	67	ALA	2.1
1	A	545	ALA	2.1
1	A	546	ALA	2.1
1	A	719	TRP	2.0
1	A	744	THR	2.0

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.


### 6.3 Carbohydrates [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.



Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
2	GAD	B	2	11/11	0.90	0.14	21,25,29,31	0
2	NAG	B	1	15/15	0.95	0.09	12,17,27,31	0

The following is a graphical depiction of the model fit to experimental electron density for oligosaccharide. Each fit is shown from different orientation to approximate a three-dimensional view.



## 6.4 Ligands ⓘ

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
5	GOL	A	801[A]	6/6	0.86	0.19	27,29,30,31	6
5	GOL	A	801[B]	6/6	0.86	0.19	22,22,25,25	6
4	PO4	A	850	5/5	0.96	0.10	21,26,28,29	0
3	NA	A	900	1/1	0.97	0.15	24,24,24,24	0

## 6.5 Other polymers

There are no such residues in this entry.