



wwPDB X-ray Structure Validation Summary Report ⓘ

May 15, 2020 – 01:18 am BST

PDB ID : 3RWA
Title : Crystal structure of circular-permuted mKate
Authors : Wang, Q.; Byrnes, L.; Sondermann, H.
Deposited on : 2011-05-08
Resolution : 1.67 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.11
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.11


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.67 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.





Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	6780 (1.70-1.66)
Clashscore	141614	7310 (1.70-1.66)
Ramachandran outliers	138981	7173 (1.70-1.66)
Sidechain outliers	138945	7172 (1.70-1.66)
RSRZ outliers	127900	6661 (1.70-1.66)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	233	<div> <div>4%</div> <div> <div></div> <div>93%</div> <div>6%</div> </div> </div>
1	B	233	<div> <div>2%</div> <div> <div></div> <div>90%</div> <div>9%</div> </div> </div>
1	C	233	<div> <div>12%</div> <div> <div></div> <div>76%</div> <div>21%</div> </div> </div>
1	D	233	<div> <div>6%</div> <div> <div></div> <div>93%</div> <div>6%</div> </div> </div>
1	E	233	<div> <div>6%</div> <div> <div></div> <div>90%</div> <div>9%</div> </div> </div>
1	F	233	<div> <div>12%</div> <div> <div></div> <div>94%</div> <div>6%</div> </div> </div>

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
1	G	233	
1	H	233	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
1	NRQ	A	133	-	-	X	-
1	NRQ	C	133	-	-	X	-
1	NRQ	D	133	-	-	X	-
1	NRQ	F	133	-	-	X	-
1	NRQ	G	133	-	-	X	-
1	NRQ	H	133	-	-	X	-

2 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 16152 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Fluorescent protein FP480.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	233	Total	C	N	O	S	0	0	0
			1839	1164	312	349	14			
1	B	233	Total	C	N	O	S	0	0	0
			1839	1164	312	349	14			
1	C	233	Total	C	N	O	S	0	0	0
			1839	1164	312	349	14			
1	D	233	Total	C	N	O	S	0	0	0
			1839	1164	312	349	14			
1	E	233	Total	C	N	O	S	0	0	0
			1839	1164	312	349	14			
1	F	233	Total	C	N	O	S	0	0	0
			1839	1164	312	349	14			
1	G	233	Total	C	N	O	S	0	0	0
			1839	1164	312	349	14			
1	H	233	Total	C	N	O	S	0	0	0
			1839	1164	312	349	14			

There are 144 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	7	LEU	PHE	See Remark 999	UNP D0VX33
A	30	ARG	TYR	See Remark 999	UNP D0VX33
A	64	ALA	-	linker	UNP D0VX33
A	65	GLY	-	linker	UNP D0VX33
A	66	GLY	-	linker	UNP D0VX33
A	67	THR	-	linker	UNP D0VX33
A	68	GLY	-	linker	UNP D0VX33
A	69	GLY	-	linker	UNP D0VX33
A	70	SER	-	linker	UNP D0VX33
A	111	MET	GLN	See Remark 999	UNP D0VX33
A	?	-	MET	chromophore	UNP D0VX33
A	?	-	TYR	chromophore	UNP D0VX33
A	133	NRQ	GLY	chromophore	UNP D0VX33

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
A	137	LYS	HIS	See Remark 999	UNP D0VX33
A	150	PHE	TRP	See Remark 999	UNP D0VX33
A	213	SER	HIS	See Remark 999	UNP D0VX33
A	228	SER	ALA	See Remark 999	UNP D0VX33
A	230	MET	LEU	See Remark 999	UNP D0VX33
B	7	LEU	PHE	See Remark 999	UNP D0VX33
B	30	ARG	TYR	See Remark 999	UNP D0VX33
B	64	ALA	-	linker	UNP D0VX33
B	65	GLY	-	linker	UNP D0VX33
B	66	GLY	-	linker	UNP D0VX33
B	67	THR	-	linker	UNP D0VX33
B	68	GLY	-	linker	UNP D0VX33
B	69	GLY	-	linker	UNP D0VX33
B	70	SER	-	linker	UNP D0VX33
B	111	MET	GLN	See Remark 999	UNP D0VX33
B	?	-	MET	chromophore	UNP D0VX33
B	?	-	TYR	chromophore	UNP D0VX33
B	133	NRQ	GLY	chromophore	UNP D0VX33
B	137	LYS	HIS	See Remark 999	UNP D0VX33
B	150	PHE	TRP	See Remark 999	UNP D0VX33
B	213	SER	HIS	See Remark 999	UNP D0VX33
B	228	SER	ALA	See Remark 999	UNP D0VX33
B	230	MET	LEU	See Remark 999	UNP D0VX33
C	7	LEU	PHE	See Remark 999	UNP D0VX33
C	30	ARG	TYR	See Remark 999	UNP D0VX33
C	64	ALA	-	linker	UNP D0VX33
C	65	GLY	-	linker	UNP D0VX33
C	66	GLY	-	linker	UNP D0VX33
C	67	THR	-	linker	UNP D0VX33
C	68	GLY	-	linker	UNP D0VX33
C	69	GLY	-	linker	UNP D0VX33
C	70	SER	-	linker	UNP D0VX33
C	111	MET	GLN	See Remark 999	UNP D0VX33
C	?	-	MET	chromophore	UNP D0VX33
C	?	-	TYR	chromophore	UNP D0VX33
C	133	NRQ	GLY	chromophore	UNP D0VX33
C	137	LYS	HIS	See Remark 999	UNP D0VX33
C	150	PHE	TRP	See Remark 999	UNP D0VX33
C	213	SER	HIS	See Remark 999	UNP D0VX33
C	228	SER	ALA	See Remark 999	UNP D0VX33
C	230	MET	LEU	See Remark 999	UNP D0VX33
D	7	LEU	PHE	See Remark 999	UNP D0VX33

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
D	30	ARG	TYR	See Remark 999	UNP D0VX33
D	64	ALA	-	linker	UNP D0VX33
D	65	GLY	-	linker	UNP D0VX33
D	66	GLY	-	linker	UNP D0VX33
D	67	THR	-	linker	UNP D0VX33
D	68	GLY	-	linker	UNP D0VX33
D	69	GLY	-	linker	UNP D0VX33
D	70	SER	-	linker	UNP D0VX33
D	111	MET	GLN	See Remark 999	UNP D0VX33
D	?	-	MET	chromophore	UNP D0VX33
D	?	-	TYR	chromophore	UNP D0VX33
D	133	NRQ	GLY	chromophore	UNP D0VX33
D	137	LYS	HIS	See Remark 999	UNP D0VX33
D	150	PHE	TRP	See Remark 999	UNP D0VX33
D	213	SER	HIS	See Remark 999	UNP D0VX33
D	228	SER	ALA	See Remark 999	UNP D0VX33
D	230	MET	LEU	See Remark 999	UNP D0VX33
E	7	LEU	PHE	See Remark 999	UNP D0VX33
E	30	ARG	TYR	See Remark 999	UNP D0VX33
E	64	ALA	-	linker	UNP D0VX33
E	65	GLY	-	linker	UNP D0VX33
E	66	GLY	-	linker	UNP D0VX33
E	67	THR	-	linker	UNP D0VX33
E	68	GLY	-	linker	UNP D0VX33
E	69	GLY	-	linker	UNP D0VX33
E	70	SER	-	linker	UNP D0VX33
E	111	MET	GLN	See Remark 999	UNP D0VX33
E	?	-	MET	chromophore	UNP D0VX33
E	?	-	TYR	chromophore	UNP D0VX33
E	133	NRQ	GLY	chromophore	UNP D0VX33
E	137	LYS	HIS	See Remark 999	UNP D0VX33
E	150	PHE	TRP	See Remark 999	UNP D0VX33
E	213	SER	HIS	See Remark 999	UNP D0VX33
E	228	SER	ALA	See Remark 999	UNP D0VX33
E	230	MET	LEU	See Remark 999	UNP D0VX33
F	7	LEU	PHE	See Remark 999	UNP D0VX33
F	30	ARG	TYR	See Remark 999	UNP D0VX33
F	64	ALA	-	linker	UNP D0VX33
F	65	GLY	-	linker	UNP D0VX33
F	66	GLY	-	linker	UNP D0VX33
F	67	THR	-	linker	UNP D0VX33
F	68	GLY	-	linker	UNP D0VX33

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
F	69	GLY	-	linker	UNP D0VX33
F	70	SER	-	linker	UNP D0VX33
F	111	MET	GLN	See Remark 999	UNP D0VX33
F	?	-	MET	chromophore	UNP D0VX33
F	?	-	TYR	chromophore	UNP D0VX33
F	133	NRQ	GLY	chromophore	UNP D0VX33
F	137	LYS	HIS	See Remark 999	UNP D0VX33
F	150	PHE	TRP	See Remark 999	UNP D0VX33
F	213	SER	HIS	See Remark 999	UNP D0VX33
F	228	SER	ALA	See Remark 999	UNP D0VX33
F	230	MET	LEU	See Remark 999	UNP D0VX33
G	7	LEU	PHE	See Remark 999	UNP D0VX33
G	30	ARG	TYR	See Remark 999	UNP D0VX33
G	64	ALA	-	linker	UNP D0VX33
G	65	GLY	-	linker	UNP D0VX33
G	66	GLY	-	linker	UNP D0VX33
G	67	THR	-	linker	UNP D0VX33
G	68	GLY	-	linker	UNP D0VX33
G	69	GLY	-	linker	UNP D0VX33
G	70	SER	-	linker	UNP D0VX33
G	111	MET	GLN	See Remark 999	UNP D0VX33
G	?	-	MET	chromophore	UNP D0VX33
G	?	-	TYR	chromophore	UNP D0VX33
G	133	NRQ	GLY	chromophore	UNP D0VX33
G	137	LYS	HIS	See Remark 999	UNP D0VX33
G	150	PHE	TRP	See Remark 999	UNP D0VX33
G	213	SER	HIS	See Remark 999	UNP D0VX33
G	228	SER	ALA	See Remark 999	UNP D0VX33
G	230	MET	LEU	See Remark 999	UNP D0VX33
H	7	LEU	PHE	See Remark 999	UNP D0VX33
H	30	ARG	TYR	See Remark 999	UNP D0VX33
H	64	ALA	-	linker	UNP D0VX33
H	65	GLY	-	linker	UNP D0VX33
H	66	GLY	-	linker	UNP D0VX33
H	67	THR	-	linker	UNP D0VX33
H	68	GLY	-	linker	UNP D0VX33
H	69	GLY	-	linker	UNP D0VX33
H	70	SER	-	linker	UNP D0VX33
H	111	MET	GLN	See Remark 999	UNP D0VX33
H	?	-	MET	chromophore	UNP D0VX33
H	?	-	TYR	chromophore	UNP D0VX33
H	133	NRQ	GLY	chromophore	UNP D0VX33

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
H	137	LYS	HIS	See Remark 999	UNP D0VX33
H	150	PHE	TRP	See Remark 999	UNP D0VX33
H	213	SER	HIS	See Remark 999	UNP D0VX33
H	228	SER	ALA	See Remark 999	UNP D0VX33
H	230	MET	LEU	See Remark 999	UNP D0VX33


- Molecule 2 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	A	213	Total O 213 213	0	0
2	B	216	Total O 216 216	0	0
2	C	186	Total O 186 186	0	0
2	D	178	Total O 178 178	0	0
2	E	194	Total O 194 194	0	0
2	F	169	Total O 169 169	0	0
2	G	154	Total O 154 154	0	0
2	H	130	Total O 130 130	0	0



3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.



- Molecule 1: Fluorescent protein FP480





- Molecule 1: Fluorescent protein FP480




- Molecule 1: Fluorescent protein FP480




- Molecule 1: Fluorescent protein FP480




- Molecule 1: Fluorescent protein FP480









- Molecule 1: Fluorescent protein FP480



- Molecule 1: Fluorescent protein FP480



- Molecule 1: Fluorescent protein FP480



4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, α , β , γ	109.36Å 93.28Å 194.37Å 90.00° 96.52° 90.00°	Depositor
Resolution (Å)	38.31 – 1.67 38.31 – 1.67	Depositor EDS
% Data completeness (in resolution range)	99.2 (38.31-1.67) 97.5 (38.31-1.67)	Depositor EDS
R_{merge}	0.06	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.31 (at 1.67Å)	Xtriage
Refinement program	PHENIX 1.7.1_743, REFMAC	Depositor
R, R_{free}	0.182 , 0.215 0.187 , 0.212	Depositor DCC
R_{free} test set	2000 reflections (0.90%)	wwPDB-VP
Wilson B-factor (Å ²)	16.8	Xtriage
Anisotropy	0.103	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.36 , 45.6	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	16152	wwPDB-VP
Average B, all atoms (Å ²)	20.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.45% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: NRQ

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.38	0/1856	0.60	0/2502
1	B	0.38	0/1856	0.59	0/2502
1	C	0.31	0/1856	0.52	0/2502
1	D	0.36	0/1856	0.56	0/2502
1	E	0.34	0/1856	0.55	0/2502
1	F	0.35	0/1856	0.55	0/2502
1	G	0.34	0/1856	0.55	0/2502
1	H	0.31	0/1856	0.51	0/2502
All	All	0.35	0/14848	0.55	0/20016

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1839	0	1802	23	0
1	B	1839	0	1802	25	0
1	C	1839	0	1802	55	0
1	D	1839	0	1802	24	0
1	E	1839	0	1802	38	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	F	1839	0	1802	23	0
1	G	1839	0	1802	32	0
1	H	1839	0	1802	22	0
2	A	213	0	0	5	0
2	B	216	0	0	3	0
2	C	186	0	0	3	0
2	D	178	0	0	0	0
2	E	194	0	0	6	0
2	F	169	0	0	3	0
2	G	154	0	0	3	0
2	H	130	0	0	2	0
All	All	16152	0	14416	236	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 8.

The worst 5 of 236 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:D:63:HIS:CB	1:D:64:ALA:HB3	1.49	1.40
1:F:61:LEU:HD12	2:F:461:HOH:O	1.26	1.34
1:G:63:HIS:HB2	1:G:64:ALA:CB	1.58	1.32
1:G:63:HIS:CB	1:G:64:ALA:HB2	1.65	1.26
1:D:63:HIS:HB3	1:D:64:ALA:CB	1.72	1.18

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	228/233 (98%)	219 (96%)	9 (4%)	0	100 100

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	B	228/233 (98%)	224 (98%)	4 (2%)	0	100	100
1	C	228/233 (98%)	220 (96%)	6 (3%)	2 (1%)	17	4
1	D	228/233 (98%)	221 (97%)	6 (3%)	1 (0%)	34	17
1	E	228/233 (98%)	218 (96%)	9 (4%)	1 (0%)	34	17
1	F	228/233 (98%)	223 (98%)	5 (2%)	0	100	100
1	G	228/233 (98%)	222 (97%)	5 (2%)	1 (0%)	34	17
1	H	228/233 (98%)	223 (98%)	5 (2%)	0	100	100
All	All	1824/1864 (98%)	1770 (97%)	49 (3%)	5 (0%)	41	23

All (5) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	D	64	ALA
1	G	64	ALA
1	C	221	ASP
1	C	65	GLY
1	E	64	ALA

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	197/197 (100%)	191 (97%)	6 (3%)	41	20
1	B	197/197 (100%)	193 (98%)	4 (2%)	55	36
1	C	197/197 (100%)	187 (95%)	10 (5%)	24	7
1	D	197/197 (100%)	191 (97%)	6 (3%)	41	20
1	E	197/197 (100%)	193 (98%)	4 (2%)	55	36
1	F	197/197 (100%)	194 (98%)	3 (2%)	65	48
1	G	197/197 (100%)	194 (98%)	3 (2%)	65	48
1	H	197/197 (100%)	188 (95%)	9 (5%)	27	9

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
All	All	1576/1576 (100%)	1531 (97%)	45 (3%)	42 21

5 of 45 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	D	36	LYS
1	D	221	ASP
1	H	81	MET
1	D	56	ASP
1	E	36	LYS

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
1	B	63	HIS
1	F	63	HIS

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

8 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
1	NRQ	D	133	1	23,24,25	5.97	7 (30%)	23,32,34	4.27	6 (26%)
1	NRQ	B	133	1	23,24,25	6.00	7 (30%)	23,32,34	4.24	6 (26%)
1	NRQ	A	133	1	23,24,25	5.99	7 (30%)	23,32,34	4.19	6 (26%)
1	NRQ	C	133	1	23,24,25	5.97	7 (30%)	23,32,34	4.21	6 (26%)
1	NRQ	F	133	1	23,24,25	6.00	7 (30%)	23,32,34	4.18	6 (26%)

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
1	NRQ	E	133	1	23,24,25	5.98	7 (30%)	23,32,34	4.26	6 (26%)
1	NRQ	G	133	1	23,24,25	5.98	7 (30%)	23,32,34	4.19	6 (26%)
1	NRQ	H	133	1	23,24,25	6.00	7 (30%)	23,32,34	4.28	6 (26%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	NRQ	D	133	1	-	2/9/31/32	0/2/2/2
1	NRQ	B	133	1	-	1/9/31/32	0/2/2/2
1	NRQ	A	133	1	-	2/9/31/32	0/2/2/2
1	NRQ	C	133	1	-	2/9/31/32	0/2/2/2
1	NRQ	F	133	1	-	3/9/31/32	0/2/2/2
1	NRQ	E	133	1	-	3/9/31/32	0/2/2/2
1	NRQ	G	133	1	-	2/9/31/32	0/2/2/2
1	NRQ	H	133	1	-	2/9/31/32	0/2/2/2

The worst 5 of 56 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	F	133	NRQ	CB2-CA2	21.80	1.53	1.35
1	B	133	NRQ	CB2-CA2	21.76	1.53	1.35
1	H	133	NRQ	CB2-CA2	21.74	1.53	1.35
1	G	133	NRQ	CB2-CA2	21.72	1.53	1.35
1	A	133	NRQ	CB2-CA2	21.68	1.53	1.35

The worst 5 of 48 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	H	133	NRQ	CG2-CB2-CA2	-13.54	113.35	129.94
1	D	133	NRQ	CG2-CB2-CA2	-13.23	113.73	129.94
1	E	133	NRQ	CG2-CB2-CA2	-13.21	113.75	129.94
1	G	133	NRQ	CG2-CB2-CA2	-13.07	113.92	129.94
1	B	133	NRQ	CG2-CB2-CA2	-12.93	114.10	129.94

There are no chirality outliers.

5 of 17 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
1	D	133	NRQ	CA1-CB1-CG1-SD
1	B	133	NRQ	CA1-CB1-CG1-SD
1	A	133	NRQ	CA1-CB1-CG1-SD
1	A	133	NRQ	C1-CA1-CB1-CG1
1	C	133	NRQ	CA1-CB1-CG1-SD

There are no ring outliers.

8 monomers are involved in 71 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
1	D	133	NRQ	9	0
1	B	133	NRQ	8	0
1	A	133	NRQ	9	0
1	C	133	NRQ	9	0
1	F	133	NRQ	9	0
1	E	133	NRQ	8	0
1	G	133	NRQ	9	0
1	H	133	NRQ	10	0

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

The following chains have linkage breaks:

Mol	Chain	Number of breaks
1	A	1
1	D	1
1	E	1

All chain breaks are listed below:

Model	Chain	Residue-1	Atom-1	Residue-2	Atom-2	Distance (Å)
1	A	132:PHE	C	133:NRQ	N1	1.65
1	D	132:PHE	C	133:NRQ	N1	1.62
1	E	133:NRQ	C3	136:SER	N	1.60

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	232/233 (99%)	0.31	9 (3%) 39 42	9, 15, 29, 65	0
1	B	232/233 (99%)	0.06	5 (2%) 62 65	9, 15, 28, 37	0
1	C	232/233 (99%)	0.75	27 (11%) 4 5	11, 17, 34, 67	0
1	D	232/233 (99%)	0.29	13 (5%) 24 25	11, 17, 34, 65	0
1	E	232/233 (99%)	0.35	14 (6%) 21 22	11, 17, 32, 69	0
1	F	232/233 (99%)	0.61	27 (11%) 4 5	11, 18, 40, 66	0
1	G	232/233 (99%)	0.56	21 (9%) 9 9	12, 19, 41, 71	0
1	H	232/233 (99%)	0.96	35 (15%) 2 2	13, 25, 43, 69	0
All	All	1856/1864 (99%)	0.49	151 (8%) 12 13	9, 18, 37, 71	0

The worst 5 of 151 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	C	64	ALA	23.4
1	G	65	GLY	20.1
1	H	64	ALA	19.2
1	A	64	ALA	19.0
1	H	63	HIS	16.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
1	NRQ	C	133	23/24	0.88	0.15	15,17,25,27	0

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
1	NRQ	H	133	23/24	0.88	0.14	19,23,31,34	0
1	NRQ	G	133	23/24	0.89	0.13	16,19,25,27	0
1	NRQ	E	133	23/24	0.89	0.15	15,17,25,28	0
1	NRQ	D	133	23/24	0.90	0.14	14,18,24,25	0
1	NRQ	A	133	23/24	0.91	0.14	12,15,22,25	0
1	NRQ	B	133	23/24	0.91	0.13	11,15,22,23	0
1	NRQ	F	133	23/24	0.91	0.13	14,18,26,28	0

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.