



# Full wwPDB X-ray Structure Validation Report ⓘ

Dec 10, 2022 – 06:43 PM EST

PDB ID : 1RY6  
Title : Crystal Structure of Internal Kinesin Motor Domain  
Authors : Shipley, K.; Hekmat-Nejad, M.; Turner, J.; Moores, C.; Anderson, R.; Milligan, R.; Sakowicz, R.; Fletterick, R.  
Deposited on : 2003-12-19  
Resolution : 1.60 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix)	:	1.13
EDS	:	2.31.2
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac	:	5.8.0158
CCP4	:	7.0.044 (Gargrove)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.31.2


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*


The reported resolution of this entry is 1.60 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	3398 (1.60-1.60)
Clashscore	141614	3665 (1.60-1.60)
Ramachandran outliers	138981	3564 (1.60-1.60)
Sidechain outliers	138945	3563 (1.60-1.60)
RSRZ outliers	127900	3321 (1.60-1.60)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	360	

## 2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 2829 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called INTERNAL KINESIN.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	319	2518	1591	426	489	12	0	0	0

There are 31 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	1	MET	-	initiating methionine	UNP Q8I4Y0
A	331	LYS	-	cloning artifact	UNP Q8I4Y0
A	332	GLY	-	cloning artifact	UNP Q8I4Y0
A	333	ASN	-	cloning artifact	UNP Q8I4Y0
A	334	SER	-	cloning artifact	UNP Q8I4Y0
A	335	LYS	-	cloning artifact	UNP Q8I4Y0
A	336	LEU	-	cloning artifact	UNP Q8I4Y0
A	337	GLU	-	cloning artifact	UNP Q8I4Y0
A	338	GLY	-	cloning artifact	UNP Q8I4Y0
A	339	LYS	-	cloning artifact	UNP Q8I4Y0
A	340	PRO	-	cloning artifact	UNP Q8I4Y0
A	341	ILE	-	cloning artifact	UNP Q8I4Y0
A	342	PRO	-	cloning artifact	UNP Q8I4Y0
A	343	ASN	-	cloning artifact	UNP Q8I4Y0
A	344	PRO	-	cloning artifact	UNP Q8I4Y0
A	345	LEU	-	cloning artifact	UNP Q8I4Y0
A	346	LEU	-	cloning artifact	UNP Q8I4Y0
A	347	GLY	-	cloning artifact	UNP Q8I4Y0
A	348	LEU	-	cloning artifact	UNP Q8I4Y0
A	349	ASP	-	cloning artifact	UNP Q8I4Y0
A	350	SER	-	cloning artifact	UNP Q8I4Y0
A	351	SER	-	cloning artifact	UNP Q8I4Y0
A	352	ARG	-	cloning artifact	UNP Q8I4Y0
A	353	THR	-	cloning artifact	UNP Q8I4Y0
A	354	GLY	-	cloning artifact	UNP Q8I4Y0
A	355	HIS	-	cloning artifact	UNP Q8I4Y0
A	356	HIS	-	cloning artifact	UNP Q8I4Y0

*Continued on next page...*

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
A	357	HIS	-	cloning artifact	UNP Q8I4Y0
A	358	HIS	-	cloning artifact	UNP Q8I4Y0
A	359	HIS	-	cloning artifact	UNP Q8I4Y0
A	360	HIS	-	cloning artifact	UNP Q8I4Y0

- Molecule 2 is SULFATE ION (three-letter code: SO4) (formula: O<sub>4</sub>S).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	O	S	0	0
			5	4	1		


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	306	Total	O	0	0
			306	306		

### 3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

#### • Molecule 1: INTERNAL KINESIN



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 32 2 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	105.59Å 105.59Å 84.77Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	24.92 – 1.60 24.91 – 1.50	Depositor EDS
% Data completeness (in resolution range)	91.6 (24.92-1.60) 87.9 (24.91-1.50)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	0.05	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.13 (at 1.50Å)	Xtriage
Refinement program	REFMAC 5.1.24	Depositor
R, $R_{free}$	0.205 , 0.231 0.217 , 0.214	Depositor DCC
$R_{free}$ test set	6715 reflections (8.74%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	15.4	Xtriage
Anisotropy	0.131	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.36 , 45.9	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	0.028 for -h,-k,l	Xtriage
$F_o, F_c$ correlation	0.96	EDS
Total number of atoms	2829	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	31.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.73% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality

### 5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: SO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	1.06	4/2551 (0.2%)	1.04	8/3437 (0.2%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1

All (4) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	84	CYS	CB-SG	-6.69	1.70	1.82
1	A	186	GLU	CG-CD	5.51	1.60	1.51
1	A	209	SER	CB-OG	-5.27	1.35	1.42
1	A	211	ARG	CZ-NH1	5.26	1.39	1.33

All (8) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	55	ASP	CB-CG-OD2	8.65	126.09	118.30
1	A	219	ASP	CB-CG-OD1	6.56	124.21	118.30
1	A	277	ASP	CB-CG-OD2	6.47	124.12	118.30
1	A	256	ASP	CB-CG-OD1	6.33	124.00	118.30
1	A	143	PHE	CB-CG-CD1	5.57	124.70	120.80
1	A	20	ASP	CB-CG-OD2	5.50	123.25	118.30
1	A	68	TYR	CB-CG-CD2	5.24	124.14	121.00
1	A	60	ASP	CB-CG-OD2	5.03	122.83	118.30

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	277	ASP	Peptide

## 5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2518	0	2556	29	0
2	A	5	0	0	0	0
3	A	306	0	0	6	0
All	All	2829	0	2556	29	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

All (29) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:156:GLN:NE2	1:A:177:LYS:HZ1	1.54	1.04
1:A:38:ARG:HH11	1:A:38:ARG:HG3	1.21	1.03
1:A:156:GLN:HE22	1:A:177:LYS:HZ1	0.98	0.96
1:A:156:GLN:NE2	1:A:177:LYS:NZ	2.23	0.85
1:A:299:LYS:HE3	3:A:497:HOH:O	1.92	0.70
1:A:38:ARG:HG3	1:A:38:ARG:NH1	1.98	0.69
1:A:38:ARG:NH1	1:A:48:GLU:OE1	2.25	0.69
1:A:137:LYS:NZ	3:A:687:HOH:O	2.30	0.63
1:A:73:LYS:NZ	3:A:554:HOH:O	2.34	0.60
1:A:142:SER:OG	1:A:178:ILE:HD13	2.03	0.59
1:A:205:GLN:OE1	1:A:252:GLN:HG3	2.02	0.59
1:A:105:SER:HG	1:A:111:SER:HG	1.53	0.56
1:A:246:THR:HG21	3:A:694:HOH:O	2.08	0.53
1:A:274:MET:HG3	1:A:296:PHE:O	2.12	0.49
1:A:24:ILE:HD13	1:A:308:ILE:HD12	1.95	0.49
1:A:274:MET:HE2	1:A:295:ILE:O	2.13	0.49
1:A:279:ASN:HB3	1:A:280:HIS:ND1	2.28	0.48

*Continued on next page...*



Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:71:THR:HB	1:A:303:ILE:HD13	1.95	0.47
1:A:16:LYS:O	1:A:19:LYS:NZ	2.33	0.46
1:A:297:VAL:HG21	1:A:299:LYS:NZ	2.31	0.45
1:A:323:ARG:HD2	3:A:458:HOH:O	2.16	0.45
1:A:172:VAL:HG23	1:A:174:LYS:HZ3	1.83	0.43
1:A:172:VAL:CG2	1:A:174:LYS:NZ	2.82	0.43
1:A:297:VAL:HG23	1:A:297:VAL:O	2.18	0.43
1:A:142:SER:OG	1:A:178:ILE:CD1	2.66	0.42
1:A:168:LYS:HG3	1:A:169:LYS:N	2.32	0.42
1:A:36:GLU:HA	1:A:37:PRO:HD3	1.83	0.41
1:A:51:GLU:O	3:A:603:HOH:O	2.22	0.41
1:A:42:ASP:O	1:A:43:MET:CB	2.69	0.41

There are no symmetry-related clashes.

## 5.3 Torsion angles

### 5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	315/360 (88%)	307 (98%)	4 (1%)	4 (1%)	<b>12</b> <b>2</b>

All (4) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	42	ASP
1	A	43	MET
1	A	276	SER
1	A	41	VAL

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	286/325 (88%)	263 (92%)	23 (8%)	12 2

All (23) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	19	LYS
1	A	21	SER
1	A	38	ARG
1	A	41	VAL
1	A	42	ASP
1	A	44	THR
1	A	129	ASN
1	A	156	GLN
1	A	168	LYS
1	A	174	LYS
1	A	180	ARG
1	A	191	MET
1	A	205	GLN
1	A	206	ASN
1	A	256	ASP
1	A	274	MET
1	A	276	SER
1	A	277	ASP
1	A	278	LYS
1	A	279	ASN
1	A	280	HIS
1	A	299	LYS
1	A	329	LYS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (4) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	156	GLN
1	A	206	ASN

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	A	261	ASN
1	A	279	ASN

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates ⓘ

There are no monosaccharides in this entry.

## 5.6 Ligand geometry ⓘ

1 ligand is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
2	SO4	A	401	-	4,4,4	0.43	0	6,6,6	1.14	1 (16%)

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed( $^{\circ}$ )	Ideal( $^{\circ}$ )
2	A	401	SO4	O4-S-O1	-2.45	96.54	109.31

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

## 5.7 Other polymers

There are no such residues in this entry.

## 5.8 Polymer linkage issues

There are no chain breaks in this entry.

## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	319/360 (88%)	0.07	24 (7%) <b>14</b> <b>12</b>	15, 26, 61, 82	0

All (24) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	41	VAL	10.2
1	A	39	TYR	10.0
1	A	43	MET	7.7
1	A	40	LYS	7.5
1	A	44	THR	7.1
1	A	277	ASP	6.5
1	A	279	ASN	5.8
1	A	280	HIS	5.1
1	A	330	ASN	4.7
1	A	42	ASP	4.3
1	A	45	LYS	4.3
1	A	276	SER	3.6
1	A	46	TYR	3.5
1	A	298	GLY	3.2
1	A	278	LYS	2.8
1	A	297	VAL	2.7
1	A	174	LYS	2.7
1	A	168	LYS	2.6
1	A	169	LYS	2.6
1	A	1	MET	2.6
1	A	224	ASN	2.3
1	A	327	ARG	2.2
1	A	47	ILE	2.2
1	A	156	GLN	2.1

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
2	SO4	A	401	5/5	0.99	0.05	16,18,18,21	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.