



wwPDB X-ray Structure Validation Summary Report ⓘ

May 16, 2020 – 12:55 pm BST

PDB ID : 1RZ1
Title : Reduced flavin reductase PheA2 in complex with NAD
Authors : Van Den Heuvel, R.H.; Westphal, A.H.; Heck, A.J.; Walsh, M.A.; Rovida, S.;
Van Berkel, W.J.; Mattevi, A.
Deposited on : 2003-12-23
Resolution : 2.10 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.11
buster-report : 1.1.7 (2018)
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.11


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.10 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	5197 (2.10-2.10)
Clashscore	141614	5710 (2.10-2.10)
Ramachandran outliers	138981	5647 (2.10-2.10)
Sidechain outliers	138945	5648 (2.10-2.10)
RSRZ outliers	127900	5083 (2.10-2.10)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	161	<div> <div>2%</div> <div>78% 16% • 5%</div> </div>
1	B	161	<div> <div>2%</div> <div>75% 18% • 5%</div> </div>
1	C	161	<div> <div>2%</div> <div>76% 17% • 5%</div> </div>
1	D	161	<div> <div>2%</div> <div>75% 19% • 5%</div> </div>
1	E	161	<div> <div>2%</div> <div>78% 15% • 5%</div> </div>
1	F	161	<div> <div>3%</div> <div>75% 19% • 5%</div> </div>

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
1	G	161	<div><div></div><div>4%</div><div>74%</div><div>20%</div><div>• 5%</div></div>
1	H	161	<div><div></div><div>5%</div><div>78%</div><div>16%</div><div>• 5%</div></div>

2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 10581 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called phenol 2-hydroxylase component B.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	153	Total	C	N	O	S	Se	0	0	0
			1176	749	196	224	1	6			
1	B	153	Total	C	N	O	S	Se	0	0	0
			1176	749	196	224	1	6			
1	C	153	Total	C	N	O	S	Se	0	0	0
			1176	749	196	224	1	6			
1	D	153	Total	C	N	O	S	Se	0	0	0
			1176	749	196	224	1	6			
1	E	153	Total	C	N	O	S	Se	0	0	0
			1176	749	196	224	1	6			
1	F	153	Total	C	N	O	S	Se	0	0	0
			1176	749	196	224	1	6			
1	G	153	Total	C	N	O	S	Se	0	0	0
			1176	749	196	224	1	6			
1	H	153	Total	C	N	O	S	Se	0	0	0
			1176	749	196	224	1	6			

There are 48 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	1	MSE	MET	MODIFIED RESIDUE	UNP Q9LAG2
A	10	MSE	MET	MODIFIED RESIDUE	UNP Q9LAG2
A	31	MSE	MET	MODIFIED RESIDUE	UNP Q9LAG2
A	37	MSE	MET	MODIFIED RESIDUE	UNP Q9LAG2
A	56	MSE	MET	MODIFIED RESIDUE	UNP Q9LAG2
A	80	MSE	MET	MODIFIED RESIDUE	UNP Q9LAG2
B	1	MSE	MET	MODIFIED RESIDUE	UNP Q9LAG2
B	10	MSE	MET	MODIFIED RESIDUE	UNP Q9LAG2
B	31	MSE	MET	MODIFIED RESIDUE	UNP Q9LAG2
B	37	MSE	MET	MODIFIED RESIDUE	UNP Q9LAG2
B	56	MSE	MET	MODIFIED RESIDUE	UNP Q9LAG2
B	80	MSE	MET	MODIFIED RESIDUE	UNP Q9LAG2
C	1	MSE	MET	MODIFIED RESIDUE	UNP Q9LAG2

Continued on next page...

Continued from previous page...


Chain	Residue	Modelled	Actual	Comment	Reference
C	10	MSE	MET	MODIFIED RESIDUE	UNP Q9LAG2
C	31	MSE	MET	MODIFIED RESIDUE	UNP Q9LAG2
C	37	MSE	MET	MODIFIED RESIDUE	UNP Q9LAG2
C	56	MSE	MET	MODIFIED RESIDUE	UNP Q9LAG2
C	80	MSE	MET	MODIFIED RESIDUE	UNP Q9LAG2
D	1	MSE	MET	MODIFIED RESIDUE	UNP Q9LAG2
D	10	MSE	MET	MODIFIED RESIDUE	UNP Q9LAG2
D	31	MSE	MET	MODIFIED RESIDUE	UNP Q9LAG2
D	37	MSE	MET	MODIFIED RESIDUE	UNP Q9LAG2
D	56	MSE	MET	MODIFIED RESIDUE	UNP Q9LAG2
D	80	MSE	MET	MODIFIED RESIDUE	UNP Q9LAG2
E	1	MSE	MET	MODIFIED RESIDUE	UNP Q9LAG2
E	10	MSE	MET	MODIFIED RESIDUE	UNP Q9LAG2
E	31	MSE	MET	MODIFIED RESIDUE	UNP Q9LAG2
E	37	MSE	MET	MODIFIED RESIDUE	UNP Q9LAG2
E	56	MSE	MET	MODIFIED RESIDUE	UNP Q9LAG2
E	80	MSE	MET	MODIFIED RESIDUE	UNP Q9LAG2
F	1	MSE	MET	MODIFIED RESIDUE	UNP Q9LAG2
F	10	MSE	MET	MODIFIED RESIDUE	UNP Q9LAG2
F	31	MSE	MET	MODIFIED RESIDUE	UNP Q9LAG2
F	37	MSE	MET	MODIFIED RESIDUE	UNP Q9LAG2
F	56	MSE	MET	MODIFIED RESIDUE	UNP Q9LAG2
F	80	MSE	MET	MODIFIED RESIDUE	UNP Q9LAG2
G	1	MSE	MET	MODIFIED RESIDUE	UNP Q9LAG2
G	10	MSE	MET	MODIFIED RESIDUE	UNP Q9LAG2
G	31	MSE	MET	MODIFIED RESIDUE	UNP Q9LAG2
G	37	MSE	MET	MODIFIED RESIDUE	UNP Q9LAG2
G	56	MSE	MET	MODIFIED RESIDUE	UNP Q9LAG2
G	80	MSE	MET	MODIFIED RESIDUE	UNP Q9LAG2
H	1	MSE	MET	MODIFIED RESIDUE	UNP Q9LAG2
H	10	MSE	MET	MODIFIED RESIDUE	UNP Q9LAG2
H	31	MSE	MET	MODIFIED RESIDUE	UNP Q9LAG2
H	37	MSE	MET	MODIFIED RESIDUE	UNP Q9LAG2
H	56	MSE	MET	MODIFIED RESIDUE	UNP Q9LAG2
H	80	MSE	MET	MODIFIED RESIDUE	UNP Q9LAG2

- Molecule 2 is FLAVIN-ADENINE DINUCLEOTIDE (three-letter code: FAD) (formula: C₂₇H₃₃N₉O₁₅P₂).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total	C	N	O	P	0	0
			53	27	9	15	2		
2	B	1	Total	C	N	O	P	0	0
			53	27	9	15	2		
2	C	1	Total	C	N	O	P	0	0
			53	27	9	15	2		
2	D	1	Total	C	N	O	P	0	0
			53	27	9	15	2		
2	E	1	Total	C	N	O	P	0	0
			53	27	9	15	2		
2	F	1	Total	C	N	O	P	0	0
			53	27	9	15	2		
2	G	1	Total	C	N	O	P	0	0
			53	27	9	15	2		
2	H	1	Total	C	N	O	P	0	0
			53	27	9	15	2		

- Molecule 3 is NICOTINAMIDE-ADENINE-DINUCLEOTIDE (three-letter code: NAD) (formula: C₂₁H₂₇N₇O₁₄P₂).



- Molecule 4 is water.

Continued on next page...


Continued from previous page...

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	E	40	Total 40	O 40	0	0
4	F	44	Total 44	O 44	0	0
4	G	47	Total 47	O 47	0	0
4	H	43	Total 43	O 43	0	0



3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.



- Molecule 1: phenol 2-hydroxylase component B




- Molecule 1: phenol 2-hydroxylase component B




- Molecule 1: phenol 2-hydroxylase component B




- Molecule 1: phenol 2-hydroxylase component B




- Molecule 1: phenol 2-hydroxylase component B




- Molecule 1: phenol 2-hydroxylase component B



- Molecule 1: phenol 2-hydroxylase component B



- Molecule 1: phenol 2-hydroxylase component B



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	53.69Å 156.46Å 83.92Å 90.00° 91.11° 90.00°	Depositor
Resolution (Å)	15.00 – 2.10 14.94 – 2.10	Depositor EDS
% Data completeness (in resolution range)	95.2 (15.00-2.10) 95.1 (14.94-2.10)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.50 (at 2.10Å)	Xtriage
Refinement program	REFMAC 5.1.24	Depositor
R, R_{free}	0.237 , 0.256 0.234 , 0.260	Depositor DCC
R_{free} test set	1560 reflections (2.04%)	wwPDB-VP
Wilson B-factor (Å ²)	34.1	Xtriage
Anisotropy	0.288	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.41 , 34.5	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtriage
Estimated twinning fraction	0.229 for h,-k,-l	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	10581	wwPDB-VP
Average B, all atoms (Å ²)	34.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 26.77 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 2.4791e-03. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: FAD, NAD

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.67	0/1187	0.78	2/1592 (0.1%)
1	B	0.66	0/1187	0.79	2/1592 (0.1%)
1	C	0.65	0/1187	0.78	3/1592 (0.2%)
1	D	0.64	0/1187	0.77	2/1592 (0.1%)
1	E	0.62	0/1187	0.77	2/1592 (0.1%)
1	F	0.68	0/1187	0.79	2/1592 (0.1%)
1	G	0.63	0/1187	0.77	4/1592 (0.3%)
1	H	0.63	0/1187	0.78	2/1592 (0.1%)
All	All	0.65	0/9496	0.78	19/12736 (0.1%)

There are no bond length outliers.

The worst 5 of 19 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	E	3	ASP	CB-CG-OD2	6.30	123.97	118.30
1	C	3	ASP	CB-CG-OD2	6.02	123.71	118.30
1	G	2	ASP	CB-CG-OD2	5.81	123.53	118.30
1	E	139	ASP	CB-CG-OD2	5.80	123.52	118.30
1	A	3	ASP	CB-CG-OD2	5.79	123.51	118.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1176	0	1205	22	0
1	B	1176	0	1205	31	4
1	C	1176	0	1205	24	0
1	D	1176	0	1205	33	4
1	E	1176	0	1205	27	0
1	F	1176	0	1205	37	1
1	G	1176	0	1205	28	0
1	H	1176	0	1205	29	1
2	A	53	0	31	3	0
2	B	53	0	31	4	0
2	C	53	0	31	4	0
2	D	53	0	31	4	0
2	E	53	0	31	4	0
2	F	53	0	31	4	0
2	G	53	0	31	5	0
2	H	53	0	31	3	0
3	A	44	0	26	0	0
3	B	44	0	26	0	0
3	C	44	0	26	0	0
3	D	44	0	26	0	0
3	E	44	0	26	0	0
3	F	44	0	26	0	0
3	G	44	0	26	0	0
3	H	44	0	26	0	0
4	A	62	0	0	2	0
4	B	53	0	0	6	0
4	C	57	0	0	3	0
4	D	51	0	0	2	0
4	E	40	0	0	3	0
4	F	44	0	0	3	0
4	G	47	0	0	3	0
4	H	43	0	0	0	0
All	All	10581	0	10096	188	5

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 9.

The worst 5 of 188 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:112:CYS:HB2	4:A:1263:HOH:O	1.35	1.22
1:D:31:MSE:HE3	2:D:4200:FAD:H5'2	1.33	1.09

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:H:31:MSE:HE3	2:H:8200:FAD:H5'2	1.31	1.09
1:B:31:MSE:HE3	2:B:2200:FAD:H5'2	1.24	1.08
1:E:31:MSE:HE3	2:E:5200:FAD:H5'2	1.37	1.05

All (5) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:62:GLN:NE2	1:D:62:GLN:CG[1_455]	1.24	0.96
1:B:58:GLU:OE2	1:D:25:ASN:ND2[1_455]	1.84	0.36
1:B:62:GLN:NE2	1:D:62:GLN:CB[1_455]	2.04	0.16
1:F:62:GLN:OE1	1:H:62:GLN:NE2[1_456]	2.08	0.12
1:B:62:GLN:NE2	1:D:62:GLN:CD[1_455]	2.11	0.09

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	151/161 (94%)	149 (99%)	2 (1%)	0	100	100
1	B	151/161 (94%)	149 (99%)	2 (1%)	0	100	100
1	C	151/161 (94%)	148 (98%)	3 (2%)	0	100	100
1	D	151/161 (94%)	147 (97%)	3 (2%)	1 (1%)	22	18
1	E	151/161 (94%)	147 (97%)	4 (3%)	0	100	100
1	F	151/161 (94%)	147 (97%)	4 (3%)	0	100	100
1	G	151/161 (94%)	149 (99%)	2 (1%)	0	100	100
1	H	151/161 (94%)	149 (99%)	2 (1%)	0	100	100
All	All	1208/1288 (94%)	1185 (98%)	22 (2%)	1 (0%)	51	54

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	D	122	ASP

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	130/132 (98%)	123 (95%)	7 (5%)	22	20
1	B	130/132 (98%)	123 (95%)	7 (5%)	22	20
1	C	130/132 (98%)	123 (95%)	7 (5%)	22	20
1	D	130/132 (98%)	122 (94%)	8 (6%)	18	15
1	E	130/132 (98%)	122 (94%)	8 (6%)	18	15
1	F	130/132 (98%)	122 (94%)	8 (6%)	18	15
1	G	130/132 (98%)	123 (95%)	7 (5%)	22	20
1	H	130/132 (98%)	123 (95%)	7 (5%)	22	20
All	All	1040/1056 (98%)	981 (94%)	59 (6%)	20	18

5 of 59 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	D	100	LEU
1	E	90	VAL
1	H	73	GLN
1	D	116	ASN
1	E	1	MSE

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 12 such sidechains are listed below:

Mol	Chain	Res	Type
1	E	116	ASN
1	F	25	ASN
1	G	116	ASN
1	D	116	ASN
1	F	116	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

5.6 Ligand geometry ⓘ

16 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z > 2$	Counts	RMSZ	# $ Z > 2$
3	NAD	A	1201	-	42,48,48	1.77	3 (7%)	50,73,73	1.18	3 (6%)
2	FAD	F	6200	-	51,58,58	1.47	7 (13%)	60,89,89	1.67	7 (11%)
2	FAD	A	1200	-	51,58,58	1.54	6 (11%)	60,89,89	1.63	9 (15%)
2	FAD	E	5200	-	51,58,58	1.54	8 (15%)	60,89,89	1.61	8 (13%)
2	FAD	D	4200	-	51,58,58	1.55	8 (15%)	60,89,89	1.58	7 (11%)
3	NAD	B	2201	-	42,48,48	1.82	4 (9%)	50,73,73	1.21	4 (8%)
3	NAD	G	7201	-	42,48,48	1.73	3 (7%)	50,73,73	1.16	3 (6%)
2	FAD	B	2200	-	51,58,58	1.57	8 (15%)	60,89,89	1.60	7 (11%)
3	NAD	H	8201	-	42,48,48	1.83	4 (9%)	50,73,73	1.22	5 (10%)
2	FAD	G	7200	-	51,58,58	1.49	6 (11%)	60,89,89	1.55	8 (13%)
2	FAD	C	3200	-	51,58,58	1.49	9 (17%)	60,89,89	1.74	9 (15%)
2	FAD	H	8200	-	51,58,58	1.48	7 (13%)	60,89,89	1.55	8 (13%)
3	NAD	E	5201	-	42,48,48	1.69	3 (7%)	50,73,73	1.22	3 (6%)
3	NAD	C	3201	-	42,48,48	1.79	3 (7%)	50,73,73	1.21	3 (6%)

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	NAD	D	4201	-	42,48,48	1.73	3 (7%)	50,73,73	1.26	5 (10%)
3	NAD	F	6201	-	42,48,48	1.81	3 (7%)	50,73,73	1.26	5 (10%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	NAD	A	1201	-	-	6/26/62/62	0/5/5/5
2	FAD	F	6200	-	-	11/30/50/50	0/6/6/6
2	FAD	A	1200	-	-	11/30/50/50	0/6/6/6
2	FAD	E	5200	-	-	9/30/50/50	0/6/6/6
2	FAD	D	4200	-	-	10/30/50/50	0/6/6/6
3	NAD	B	2201	-	-	6/26/62/62	0/5/5/5
3	NAD	G	7201	-	-	6/26/62/62	0/5/5/5
2	FAD	B	2200	-	-	11/30/50/50	0/6/6/6
3	NAD	H	8201	-	-	6/26/62/62	0/5/5/5
2	FAD	G	7200	-	-	11/30/50/50	0/6/6/6
2	FAD	C	3200	-	-	9/30/50/50	0/6/6/6
2	FAD	H	8200	-	-	11/30/50/50	0/6/6/6
3	NAD	E	5201	-	-	6/26/62/62	0/5/5/5
3	NAD	C	3201	-	-	6/26/62/62	0/5/5/5
3	NAD	D	4201	-	-	6/26/62/62	0/5/5/5
3	NAD	F	6201	-	-	6/26/62/62	0/5/5/5

The worst 5 of 85 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	F	6201	NAD	O7N-C7N	9.60	1.42	1.24
3	A	1201	NAD	O7N-C7N	9.32	1.42	1.24
3	G	7201	NAD	O7N-C7N	9.32	1.42	1.24
3	B	2201	NAD	O7N-C7N	9.17	1.41	1.24
3	H	8201	NAD	O7N-C7N	9.16	1.41	1.24

The worst 5 of 94 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	H	8200	FAD	N3A-C2A-N1A	-5.81	119.60	128.68
2	F	6200	FAD	N3A-C2A-N1A	-5.80	119.61	128.68
2	E	5200	FAD	N3A-C2A-N1A	-5.72	119.74	128.68
3	E	5201	NAD	N3A-C2A-N1A	-5.70	119.77	128.68
2	B	2200	FAD	N3A-C2A-N1A	-5.68	119.80	128.68

There are no chirality outliers.

5 of 131 torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
3	A	1201	NAD	C5B-O5B-PA-O1A
3	A	1201	NAD	C5B-O5B-PA-O2A
2	F	6200	FAD	C5B-O5B-PA-O3P
2	F	6200	FAD	C5'-O5'-P-O1P
2	F	6200	FAD	C5'-O5'-P-O2P


There are no ring outliers.


8 monomers are involved in 31 short contacts:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	F	6200	FAD	4	0
2	A	1200	FAD	3	0
2	E	5200	FAD	4	0
2	D	4200	FAD	4	0
2	B	2200	FAD	4	0
2	G	7200	FAD	5	0
2	C	3200	FAD	4	0
2	H	8200	FAD	3	0


The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.












































5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	147/161 (91%)	0.30	2 (1%) 75 78	18, 33, 48, 70	0
1	B	147/161 (91%)	0.36	4 (2%) 54 60	18, 33, 47, 71	0
1	C	147/161 (91%)	0.32	2 (1%) 75 78	18, 33, 47, 70	0
1	D	147/161 (91%)	0.33	4 (2%) 54 60	18, 33, 47, 71	0
1	E	147/161 (91%)	0.35	4 (2%) 54 60	18, 33, 48, 71	0
1	F	147/161 (91%)	0.36	5 (3%) 45 51	18, 33, 48, 71	0
1	G	147/161 (91%)	0.32	6 (4%) 37 43	18, 33, 48, 70	0
1	H	147/161 (91%)	0.30	8 (5%) 25 31	18, 33, 48, 72	0
All	All	1176/1288 (91%)	0.33	35 (2%) 50 56	18, 33, 48, 72	0

The worst 5 of 35 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	H	88	LYS	3.7
1	D	88	LYS	3.4
1	E	153	GLN	3.2
1	B	88	LYS	3.0
1	G	3	ASP	2.8

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates ⓘ

There are no carbohydrates in this entry.

6.4 Ligands




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	FAD	B	2200	53/53	0.89	0.15	30,39,48,50	0
2	FAD	D	4200	53/53	0.90	0.15	31,39,48,50	0
3	NAD	H	8201	44/44	0.90	0.16	37,42,48,49	0
2	FAD	C	3200	53/53	0.91	0.14	30,39,48,50	0
2	FAD	F	6200	53/53	0.92	0.14	30,39,48,50	0
2	FAD	E	5200	53/53	0.92	0.13	30,39,48,50	0
2	FAD	H	8200	53/53	0.92	0.14	31,39,48,50	0
3	NAD	D	4201	44/44	0.92	0.15	38,42,48,49	0
3	NAD	B	2201	44/44	0.93	0.14	37,42,48,49	0
3	NAD	F	6201	44/44	0.93	0.14	37,42,48,49	0
2	FAD	A	1200	53/53	0.94	0.12	30,39,48,50	0
3	NAD	C	3201	44/44	0.94	0.13	37,42,48,48	0
2	FAD	G	7200	53/53	0.95	0.12	30,39,48,50	0
3	NAD	A	1201	44/44	0.95	0.12	37,41,48,49	0
3	NAD	E	5201	44/44	0.95	0.13	37,42,48,48	0
3	NAD	G	7201	44/44	0.96	0.13	37,42,48,49	0




The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

Electron density around FAD B 2200:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)




**Electron density around FAD D 4200:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around NAD H 8201:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around FAD C 3200:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)




**Electron density around FAD F 6200:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around FAD E 5200:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)




**Electron density around FAD H 8200:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around NAD D 4201:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around NAD B 2201:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around NAD F 6201:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around FAD A 1200:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around NAD C 3201:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around FAD G 7200:

$2mF_o - DF_c$ (at 0.7 rmsd) in gray
 $mF_o - DF_c$ (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around NAD A 1201:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around NAD E 5201:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



Electron density around NAD G 7201:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers [i](#)

There are no such residues in this entry.