



Full wwPDB X-ray Structure Validation Report ⓘ

May 22, 2020 – 01:19 pm BST

PDB ID : 4S15
Title : Crystal structure of the orphan nuclear receptor RORalpha ligand-binding domain in complex with 4alpha-carboxyl, 4beta-methyl-zymosterol (4ACD8)
Authors : Huang, P.; Santori, F.R.; Littman, D.R.; Rastinejad, F.
Deposited on : 2015-01-08
Resolution : 1.90 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.11
buster-report : 1.1.7 (2018)
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.11


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.90 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	6207 (1.90-1.90)
Clashscore	141614	6847 (1.90-1.90)
Ramachandran outliers	138981	6760 (1.90-1.90)
Sidechain outliers	138945	6760 (1.90-1.90)
RSRZ outliers	127900	6082 (1.90-1.90)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	256	<div> <div style="width: 100%; height: 10px; background-color: red;"></div> <div style="display: flex; justify-content: space-between; align-items: center;"> % <div style="width: 87%; height: 10px; background-color: green;"></div> <div style="width: 8%; height: 10px; background-color: yellow;"></div> <div style="width: 5%; height: 10px; background-color: grey;"></div> </div> <div style="display: flex; justify-content: space-between; align-items: center;"> 87% 8% 5% </div> </div>
1	B	256	<div> <div style="width: 100%; height: 10px; background-color: red;"></div> <div style="display: flex; justify-content: space-between; align-items: center;"> 3% <div style="width: 86%; height: 10px; background-color: green;"></div> <div style="width: 11%; height: 10px; background-color: yellow;"></div> <div style="width: 4%; height: 10px; background-color: grey;"></div> </div> <div style="display: flex; justify-content: space-between; align-items: center;"> 86% 11% • </div> </div>
2	C	12	<div> <div style="width: 100%; height: 10px; background-color: red;"></div> <div style="display: flex; justify-content: space-between; align-items: center;"> 8% <div style="width: 75%; height: 10px; background-color: green;"></div> <div style="width: 25%; height: 10px; background-color: grey;"></div> </div> <div style="display: flex; justify-content: space-between; align-items: center;"> 75% 25% </div> </div>
2	D	12	<div> <div style="width: 100%; height: 10px; background-color: red;"></div> <div style="display: flex; justify-content: space-between; align-items: center;"> <div style="width: 75%; height: 10px; background-color: green;"></div> <div style="width: 25%; height: 10px; background-color: grey;"></div> </div> <div style="display: flex; justify-content: space-between; align-items: center;"> 75% 25% </div> </div>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit crite-

ria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
4	GOL	B	602	-	-	X	-

2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 4630 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Nuclear receptor ROR-alpha.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	244	Total	C	N	O	S	0	8	0
			2035	1311	337	371	16			
1	B	250	Total	C	N	O	S	0	5	0
			2069	1331	341	380	17			


There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	268	GLY	-	EXPRESSION TAG	UNP P35398
B	268	GLY	-	EXPRESSION TAG	UNP P35398

- Molecule 2 is a protein called Nuclear receptor-interacting protein 1.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
2	C	9	Total	C	N	O	0	0	0
			70	47	12	11			
2	D	9	Total	C	N	O	0	0	0
			70	47	12	11			

- Molecule 3 is (3beta,4alpha,5beta,14beta)-3-hydroxy-4-methylcholesta-8,24-diene-4-carboxylic acid (three-letter code: 4D8) (formula: C₂₉H₄₆O₃).



- GOL
-
- The diagram shows the skeletal structure of 1,2,3-propanetriol (glycerol). The three carbon atoms are labeled C1, C2, and C3 in green. C1 is on the left, C2 is in the middle, and C3 is on the right. Each carbon is bonded to a hydroxyl group (OH) in red. The hydroxyl group on C1 is labeled O1, the one on C2 is labeled O2, and the one on C3 is labeled O3. The bonds between the carbons and the carbons to their respective hydroxyl groups are shown as single lines.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	1	Total C O 6 3 3	0	0
4	B	1	Total C O 6 3 3	0	0


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	166	Total 166	O 166	0	0
5	C	1	Total 1	O 1	0	0
5	B	142	Total 142	O 142	0	0
5	D	1	Total 1	O 1	0	0



3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Nuclear receptor ROR-alpha




- Molecule 1: Nuclear receptor ROR-alpha



- Molecule 2: Nuclear receptor-interacting protein 1



- Molecule 2: Nuclear receptor-interacting protein 1



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	71.19Å 80.31Å 113.10Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	46.24 – 1.90 48.19 – 1.90	Depositor EDS
% Data completeness (in resolution range)	99.2 (46.24-1.90) 99.2 (48.19-1.90)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	0.11	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.19 (at 1.90Å)	Xtriage
Refinement program	PHENIX (phenix.refine: 1.9_1692)	Depositor
R, R_{free}	0.174 , 0.213 0.176 , 0.215	Depositor DCC
R_{free} test set	2641 reflections (5.12%)	wwPDB-VP
Wilson B-factor (Å ²)	24.9	Xtriage
Anisotropy	0.429	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.35 , 47.7	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	4630	wwPDB-VP
Average B, all atoms (Å ²)	33.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 7.00% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: GOL, 4D8

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.29	0/2098	0.41	0/2824
1	B	0.28	0/2125	0.40	0/2861
2	C	0.23	0/70	0.40	0/94
2	D	0.25	0/70	0.34	0/94
All	All	0.28	0/4363	0.40	0/5873

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2035	0	2064	14	0
1	B	2069	0	2078	18	0
2	C	70	0	79	0	0
2	D	70	0	79	0	0
3	A	32	0	45	2	0
3	B	32	0	45	4	0
4	A	6	0	8	3	0
4	B	6	0	8	4	0
5	A	166	0	0	5	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
5	B	142	0	0	1	0
5	C	1	0	0	0	0
5	D	1	0	0	0	0
All	All	4630	0	4406	35	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (35) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:367:ARG:HB3	4:B:602:GOL:H32	1.57	0.84
1:A:367:ARG:HB3	4:A:602:GOL:H32	1.71	0.73
1:B:316:ARG:NH1	5:B:770:HOH:O	2.23	0.71
1:A:290:TYR:H	4:A:602:GOL:HO1	1.38	0.69
1:A:288[B]:CYS:SG	5:A:828:HOH:O	2.52	0.66
1:B:311:TYR:OH	1:B:382:ASP:OD2	2.24	0.55
1:A:367:ARG:HB3	4:A:602:GOL:C3	2.39	0.52
1:B:430[A]:SER:OG	1:B:432:ASP:OD1	2.26	0.51
3:A:601:4D8:H31	3:A:601:4D8:H29	1.92	0.50
1:A:347:LEU:HD11	1:A:429:MET:HE1	1.94	0.49
1:A:457:LEU:HD22	1:A:466:ILE:HG12	1.95	0.49
1:A:412[A]:SER:OG	5:A:755:HOH:O	2.20	0.49
1:B:290:TYR:H	4:B:602:GOL:HO1	1.59	0.48
3:A:601:4D8:H32	5:A:764:HOH:O	2.12	0.48
1:B:370:ARG:HH11	4:B:602:GOL:H31	1.80	0.47
1:A:284:HIS:NE2	1:A:288[A]:CYS:SG	2.89	0.45
1:B:368:MET:HG3	3:B:601:4D8:H27	1.97	0.45
1:A:371:ALA:HB1	1:A:380:TYR:HB3	1.97	0.45
1:B:328:THR:O	1:B:332:GLN:HG3	2.16	0.45
1:A:499:VAL:O	1:A:508:LYS:HD3	2.17	0.44
1:B:467:LEU:O	1:B:471:ILE:HG12	2.18	0.44
3:B:601:4D8:H29	3:B:601:4D8:H31	1.98	0.44
1:B:338:ALA:HB2	1:B:428:LEU:HD21	1.99	0.43
1:A:289:GLN:HB3	5:A:740:HOH:O	2.18	0.43
1:A:504:PRO:HA	1:A:505:PRO:HD3	1.94	0.43
1:B:282:LYS:HE2	1:B:286:GLU:OE2	2.19	0.43
1:A:336:GLU:OE2	5:A:784:HOH:O	2.21	0.42
1:B:289:GLN:HG2	1:B:333:TYR:CD2	2.55	0.42
1:B:370:ARG:NH1	4:B:602:GOL:H31	2.35	0.42
1:B:505:PRO:O	1:B:509[A]:GLU:HG3	2.20	0.42

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:482:GLY:O	1:B:486:GLU:HG3	2.20	0.42
1:B:320:TRP:CH2	3:B:601:4D8:H44	2.56	0.41
1:B:489:MET:HE1	1:B:512:THR:HA	2.02	0.41
1:B:290:TYR:HE2	3:B:601:4D8:OC2	2.04	0.41
1:A:469:LYS:O	1:A:473:LYS:HG2	2.21	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	250/256 (98%)	246 (98%)	4 (2%)	0	100	100
1	B	253/256 (99%)	251 (99%)	2 (1%)	0	100	100
2	C	7/12 (58%)	7 (100%)	0	0	100	100
2	D	7/12 (58%)	7 (100%)	0	0	100	100
All	All	517/536 (96%)	511 (99%)	6 (1%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	230/231 (100%)	228 (99%)	2 (1%)	78	79

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	B	232/231 (100%)	229 (99%)	3 (1%)	69	68
2	C	8/11 (73%)	8 (100%)	0	100	100
2	D	8/11 (73%)	8 (100%)	0	100	100
All	All	478/484 (99%)	473 (99%)	5 (1%)	73	76

All (5) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	315	GLN
1	A	369	CYS
1	B	292	ARG
1	B	369	CYS
1	B	445	LEU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
1	B	289	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

5.6 Ligand geometry ⓘ

4 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The

Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	4D8	B	601	-	32,35,35	1.61	6 (18%)	48,56,56	1.71	11 (22%)
4	GOL	B	602	-	5,5,5	0.40	0	5,5,5	0.60	0
3	4D8	A	601	-	32,35,35	0.83	1 (3%)	48,56,56	1.33	5 (10%)
4	GOL	A	602	-	5,5,5	0.46	0	5,5,5	0.47	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	4D8	B	601	-	-	1/10/82/82	0/4/4/4
4	GOL	B	602	-	-	1/4/4/4	-
3	4D8	A	601	-	-	1/10/82/82	0/4/4/4
4	GOL	A	602	-	-	4/4/4/4	-

All (7) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	B	601	4D8	C8-C9	-5.45	1.28	1.34
3	B	601	4D8	C11-C9	-3.26	1.45	1.51
3	B	601	4D8	C14-C8	-3.15	1.47	1.51
3	B	601	4D8	C13-C14	-2.69	1.51	1.56
3	A	601	4D8	C13-C14	-2.53	1.51	1.56
3	B	601	4D8	C10-C5	-2.40	1.52	1.56
3	B	601	4D8	C7-C8	2.11	1.54	1.50

All (16) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	B	601	4D8	C7-C6-C5	-4.34	103.86	110.87
3	B	601	4D8	C14-C8-C9	-4.34	116.21	122.44
3	A	601	4D8	C14-C8-C9	-3.90	116.85	122.44
3	B	601	4D8	C1-C10-C5	3.20	111.99	108.34
3	A	601	4D8	C7-C6-C5	-3.19	105.72	110.87
3	B	601	4D8	C5-C10-C9	-3.07	103.27	108.39
3	B	601	4D8	C11-C9-C8	-2.94	118.43	120.73

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	B	601	4D8	C2-C3-C4	2.81	115.97	112.33
3	B	601	4D8	C13-C14-C8	-2.50	109.30	113.24
3	A	601	4D8	C1-C10-C5	2.35	111.02	108.34
3	B	601	4D8	O3-C3-C4	-2.31	104.64	110.61
3	B	601	4D8	C12-C13-C17	2.30	120.02	116.57
3	A	601	4D8	C27-C25-C26	2.29	119.67	114.60
3	B	601	4D8	C27-C25-C26	2.25	119.58	114.60
3	B	601	4D8	C5-C4-C3	2.21	111.98	107.95
3	A	601	4D8	C10-C9-C8	-2.12	121.38	123.16

There are no chirality outliers.

All (7) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	A	602	GOL	O1-C1-C2-C3
4	A	602	GOL	C1-C2-C3-O3
4	A	602	GOL	O1-C1-C2-O2
4	B	602	GOL	O1-C1-C2-C3
4	A	602	GOL	O2-C2-C3-O3
3	B	601	4D8	C20-C22-C23-C24
3	A	601	4D8	C20-C22-C23-C24



There are no ring outliers.

4 monomers are involved in 13 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	B	601	4D8	4	0
4	B	602	GOL	4	0
3	A	601	4D8	2	0
4	A	602	GOL	3	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient

equivalents in the CSD to analyse the geometry.



5.7 Other polymers

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	244/256 (95%)	-0.04	2 (0%) 86 87	16, 26, 54, 76	0
1	B	250/256 (97%)	0.19	8 (3%) 47 50	17, 30, 64, 107	0
2	C	9/12 (75%)	0.01	1 (11%) 5 6	25, 28, 43, 49	0
2	D	9/12 (75%)	-0.05	0 100 100	24, 29, 45, 49	0
All	All	512/536 (95%)	0.07	11 (2%) 63 66	16, 28, 58, 107	0

All (11) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	464	ASP	5.7
1	B	461	HIS	4.7
1	B	463	GLU	4.1
1	B	270	MET	3.6
1	A	466	ILE	3.0
1	B	462	ARG	2.6
1	B	346	GLU	2.4
1	B	269	SER	2.3
2	C	498	THR	2.2
1	A	288[A]	CYS	2.1
1	B	445	LEU	2.1

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates ⓘ


There are no carbohydrates in this entry.


6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q<0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
3	4D8	B	601	32/32	0.78	0.31	41,51,89,94	0
3	4D8	A	601	32/32	0.82	0.24	34,41,76,83	0
4	GOL	A	602	6/6	0.84	0.35	45,52,56,63	0
4	GOL	B	602	6/6	0.85	0.31	41,46,60,75	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.





6.5 Other polymers [i](#)

There are no such residues in this entry.