



# Full wwPDB NMR Structure Validation Report ⓘ

May 28, 2020 – 08:27 pm BST

PDB ID : 1S37  
Title : Accomodation of Mispair-Aligned N3T-Ethyl-N3T DNA Interstrand Crosslink  
Authors : da Silva, M.W.; Noronha, A.M.; Noll, D.M.; Miller, P.S.; Colvin, O.M.; Gamcsik, M.P.  
Deposited on : 2004-01-12

This is a Full wwPDB NMR Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/NMRValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

Cyrange	:	Kirchner and Güntert (2011)
NmrClust	:	Kelley et al. (1996)
MolProbity	:	4.02b-467
Mogul	:	1.8.5 (274361), CSD as541be (2020)
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
RCI	:	v_1n_11_5_13_A (Berjanski et al., 2005)
PANAV	:	Wang et al. (2010)
ShiftChecker	:	2.11
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.11


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*SOLUTION NMR*



The overall completeness of chemical shifts assignment was not calculated.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	NMR archive (#Entries)
Clashscore	158937	12864

The table below summarises the geometric issues observed across the polymeric chains and their fit to the experimental data. The red, orange, yellow and green segments indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A cyan segment indicates the fraction of residues that are not part of the well-defined cores, and a grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$

Mol	Chain	Length	Quality of chain
1	A	11	 18% 82%
2	B	11	 9% 91%

## 2 Ensemble composition and analysis ⓘ

This entry contains 1 models. Identification of well-defined residues and clustering analysis are not possible.

### 3 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 702 atoms, of which 256 are hydrogens and 0 are deuteriums.

- Molecule 1 is a DNA chain called DNA (5'-D(\*CP\*GP\*AP\*AP\*AP\*(TTM)P\*TP\*TP\*TP\*CP\*G)-3').

Mol	Chain	Residues	Atoms						Trace
			Total	C	H	N	O	P	
1	A	11	354	110	130	39	65	10	0

- Molecule 2 is a DNA chain called DNA (5'-D(\*CP\*GP\*AP\*AP\*AP\*TP\*TP\*TP\*TP\*CP\*G)-3').

Mol	Chain	Residues	Atoms						Trace
			Total	C	H	N	O	P	
2	B	11	348	108	126	39	65	10	0

## 4 Residue-property plots

These plots are provided for all protein, RNA and DNA chains in the entry. The first graphic is the same as shown in the summary in section 1 of this report. The second graphic shows the sequence where residues are colour-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outliers are shown as green connectors. Residues which are classified as ill-defined in the NMR ensemble, are shown in cyan with an underline colour-coded according to the previous scheme. Residues which were present in the experimental sample, but not modelled in the final structure are shown in grey.

- Molecule 1: DNA (5'-D(\*CP\*GP\*AP\*AP\*AP\*(TTM)P\*TP\*TP\*TP\*CP\*G)-3')

Chain A:  18% 82%



- Molecule 2: DNA (5'-D(\*CP\*GP\*AP\*AP\*AP\*TP\*TP\*TP\*TP\*CP\*G)-3')

Chain B:  9% 91%



## 5 Refinement protocol and experimental data overview [i](#)

The models were refined using the following method: *distance geometry, simulated annealing, molecular dynamics*.

Of the 8 calculated structures, 1 were deposited, based on the following criterion: *structures with the lowest energy*.

The following table shows the software used for structure solution, optimisation and refinement.

Software name	Classification	Version
X-PLOR	refinement	3.2

No chemical shift data was provided. No validations of the models with respect to experimental NMR restraints is performed at this time.

COVALENT-GEOMETRY INFOmissingINFO

### 5.1 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in each chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes averaged over the ensemble.

Mol	Chain	Non-H	H(model)	H(added)	Clashes
1	A	224	130	127	13
2	B	222	126	126	13
All	All	446	256	253	26

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 37.

All clashes are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Clash(Å)	Distance(Å)
2:B:6:DT:H2'	2:B:7:DT:H71	0.58	1.75
2:B:5:DA:C2	2:B:6:DT:C2	0.56	2.93
1:A:7:DT:H2''	1:A:8:DT:H72	0.56	1.77
2:B:7:DT:H2''	2:B:8:DT:H72	0.54	1.80
1:A:5:DA:C2'	1:A:6:TTM:H73	0.53	2.34
1:A:7:DT:H2''	1:A:8:DT:C7	0.51	2.35
1:A:5:DA:C2	1:A:6:TTM:C2	0.51	2.93
2:B:6:DT:OP1	2:B:6:DT:H4'	0.51	2.06
1:A:6:TTM:H4'	1:A:6:TTM:OP1	0.49	2.07
2:B:7:DT:H2''	2:B:8:DT:C7	0.49	2.36

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Clash(Å)	Distance(Å)
2:B:5:DA:C8	2:B:6:DT:H72	0.47	2.44
1:A:6:TTM:C6	1:A:7:DT:H72	0.46	2.40
1:A:4:DA:C2	1:A:5:DA:C5	0.45	3.05
2:B:4:DA:C2	2:B:5:DA:C5	0.45	3.04
2:B:10:DC:O5'	2:B:10:DC:H2'	0.44	2.11
1:A:10:DC:H2''	1:A:11:DG:C8	0.43	2.49
1:A:10:DC:O5'	1:A:10:DC:H2'	0.43	2.13
2:B:5:DA:C4	2:B:6:DT:C5	0.42	3.07
1:A:5:DA:C2	1:A:6:TTM:N3	0.42	2.87
2:B:5:DA:C2	2:B:6:DT:N3	0.42	2.87
2:B:9:DT:C2	2:B:10:DC:C5	0.41	3.08
2:B:10:DC:H2''	2:B:11:DG:C8	0.41	2.50
1:A:9:DT:C2	1:A:10:DC:C5	0.41	3.08
1:A:1:DC:C2	1:A:2:DG:C5	0.40	3.09
2:B:1:DC:C2	2:B:2:DG:C5	0.40	3.09
1:A:7:DT:C2'	1:A:8:DT:H72	0.40	2.45

## 5.2 Torsion angles [i](#)

### 5.2.1 Protein backbone [i](#)

There are no protein molecules in this entry.

### 5.2.2 Protein sidechains [i](#)

There are no protein molecules in this entry.

### 5.2.3 RNA [i](#)

There are no RNA molecules in this entry.

MODRES-GEOMETRY INFOmissingINFO

## 5.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 5.4 Ligand geometry [i](#)

There are no ligands in this entry.

## 5.5 Other polymers [i](#)

There are no such molecules in this entry.

## 5.6 Polymer linkage issues [i](#)

There are no chain breaks in this entry.



## 6 Chemical shift validation

No chemical shift data were provided