



# Full wwPDB X-ray Structure Validation Report ⓘ

May 24, 2020 – 01:20 am BST

PDB ID : 4S30  
Title : ERK2 intrinsically active mutant (I84A)  
Authors : Livnah, O.; Karamansha, Y.  
Deposited on : 2015-01-26  
Resolution : 2.00 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Xtriage (Phenix)	:	1.13
EDS	:	2.11
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac	:	5.8.0158
CCP4	:	7.0.044 (Gargrove)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.11


# 1 Overall quality at a glance ⓘ

The following experimental techniques were used to determine the structure:

## *X-RAY DIFFRACTION*

The reported resolution of this entry is 2.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	8085 (2.00-2.00)
Clashscore	141614	9178 (2.00-2.00)
Ramachandran outliers	138981	9054 (2.00-2.00)
Sidechain outliers	138945	9053 (2.00-2.00)
RSRZ outliers	127900	7900 (2.00-2.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	358	<div> <div>8%</div> <div>73%</div> <div>19%</div> <div>...</div> </div>

## 2 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 2856 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Mitogen-activated protein kinase 1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	342	Total	C	N	O	S	0	0	0
			2774	1781	476	503	14			

There is a discrepancy between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	84	ALA	ILE	ENGINEERED MUTATION	UNP P63086


- Molecule 2 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	82	Total	O	0	0
			82	82		

### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Mitogen-activated protein kinase 1



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	49.02Å 69.54Å 59.80Å 90.00° 109.06° 90.00°	Depositor
Resolution (Å)	28.64 – 2.00 28.64 – 2.00	Depositor EDS
% Data completeness (in resolution range)	99.5 (28.64-2.00) 99.6 (28.64-2.00)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.57 (at 2.00Å)	Xtriage
Refinement program	REFMAC 5.7.0029	Depositor
R, $R_{free}$	0.194 , 0.249 0.201 , 0.243	Depositor DCC
$R_{free}$ test set	1306 reflections (5.07%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	33.0	Xtriage
Anisotropy	0.069	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.37 , 51.1	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.50$ , $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.95	EDS
Total number of atoms	2856	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	44.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 7.49% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality

### 5.1 Standard geometry

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.77	1/2838 (0.0%)	0.94	7/3842 (0.2%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	211	SER	CB-OG	-5.23	1.35	1.42

All (7) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	89	ARG	NE-CZ-NH1	12.21	126.40	120.30
1	A	89	ARG	NE-CZ-NH2	-8.83	115.88	120.30
1	A	86	ASP	CB-CG-OD1	5.72	123.45	118.30
1	A	110	LEU	CB-CG-CD2	5.56	120.45	111.00
1	A	189	ARG	NE-CZ-NH1	5.42	123.01	120.30
1	A	216	LEU	CB-CG-CD2	5.21	119.86	111.00
1	A	146	ARG	NE-CZ-NH2	-5.20	117.70	120.30

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	29	ILE	Peptide

## 5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2774	0	2776	47	0
2	A	82	0	0	6	0
All	All	2856	0	2776	47	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 8.

All (47) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:33:ALA:O	1:A:35:GLY:N	1.99	0.94
1:A:78:HIS:HD2	1:A:80:ASN:H	1.37	0.70
1:A:237:HIS:HD2	2:A:448:HOH:O	1.76	0.68
1:A:51:ILE:HD12	1:A:102:VAL:HG22	1.77	0.67
1:A:107:GLU:HG3	1:A:157:THR:HG23	1.79	0.65
1:A:78:HIS:CD2	1:A:80:ASN:H	2.14	0.64
1:A:330:ASP:C	1:A:331:MET:HE3	2.16	0.64
1:A:45:ASN:O	1:A:47:VAL:HG13	1.98	0.62
1:A:42:ASP:C	1:A:42:ASP:OD1	2.38	0.62
1:A:68:ARG:HG3	1:A:330:ASP:HB3	1.83	0.61
1:A:185:TYR:CE2	1:A:187:ALA:HB3	2.37	0.60
1:A:185:TYR:HE2	1:A:187:ALA:HB3	1.70	0.57
1:A:89:ARG:HD3	1:A:349:THR:OG1	2.08	0.54
1:A:170:ARG:HG3	1:A:331:MET:CG	2.38	0.53
1:A:51:ILE:CD1	1:A:102:VAL:HG22	2.38	0.53
1:A:272:PRO:HG2	1:A:275:ARG:HD2	1.90	0.53
1:A:170:ARG:HG3	1:A:331:MET:HG3	1.90	0.52
1:A:24:THR:HG1	1:A:43:ASN:HD21	1.57	0.51
1:A:107:GLU:CG	1:A:157:THR:HG23	2.40	0.50
1:A:230:HIS:HD2	2:A:427:HOH:O	1.94	0.50
1:A:272:PRO:HG2	1:A:275:ARG:CD	2.42	0.49
1:A:22:ARG:O	1:A:22:ARG:HG3	2.12	0.49
1:A:107:GLU:N	1:A:107:GLU:CD	2.64	0.49
1:A:325:ALA:HB1	1:A:326:PRO:HD2	1.94	0.48
1:A:330:ASP:O	1:A:332:GLU:OE2	2.32	0.48

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:330:ASP:CA	1:A:331:MET:HE3	2.45	0.47
1:A:110:LEU:HD22	1:A:114:LEU:HG	1.97	0.46
1:A:300:ILE:CG1	1:A:304:GLN:HB2	2.46	0.45
1:A:64:GLN:NE2	1:A:332:GLU:O	2.50	0.45
1:A:171:VAL:H	1:A:331:MET:CE	2.31	0.43
1:A:351:ARG:HD2	2:A:453:HOH:O	2.17	0.43
1:A:24:THR:OG1	1:A:43:ASN:ND2	2.37	0.43
1:A:28:TYR:CE1	1:A:30:GLY:HA3	2.54	0.43
1:A:124:ILE:HD13	1:A:220:LEU:HD23	2.01	0.43
1:A:89:ARG:NH2	2:A:436:HOH:O	2.42	0.43
1:A:121:ASN:ND2	2:A:425:HOH:O	2.52	0.42
1:A:254:ILE:HG22	2:A:461:HOH:O	2.18	0.42
1:A:107:GLU:HG3	1:A:157:THR:CG2	2.45	0.42
1:A:136:LYS:HE2	1:A:322:ILE:HG13	2.00	0.42
1:A:107:GLU:CD	1:A:107:GLU:H	2.24	0.42
1:A:225:ILE:HG23	1:A:226:PHE:CD2	2.55	0.41
1:A:272:PRO:CG	1:A:275:ARG:HD2	2.51	0.41
1:A:216:LEU:HD13	1:A:288:LEU:HB2	2.01	0.41
1:A:255:ASN:HD22	1:A:255:ASN:C	2.24	0.41
1:A:42:ASP:OD1	1:A:44:LEU:N	2.54	0.41
1:A:330:ASP:N	1:A:331:MET:HE3	2.36	0.40
1:A:30:GLY:HA2	1:A:37:VAL:H	1.86	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	334/358 (93%)	317 (95%)	12 (4%)	5 (2%)	<b>10</b>   <b>4</b>

All (5) Ramachandran outliers are listed below:



Mol	Chain	Res	Type
1	A	330	ASP
1	A	255	ASN
1	A	5	ALA
1	A	29	ILE
1	A	56	PRO

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	304/317 (96%)	272 (90%)	32 (10%)	7 4

All (32) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	24	THR
1	A	26	LEU
1	A	27	SER
1	A	42	ASP
1	A	44	LEU
1	A	59	HIS
1	A	60	GLN
1	A	65	ARG
1	A	67	LEU
1	A	75	ARG
1	A	89	ARG
1	A	107	GLU
1	A	110	LEU
1	A	113	LEU
1	A	117	GLN
1	A	142	ASN
1	A	155	LEU
1	A	170	ARG
1	A	199	ASN
1	A	207	ILE
1	A	216	LEU
1	A	248	GLU

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	A	250	LEU
1	A	255	ASN
1	A	256	LEU
1	A	262	LEU
1	A	265	LEU
1	A	268	LYS
1	A	303	GLU
1	A	309	PRO
1	A	322	ILE
1	A	344	LEU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (14) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	45	ASN
1	A	60	GLN
1	A	78	HIS
1	A	85	ASN
1	A	117	GLN
1	A	121	ASN
1	A	152	ASN
1	A	199	ASN
1	A	230	HIS
1	A	237	HIS
1	A	251	ASN
1	A	255	ASN
1	A	260	ASN
1	A	279	ASN

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

### 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

### 5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

## 5.6 Ligand geometry

There are no ligands in this entry.

## 5.7 Other polymers

There are no such residues in this entry.

## 5.8 Polymer linkage issues

There are no chain breaks in this entry.

## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	342/358 (95%)	0.20	28 (8%) <b>11</b> <b>11</b>	20, 40, 78, 112	0

All (28) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	59	HIS	6.3
1	A	355	GLY	5.6
1	A	62	TYR	5.5
1	A	8	GLY	5.1
1	A	329	PHE	4.5
1	A	186	VAL	4.3
1	A	44	LEU	4.1
1	A	60	GLN	4.1
1	A	47	VAL	3.6
1	A	15	GLN	3.3
1	A	338	LYS	3.0
1	A	28	TYR	3.0
1	A	32	GLY	2.9
1	A	46	LYS	2.7
1	A	58	GLU	2.7
1	A	16	VAL	2.6
1	A	328	LYS	2.6
1	A	176	HIS	2.6
1	A	31	GLU	2.6
1	A	212	VAL	2.5
1	A	41	TYR	2.5
1	A	61	THR	2.4
1	A	184	GLU	2.2
1	A	254	ILE	2.2
1	A	223	ARG	2.1
1	A	148	LEU	2.1
1	A	115	LYS	2.1

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	A	30	GLY	2.1

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 6.4 Ligands [i](#)

There are no ligands in this entry.

## 6.5 Other polymers [i](#)

There are no such residues in this entry.