



# Full wwPDB X-ray Structure Validation Report ⓘ

Aug 6, 2020 – 08:33 AM BST

PDB ID : 3S4A  
Title : Cellobiose phosphorylase from *Cellulomonas uda* in complex with cellobiose  
Authors : Van Hoorebeke, A.; Stout, J.; Soetaert, W.; Van Beeumen, J.; Desmet, T.; Savvides, S.  
Deposited on : 2011-05-19  
Resolution : 1.99 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.13.1  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.13.1


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

## *X-RAY DIFFRACTION*

The reported resolution of this entry is 1.99 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	8085 (2.00-2.00)
Clashscore	141614	9178 (2.00-2.00)
Ramachandran outliers	138981	9054 (2.00-2.00)
Sidechain outliers	138945	9053 (2.00-2.00)
RSRZ outliers	127900	7900 (2.00-2.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	822	<div> <div>5%</div> <div>89%</div> <div>11%</div> </div>
1	B	822	<div> <div>2%</div> <div>89%</div> <div>11%</div> </div>
2	C	2	<div> <div>100%</div> </div>
2	D	2	<div> <div>100%</div> </div>

## 2 Entry composition [i](#)

There are 3 unique types of molecules in this entry. The entry contains 26625 atoms, of which 12150 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Cellobiose phosphorylase.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	822	Total	C	H	N	O	S	0	3	0
			12471	4072	6026	1111	1245	17			
1	B	822	Total	C	H	N	O	S	9	5	0
			12559	4091	6082	1112	1257	17			

- Molecule 2 is an oligosaccharide called beta-D-glucopyranose-(1-4)-beta-D-glucopyranose.



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
2	C	2	Total	C	H	O	0	0	0
			44	12	21	11			
2	D	2	Total	C	H	O	0	0	0
			44	12	21	11			


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	745	Total	O	0	0
			745	745		
3	B	762	Total	O	0	0
			762	762		


### 3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

#### • Molecule 1: Cellobiose phosphorylase



#### • Molecule 1: Cellobiose phosphorylase



#### • Molecule 2: beta-D-glucopyranose-(1-4)-beta-D-glucopyranose



B6C1  
B6C2

- Molecule 2: beta-D-glucopyranose-(1-4)-beta-D-glucopyranose

Chain D:

100%

B6C1  
B6C2

## 4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	85.69Å 195.56Å 103.64Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	45.79 – 1.99 45.79 – 1.99	Depositor EDS
% Data completeness (in resolution range)	88.2 (45.79-1.99) 98.8 (45.79-1.99)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.64 (at 1.98Å)	Xtriage
Refinement program	PHENIX 1.6.4 _486	Depositor
R, $R_{free}$	0.200 , 0.255 0.209 , 0.254	Depositor DCC
$R_{free}$ test set	5954 reflections (5.01%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	10.0	Xtriage
Anisotropy	0.596	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.42 , 57.4	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.57$ , $\langle L^2 \rangle = 0.42$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.94	EDS
Total number of atoms	26625	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	14.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 61.30 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 1.3373e-05. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: BGC

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.80	1/6622 (0.0%)	0.70	3/9029 (0.0%)
1	B	0.80	0/6657	0.71	1/9075 (0.0%)
All	All	0.80	1/13279 (0.0%)	0.70	4/18104 (0.0%)

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	623	VAL	CB-CG2	5.18	1.63	1.52

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	789	ASP	CB-CG-OD1	8.56	126.01	118.30
1	A	789	ASP	CB-CG-OD1	5.31	123.08	118.30
1	A	792	ARG	NE-CZ-NH1	5.25	122.92	120.30
1	A	792	ARG	NE-CZ-NH2	-5.16	117.72	120.30

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	6445	6026	6050	56	0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	B	6477	6082	6087	67	0
2	C	23	21	21	0	0
2	D	23	21	21	0	0
3	A	745	0	0	3	0
3	B	762	0	0	6	0
All	All	14475	12150	12179	121	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (121) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:504:SER:O	1:B:508[B]:THR:HG22	1.80	0.82
1:A:51:LYS:HG2	1:A:52:MET:HG2	1.63	0.80
1:A:51:LYS:HE3	1:A:158:GLN:OE1	1.88	0.73
1:B:633:MET:H	1:B:668:ASN:HD21	1.37	0.70
1:B:503:GLU:HB3	1:B:508[A]:THR:HG21	1.74	0.69
1:A:31:GLU:O	1:A:34:PHE:CZ	2.46	0.69
1:B:715:ALA:O	1:B:722:HIS:HD2	1.76	0.68
1:B:198:TYR:CE2	1:B:286[B]:VAL:HG11	2.30	0.66
1:A:51:LYS:HD2	1:A:360:ILE:O	1.98	0.64
1:A:633:MET:H	1:A:668:ASN:HD21	1.47	0.63
1:B:28:LEU:HB2	1:B:35:SER:HB2	1.79	0.63
1:B:794:ARG:NH1	3:B:1217:HOH:O	2.33	0.62
1:A:347:PHE:HE2	1:A:383:ARG:NH2	1.97	0.62
1:B:198:TYR:CE2	1:B:286[B]:VAL:CG1	2.82	0.62
1:B:22:TYR:CG	1:B:23:PRO:HD2	2.34	0.61
1:A:322:TYR:CD2	1:A:772:ARG:HD3	2.35	0.61
1:A:668:ASN:N	1:A:669:PRO:CD	2.66	0.58
1:A:795:LEU:HD22	1:A:820:VAL:HG22	1.86	0.55
3:A:981:HOH:O	1:B:192:ARG:HD3	2.04	0.55
1:B:496:CYS:SG	1:B:508[B]:THR:HG23	2.45	0.55
1:B:191:TYR:CE2	1:B:286[B]:VAL:HG22	2.42	0.55
1:A:406:GLN:HE21	1:A:413:ASN:HB2	1.71	0.55
1:B:278:TRP:CD1	1:B:283:LYS:HG2	2.41	0.55
1:B:22:TYR:CD2	1:B:23:PRO:HD2	2.42	0.55
1:B:402:TYR:CG	1:B:412:GLY:HA3	2.41	0.54
1:A:795:LEU:HD23	1:A:820:VAL:HG13	1.88	0.54
1:A:429:ALA:HB2	1:A:528:TYR:CZ	2.44	0.52
1:B:539:ARG:HA	3:B:1497:HOH:O	2.08	0.52

*Continued on next page...*



*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:317:ASP:HB2	3:B:972:HOH:O	2.10	0.52
1:A:190:GLU:HG2	1:A:193:GLU:CD	2.31	0.51
1:B:633:MET:H	1:B:668:ASN:ND2	2.08	0.51
1:B:329:GLU:HG3	3:B:1329:HOH:O	2.09	0.51
1:A:402:TYR:CG	1:A:412:GLY:HA3	2.46	0.51
1:B:794:ARG:HH21	1:B:821:THR:HG22	1.75	0.51
1:A:743:GLN:HB3	1:A:749:ARG:HB3	1.93	0.51
1:A:485:ARG:HD3	1:A:506:GLN:O	2.11	0.51
1:A:571:ARG:HG2	1:A:591:TRP:CD2	2.47	0.50
1:A:593:GLU:N	1:A:594:PRO:CD	2.75	0.50
1:B:382:GLU:HG2	1:B:383:ARG:N	2.26	0.50
1:B:768:PHE:CE1	1:B:783:VAL:HG21	2.47	0.50
1:B:716:GLY:C	1:B:722:HIS:CD2	2.85	0.49
1:A:666:HIS:HB2	3:A:924:HOH:O	2.13	0.48
1:B:705:LEU:HD11	1:B:726:LYS:HA	1.95	0.48
1:A:771:THR:HG23	1:A:780:GLU:OE2	2.13	0.48
1:B:504:SER:O	1:B:508[A]:THR:HG23	2.13	0.48
1:A:52:MET:HA	1:A:156:ASN:HA	1.95	0.48
1:A:657:TYR:HB3	1:B:169:SER:HB3	1.96	0.48
1:A:101:LEU:HD22	1:A:341:TYR:CG	2.48	0.48
1:A:683:ARG:HD2	1:A:687:TYR:CZ	2.49	0.47
1:B:668:ASN:N	1:B:669:PRO:CD	2.77	0.47
1:B:716:GLY:O	1:B:722:HIS:HB2	2.14	0.47
1:B:136:ASN:ND2	1:B:142:LYS:HG2	2.30	0.47
1:B:1:MET:CE	3:B:1282:HOH:O	2.62	0.47
1:B:686:ASP:O	1:B:690:ARG:HG3	2.15	0.47
1:A:101:LEU:HD23	1:A:341:TYR:CD2	2.50	0.47
1:B:504:SER:O	1:B:508[B]:THR:CG2	2.59	0.46
1:B:53:ARG:HA	1:B:53:ARG:HD3	1.70	0.46
1:B:191:TYR:CD2	1:B:286[B]:VAL:HG22	2.51	0.46
1:B:405:TYR:O	1:B:407:PRO:HD3	2.16	0.46
1:A:210:PHE:O	1:A:237:ASN:HA	2.16	0.46
1:B:758:ASP:N	1:B:759:PRO:HD3	2.31	0.46
1:A:711:ALA:HA	1:A:728:SER:HA	1.98	0.45
1:B:693:PRO:HA	3:B:1357:HOH:O	2.16	0.45
1:B:483:ILE:HG21	1:B:494:LEU:HD12	1.99	0.45
1:A:703:HIS:O	1:A:704:ARG:HB2	2.16	0.45
1:B:795:LEU:HD22	1:B:820:VAL:HG22	1.99	0.44
1:A:483:ILE:HG21	1:A:494:LEU:HD12	1.98	0.44
1:B:656:GLY:HA3	1:B:719:ALA:HB2	1.99	0.44
1:B:210:PHE:O	1:B:237:ASN:HA	2.17	0.44

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:74:VAL:O	1:B:80:VAL:HA	2.17	0.44
1:B:224:LEU:HD23	1:B:224:LEU:HA	1.77	0.44
1:B:715:ALA:O	1:B:722:HIS:CD2	2.63	0.44
1:A:527:LEU:O	1:A:531:GLN:HG3	2.17	0.44
1:A:761:ILE:HG21	1:A:765:VAL:HB	1.99	0.44
1:B:448:VAL:HG12	1:B:463:HIS:CE1	2.53	0.44
1:B:22:TYR:CG	1:B:23:PRO:CD	3.01	0.43
1:A:705:LEU:HD11	1:A:726:LYS:HA	1.99	0.43
1:B:269:TYR:CE2	1:B:271:GLU:HG3	2.54	0.43
1:A:568:TRP:CZ2	1:A:583:ASP:HB2	2.53	0.43
1:A:191:TYR:CD2	1:A:286:VAL:HG12	2.54	0.43
1:B:728:SER:O	1:B:729:TRP:HB2	2.18	0.43
1:B:487:ASP:O	1:B:488:TRP:C	2.56	0.42
1:A:347:PHE:HE2	1:A:383:ARG:CZ	2.31	0.42
1:A:656:GLY:HA3	1:A:719:ALA:HB2	2.01	0.42
1:B:113:VAL:HG23	1:B:142:LYS:HE3	2.00	0.42
1:B:347:PHE:HB2	1:B:376:PHE:CD2	2.54	0.42
1:A:794:ARG:NH2	1:A:821:THR:HG22	2.35	0.42
1:B:496:CYS:SG	1:B:508[B]:THR:CG2	3.08	0.42
1:A:166:ARG:HD2	1:B:649:GLU:OE2	2.19	0.42
1:A:146:LEU:HD12	1:A:253:VAL:HG21	2.02	0.42
1:B:190:GLU:O	1:B:193:GLU:HG3	2.19	0.42
1:B:593:GLU:HB2	1:B:594:PRO:HD3	2.01	0.42
1:B:70:ARG:HD2	1:B:70:ARG:N	2.34	0.42
1:A:60:TYR:CZ	1:A:358:THR:HA	2.55	0.42
1:A:337:ILE:HG13	3:A:1260:HOH:O	2.20	0.42
1:A:458:VAL:HB	1:A:459:PRO:HD2	2.02	0.42
1:B:711:ALA:HA	1:B:728:SER:HA	2.02	0.41
1:A:229:VAL:HB	1:A:230:PRO:HD3	2.03	0.41
1:A:55:LEU:HD12	1:A:55:LEU:N	2.35	0.41
1:A:749:ARG:HA	1:A:750:PRO:HD3	1.73	0.41
1:B:113:VAL:CG2	1:B:142:LYS:HE3	2.50	0.41
1:A:52:MET:SD	1:A:156:ASN:ND2	2.94	0.41
1:A:535:LEU:HA	1:A:535:LEU:HD23	1.93	0.41
1:A:666:HIS:CE1	1:A:732:GLY:HA3	2.56	0.41
1:A:429:ALA:HB2	1:A:528:TYR:CE2	2.55	0.41
1:B:49:ASP:OD1	1:B:362:ARG:HD3	2.21	0.41
1:B:446:GLU:HA	1:B:447:PRO:HD3	1.95	0.41
1:B:665:CYS:O	1:B:669:PRO:HD3	2.21	0.41
1:A:101:LEU:CD2	1:A:341:TYR:CG	3.04	0.41
1:A:795:LEU:HD11	1:A:805:ASN:HA	2.03	0.41

*Continued on next page...*

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:757:VAL:HG11	1:B:820:VAL:HG21	2.02	0.41
1:A:380:ILE:HB	1:A:383:ARG:HD3	2.03	0.41
1:A:382:GLU:HG2	1:A:383:ARG:N	2.36	0.41
1:A:574:ASP:HB3	1:A:580[A]:ILE:HD11	2.03	0.40
1:A:672:ILE:HG12	1:A:687:TYR:HB2	2.03	0.40
1:B:175:VAL:HG21	1:B:210:PHE:CE2	2.56	0.40
1:B:193:GLU:CG	1:B:194:ARG:H	2.34	0.40
1:B:574:ASP:OD2	1:B:578:ASN:HB2	2.21	0.40
1:A:322:TYR:HD2	1:A:772:ARG:HD3	1.86	0.40
1:B:593:GLU:N	1:B:594:PRO:CD	2.84	0.40
1:B:6:PHE:CD1	1:B:333:ARG:HD3	2.56	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	823/822 (100%)	786 (96%)	37 (4%)	0	100	100
1	B	825/822 (100%)	786 (95%)	38 (5%)	1 (0%)	51	49
All	All	1648/1644 (100%)	1572 (95%)	75 (5%)	1 (0%)	51	49

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	85	TRP

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	667/675 (99%)	665 (100%)	2 (0%)	92	95
1	B	674/675 (100%)	672 (100%)	2 (0%)	92	95
All	All	1341/1350 (99%)	1337 (100%)	4 (0%)	92	95

All (4) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	543	ASP
1	A	597	PHE
1	B	597	PHE
1	B	772	ARG

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (5) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	406	GLN
1	A	668	ASN
1	B	668	ASN
1	B	722	HIS
1	B	760	GLN

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates ⓘ

4 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond

length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
2	BGC	C	1	2	12,12,12	1.12	2 (16%)	17,17,17	1.27	1 (5%)
2	BGC	C	2	2	11,11,12	0.98	0	15,15,17	1.64	3 (20%)
2	BGC	D	1	2	12,12,12	0.96	0	17,17,17	1.05	1 (5%)
2	BGC	D	2	2	11,11,12	1.07	1 (9%)	15,15,17	1.97	5 (33%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	BGC	C	1	2	-	1/2/22/22	0/1/1/1
2	BGC	C	2	2	-	2/2/19/22	0/1/1/1
2	BGC	D	1	2	-	0/2/22/22	0/1/1/1
2	BGC	D	2	2	-	1/2/19/22	0/1/1/1

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	C	1	BGC	O5-C5	-2.29	1.38	1.44
2	D	2	BGC	O5-C5	-2.10	1.39	1.43
2	C	1	BGC	O5-C1	-2.09	1.37	1.42

All (10) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	D	2	BGC	O4-C4-C5	4.11	119.51	109.30
2	D	2	BGC	C1-C2-C3	3.93	114.50	109.67
2	C	2	BGC	C1-C2-C3	3.88	114.43	109.67
2	C	2	BGC	O4-C4-C5	2.90	116.49	109.30
2	C	1	BGC	O5-C1-C2	-2.81	105.26	110.28
2	D	1	BGC	O5-C5-C4	2.41	114.08	109.69
2	D	2	BGC	O5-C1-C2	2.34	114.38	110.77
2	D	2	BGC	C3-C4-C5	-2.16	106.39	110.24
2	D	2	BGC	O2-C2-C1	2.10	113.44	109.15
2	C	2	BGC	O2-C2-C1	2.06	113.37	109.15

There are no chirality outliers.


All (4) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
2	C	2	BGC	O5-C5-C6-O6
2	C	2	BGC	C4-C5-C6-O6
2	C	1	BGC	C4-C5-C6-O6
2	D	2	BGC	C4-C5-C6-O6

There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.





## 5.6 Ligand geometry [i](#)

There are no ligands in this entry.

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data ⓘ

### 6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2		OWAB(Å <sup>2</sup> )	Q<0.9
1	A	822/822 (100%)	0.45	41 (4%)	28 28	3, 10, 30, 180	0
1	B	822/822 (100%)	0.36	20 (2%)	59 57	4, 9, 30, 140	1 (0%)
All	All	1644/1644 (100%)	0.40	61 (3%)	41 41	3, 10, 30, 180	1 (0%)

All (61) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	512	ALA	6.0
1	B	612	THR	4.5
1	B	512	ALA	4.3
1	A	612	THR	4.0
1	A	52	MET	3.5
1	A	258	GLY	3.5
1	B	610	GLN	3.4
1	B	613	ASP	3.4
1	B	274	ASP	3.3
1	A	93	ASP	3.3
1	A	547	ARG	3.2
1	A	257	PRO	3.1
1	B	382	GLU	3.0
1	A	611	ASP	3.0
1	A	448	VAL	3.0
1	A	613	ASP	3.0
1	B	1	MET	2.9
1	B	140	SER	2.9
1	A	513	GLY	2.9
1	A	10	ALA	2.9
1	A	274	ASP	2.8
1	A	537	ALA	2.8
1	A	382	GLU	2.7
1	B	802	VAL	2.7

*Continued on next page...*



*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	A	597	PHE	2.5
1	A	608	GLY	2.5
1	B	608	GLY	2.4
1	A	141	TYR	2.4
1	A	610	GLN	2.4
1	A	664	PHE	2.4
1	B	611	ASP	2.4
1	A	1	MET	2.3
1	B	597	PHE	2.3
1	B	286[A]	VAL	2.3
1	A	542	ALA	2.3
1	B	764	ASP	2.3
1	A	138	SER	2.3
1	A	135	THR	2.2
1	A	536	ALA	2.2
1	A	137	THR	2.2
1	A	557[A]	ASP	2.2
1	A	92	LEU	2.2
1	B	607	GLU	2.2
1	B	362	ARG	2.2
1	A	16	THR	2.2
1	B	135	THR	2.2
1	A	455	GLY	2.1
1	A	580[A]	ILE	2.1
1	A	606	GLY	2.1
1	A	300	THR	2.1
1	A	81	TRP	2.1
1	B	316	THR	2.1
1	B	2	ARG	2.1
1	A	67	ALA	2.1
1	A	360	ILE	2.1
1	B	356	PHE	2.1
1	A	234	GLU	2.1
1	A	114	ARG	2.0
1	A	609	PRO	2.0
1	A	281	ASP	2.0
1	A	607	GLU	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains ⓘ


There are no non-standard protein/DNA/RNA residues in this entry.


## 6.3 Carbohydrates ⓘ

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q<0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
2	BGC	C	1	12/12	0.91	0.14	10,19,23,26	0
2	BGC	C	2	11/12	0.92	0.15	5,12,19,25	0
2	BGC	D	2	11/12	0.92	0.15	5,9,15,18	0
2	BGC	D	1	12/12	0.93	0.12	6,13,21,26	0

The following is a graphical depiction of the model fit to experimental electron density for oligosaccharide. Each fit is shown from different orientation to approximate a three-dimensional view.





## 6.4 Ligands [i](#)

There are no ligands in this entry.

## 6.5 Other polymers [i](#)

There are no such residues in this entry.