



# Full wwPDB X-ray Structure Validation Report ⓘ

May 29, 2020 – 10:28 pm BST

PDB ID : 3S5S  
Title : Crystal structure of putative mandelate racemase/muconate lactonizing enzyme (PSI/COM target 200551) from *Sorangium cellulosum*  
Authors : Ramagopal, U.A.; Toro, R.; Almo, S.C.; New York Structural Genomics Research Consortium (NYSGRC)  
Deposited on : 2011-05-23  
Resolution : 2.40 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.11  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.11


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

## *X-RAY DIFFRACTION*

The reported resolution of this entry is 2.40 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	3907 (2.40-2.40)
Clashscore	141614	4398 (2.40-2.40)
Ramachandran outliers	138981	4318 (2.40-2.40)
Sidechain outliers	138945	4319 (2.40-2.40)
RSRZ outliers	127900	3811 (2.40-2.40)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	389	<div> <div>4%</div> <div> <div></div> <div>77%</div> <div>15%</div> <div>• 6%</div> </div> </div>
1	B	389	<div> <div>3%</div> <div> <div></div> <div>79%</div> <div>11%</div> <div>•• 8%</div> </div> </div>

## 2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 5361 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Mandelate racemase/muconate lactonizing enzyme family protein.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	366	Total	C	N	O	S	0	5	0
			2658	1654	486	508	10			
1	B	358	Total	C	N	O	S	0	7	0
			2617	1625	486	496	10			

There are 44 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	368	ALA	-	EXPRESSION TAG	UNP A9GEI3
A	369	GLU	-	EXPRESSION TAG	UNP A9GEI3
A	370	ASN	-	EXPRESSION TAG	UNP A9GEI3
A	371	LEU	-	EXPRESSION TAG	UNP A9GEI3
A	372	TYR	-	EXPRESSION TAG	UNP A9GEI3
A	373	PHE	-	EXPRESSION TAG	UNP A9GEI3
A	374	GLN	-	EXPRESSION TAG	UNP A9GEI3
A	375	SER	-	EXPRESSION TAG	UNP A9GEI3
A	376	HIS	-	EXPRESSION TAG	UNP A9GEI3
A	377	HIS	-	EXPRESSION TAG	UNP A9GEI3
A	378	HIS	-	EXPRESSION TAG	UNP A9GEI3
A	379	HIS	-	EXPRESSION TAG	UNP A9GEI3
A	380	HIS	-	EXPRESSION TAG	UNP A9GEI3
A	381	HIS	-	EXPRESSION TAG	UNP A9GEI3
A	382	TRP	-	EXPRESSION TAG	UNP A9GEI3
A	383	SER	-	EXPRESSION TAG	UNP A9GEI3
A	384	HIS	-	EXPRESSION TAG	UNP A9GEI3
A	385	PRO	-	EXPRESSION TAG	UNP A9GEI3
A	386	GLN	-	EXPRESSION TAG	UNP A9GEI3
A	387	PHE	-	EXPRESSION TAG	UNP A9GEI3
A	388	GLU	-	EXPRESSION TAG	UNP A9GEI3
A	389	LYS	-	EXPRESSION TAG	UNP A9GEI3
B	368	ALA	-	EXPRESSION TAG	UNP A9GEI3
B	369	GLU	-	EXPRESSION TAG	UNP A9GEI3

*Continued on next page...*

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
B	370	ASN	-	EXPRESSION TAG	UNP A9GEI3
B	371	LEU	-	EXPRESSION TAG	UNP A9GEI3
B	372	TYR	-	EXPRESSION TAG	UNP A9GEI3
B	373	PHE	-	EXPRESSION TAG	UNP A9GEI3
B	374	GLN	-	EXPRESSION TAG	UNP A9GEI3
B	375	SER	-	EXPRESSION TAG	UNP A9GEI3
B	376	HIS	-	EXPRESSION TAG	UNP A9GEI3
B	377	HIS	-	EXPRESSION TAG	UNP A9GEI3
B	378	HIS	-	EXPRESSION TAG	UNP A9GEI3
B	379	HIS	-	EXPRESSION TAG	UNP A9GEI3
B	380	HIS	-	EXPRESSION TAG	UNP A9GEI3
B	381	HIS	-	EXPRESSION TAG	UNP A9GEI3
B	382	TRP	-	EXPRESSION TAG	UNP A9GEI3
B	383	SER	-	EXPRESSION TAG	UNP A9GEI3
B	384	HIS	-	EXPRESSION TAG	UNP A9GEI3
B	385	PRO	-	EXPRESSION TAG	UNP A9GEI3
B	386	GLN	-	EXPRESSION TAG	UNP A9GEI3
B	387	PHE	-	EXPRESSION TAG	UNP A9GEI3
B	388	GLU	-	EXPRESSION TAG	UNP A9GEI3
B	389	LYS	-	EXPRESSION TAG	UNP A9GEI3

- Molecule 2 is SULFATE ION (three-letter code: SO4) (formula: O<sub>4</sub>S).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	O	S	0	0
			5	4	1		

Continued on next page...

*Continued from previous page...*

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	O	S	0	0
			5	4	1		
2	B	1	Total	O	S	0	0
			5	4	1		
2	B	1	Total	O	S	0	0
			5	4	1		


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	28	Total	O	0	0
			28	28		
3	B	38	Total	O	0	0
			38	38		


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Mandelate racemase/muconate lactonizing enzyme family protein



- Molecule 1: Mandelate racemase/muconate lactonizing enzyme family protein



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 43 21 2	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	96.36 Å 96.36 Å 152.35 Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	48.18 – 2.40 48.18 – 2.40	Depositor EDS
% Data completeness (in resolution range)	95.3 (48.18-2.40) 95.3 (48.18-2.40)	Depositor EDS
$R_{merge}$	0.16	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	4.02 (at 2.39 Å)	Xtriage
Refinement program	REFMAC	Depositor
R, $R_{free}$	0.193 , 0.263 0.189 , 0.256	Depositor DCC
$R_{free}$ test set	1386 reflections (5.04%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	31.7	Xtriage
Anisotropy	0.159	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.34 , 29.9	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.94	EDS
Total number of atoms	5361	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	32.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.92% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: SO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.85	0/2695	0.86	2/3653 (0.1%)
1	B	0.87	0/2653	0.87	1/3595 (0.0%)
All	All	0.86	0/5348	0.87	3/7248 (0.0%)

There are no bond length outliers.

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	154	ARG	NE-CZ-NH1	5.40	123.00	120.30
1	A	32	ARG	N-CA-C	-5.23	96.87	111.00
1	A	86	ARG	NE-CZ-NH2	-5.09	117.76	120.30

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2658	0	2663	45	0
1	B	2617	0	2636	34	0
2	A	10	0	0	0	0
2	B	10	0	0	0	0
3	A	28	0	0	2	0
3	B	38	0	0	0	0

*Continued on next page...*



*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
All	All	5361	0	5299	76	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 7.

All (76) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:139:THR:HG22	1:A:325:ASP:OD2	1.35	1.26
1:B:170:ARG:HH11	1:B:170:ARG:HG2	1.11	1.10
1:A:19:LEU:HD23	1:A:330:LEU:HD23	1.56	0.87
1:B:147:ARG:HH11	1:B:147:ARG:HB3	1.41	0.85
1:A:144:SER:HB2	1:A:146:GLU:HG2	1.61	0.82
1:B:170:ARG:NH1	1:B:170:ARG:HG2	1.89	0.81
1:A:372:TYR:O	1:A:373:PHE:HB2	1.81	0.80
1:A:149:GLU:HB2	1:A:181:ALA:HB1	1.73	0.69
1:A:139:THR:CG2	1:A:325:ASP:OD2	2.28	0.68
1:B:150:GLU:O	1:B:154:ARG:HG2	1.95	0.67
1:A:19:LEU:HD12	1:A:30:GLN:O	1.96	0.66
1:A:256:VAL:HG21	1:A:283:ILE:HG23	1.77	0.66
1:B:175:ASP:HB2	1:B:176:PRO:HD3	1.82	0.62
1:B:223:GLN:HG2	1:B:248:GLU:OE1	2.00	0.61
1:A:19:LEU:CD2	1:A:330:LEU:HD23	2.30	0.59
1:B:256:VAL:HG22	1:B:286:VAL:HG12	1.86	0.57
1:B:299[A]:MET:HA	1:B:299[A]:MET:CE	2.35	0.57
1:A:11:SER:OG	1:A:370:ASN:OD1	2.15	0.57
1:A:299[A]:MET:HA	1:A:299[A]:MET:CE	2.34	0.57
1:B:55:LEU:HD23	1:B:58:PHE:HB2	1.88	0.56
1:B:168:GLY:HA3	1:B:199:LEU:CD1	2.36	0.56
1:B:90:ARG:O	1:B:94:GLU:HG2	2.05	0.56
1:A:116:THR:OG1	1:A:354:GLY:HA2	2.07	0.54
1:B:170:ARG:HH11	1:B:170:ARG:CG	2.01	0.54
1:B:170:ARG:O	1:B:171:LEU:HB2	2.07	0.54
1:A:164:LYS:HE3	1:A:194:ASP:OD2	2.10	0.52
1:A:339:PHE:C	1:A:339:PHE:CD2	2.82	0.52
1:A:10:VAL:HG22	1:A:40:VAL:HG22	1.92	0.52
1:A:192:ILE:HG12	1:A:220:LEU:HB3	1.92	0.51
1:B:299[A]:MET:HE2	1:B:299[A]:MET:HA	1.92	0.51
1:A:339:PHE:HD2	1:A:339:PHE:O	1.93	0.51
1:B:147:ARG:HH11	1:B:147:ARG:CB	2.19	0.51
1:A:225:VAL:HB	1:A:226:PRO:CD	2.42	0.50

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:168:GLY:HA3	1:B:199:LEU:HD11	1.93	0.50
1:B:144:SER:HB2	1:B:147:ARG:H	1.78	0.49
1:A:230:TRP:CH2	1:A:258:ARG:HG2	2.47	0.49
1:A:339:PHE:CD2	1:A:339:PHE:O	2.65	0.49
1:A:230:TRP:CZ3	1:A:258:ARG:HG2	2.48	0.49
1:A:55:LEU:HD23	1:A:58:PHE:HB2	1.95	0.49
1:A:339:PHE:HD2	1:A:339:PHE:C	2.15	0.49
1:B:199:LEU:HB2	1:B:224:PRO:O	2.13	0.48
1:B:227:ARG:HG3	1:B:249:SER:HA	1.95	0.48
1:A:58:PHE:CD1	1:A:299[A]:MET:HE1	2.49	0.47
1:B:339:PHE:C	1:B:339:PHE:CD2	2.88	0.47
1:A:299[A]:MET:HB3	3:A:398:HOH:O	2.14	0.47
1:A:221:LEU:HD23	1:A:236:VAL:HG11	1.97	0.46
1:A:58:PHE:CG	1:A:299[A]:MET:HE1	2.49	0.46
1:A:30:GLN:HG3	1:A:32:ARG:O	2.15	0.46
1:A:19:LEU:HD23	1:A:330:LEU:CD2	2.37	0.46
1:A:87:ALA:HB2	1:B:130:SER:HA	1.97	0.46
1:A:225:VAL:HB	1:A:226:PRO:HD2	1.98	0.46
1:B:153[A]:ARG:HH11	1:B:153[A]:ARG:HB3	1.80	0.46
1:B:256:VAL:HG22	1:B:286:VAL:CG1	2.44	0.46
1:A:223:GLN:HG3	1:A:223:GLN:O	2.17	0.45
1:B:306:MET:CE	1:B:339:PHE:CE1	3.00	0.45
1:A:257:LEU:HD21	1:B:257:LEU:CD2	2.46	0.45
1:B:17:ILE:HA	1:B:18:PRO:HD3	1.85	0.44
1:A:299[A]:MET:HA	1:A:299[A]:MET:HE2	1.98	0.44
1:A:175:ASP:HB2	1:A:176:PRO:HD3	2.00	0.43
1:A:287:ALA:HB3	1:A:294:LEU:HD21	1.99	0.43
1:A:199:LEU:HB2	1:A:224:PRO:O	2.18	0.42
1:B:51:GLU:HB3	1:B:305:ALA:HB2	2.01	0.42
1:A:221:LEU:HD23	1:A:236:VAL:CG1	2.49	0.42
1:B:287:ALA:HB3	1:B:294:LEU:HD21	2.01	0.42
1:A:221:LEU:CD2	1:A:236:VAL:HG11	2.49	0.42
1:A:247:ASP:OD1	1:A:271:LYS:HE3	2.19	0.41
1:B:177:ALA:O	1:B:180:GLU:HB2	2.21	0.41
1:A:227:ARG:HG3	1:A:249:SER:HA	2.03	0.41
1:A:271:LYS:HA	3:A:418:HOH:O	2.21	0.41
1:B:18:PRO:HB2	1:B:331:ALA:HB3	2.01	0.41
1:B:306:MET:HG2	1:B:327:PRO:HG2	2.03	0.41
1:B:348:LEU:HB3	1:B:351:ILE:HD12	2.03	0.41
1:A:173:ALA:O	1:A:176:PRO:HD2	2.21	0.41
1:A:128:GLY:HA2	1:B:85:TRP:CE2	2.57	0.40

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:18:PRO:HB2	1:A:331:ALA:HB3	2.04	0.40
1:B:112:LEU:HA	1:B:112:LEU:HD23	1.80	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	367/389 (94%)	340 (93%)	27 (7%)	0	100	100
1	B	363/389 (93%)	346 (95%)	15 (4%)	2 (1%)	25	36
All	All	730/778 (94%)	686 (94%)	42 (6%)	2 (0%)	51	55

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	299[A]	MET
1	B	299[B]	MET

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	250/266 (94%)	227 (91%)	23 (9%)	9	13
1	B	246/266 (92%)	225 (92%)	21 (8%)	10	16

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
All	All	496/532 (93%)	452 (91%)	44 (9%)	13	14

All (44) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	2	THR
1	A	21	GLU
1	A	30	GLN
1	A	31	GLU
1	A	59	ASN
1	A	93	ARG
1	A	117	LYS
1	A	139	THR
1	A	144	SER
1	A	146	GLU
1	A	147	ARG
1	A	154	ARG
1	A	180	GLU
1	A	190	SER
1	A	212	ARG
1	A	288[A]	ARG
1	A	288[B]	ARG
1	A	299[A]	MET
1	A	299[B]	MET
1	A	303	VAL
1	A	339	PHE
1	A	371	LEU
1	A	373	PHE
1	B	9	ARG
1	B	20	HIS
1	B	30[A]	GLN
1	B	30[B]	GLN
1	B	93[A]	ARG
1	B	93[B]	ARG
1	B	94	GLU
1	B	141	THR
1	B	142	THR
1	B	147	ARG
1	B	153[A]	ARG
1	B	153[B]	ARG
1	B	154	ARG
1	B	170	ARG

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	B	196	ASN
1	B	256	VAL
1	B	288[A]	ARG
1	B	288[B]	ARG
1	B	299[A]	MET
1	B	299[B]	MET
1	B	306	MET

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	59	ASN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 5.6 Ligand geometry [i](#)

4 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
2	SO4	B	391	-	4,4,4	0.29	0	6,6,6	0.24	0

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
2	SO4	A	391	-	4,4,4	0.38	0	6,6,6	0.50	0
2	SO4	B	390	-	4,4,4	0.31	0	6,6,6	0.30	0
2	SO4	A	390	-	4,4,4	0.70	0	6,6,6	0.75	0

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	366/389 (94%)	-0.18	15 (4%) 37 36	15, 30, 66, 78	0
1	B	358/389 (92%)	-0.38	11 (3%) 49 47	16, 26, 48, 74	0
All	All	724/778 (93%)	-0.28	26 (3%) 42 42	15, 28, 63, 78	0

All (26) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	28	GLY	5.3
1	A	29	ALA	4.8
1	B	26	ALA	4.4
1	A	22	PRO	4.3
1	B	27	GLY	4.3
1	B	171	LEU	3.8
1	B	24	GLY	3.4
1	A	367	GLY	3.3
1	B	170	ARG	3.2
1	B	141	THR	3.1
1	A	28	GLY	3.0
1	B	30[A]	GLN	3.0
1	A	184	ALA	3.0
1	B	142	THR	3.0
1	A	213	ARG	2.9
1	A	27	GLY	2.9
1	A	147	ARG	2.7
1	A	177	ALA	2.7
1	A	21	GLU	2.7
1	A	146	GLU	2.7
1	A	373	PHE	2.6
1	B	25	ILE	2.6
1	A	144	SER	2.3
1	B	29	ALA	2.2

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	A	228[A]	ASP	2.2
1	A	151	ALA	2.2

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
2	SO4	A	390	5/5	0.79	0.21	46,48,53,57	0
2	SO4	B	390	5/5	0.96	0.11	51,55,56,56	0
2	SO4	A	391	5/5	0.96	0.18	58,60,60,62	0
2	SO4	B	391	5/5	0.97	0.22	56,58,60,60	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.