



# Full wwPDB X-ray Structure Validation Report ⓘ

May 26, 2020 – 06:39 pm BST

PDB ID : 3S8C  
Title : Structure of Yeast Ribonucleotide Reductase 1 R293A with AMPPNP and CDP  
Authors : Ahmad, M.F.; Kaushal, P.S.; Wan, Q.; Wijeratna, S.R.; Huang, M.; Dealwis, C.D.  
Deposited on : 2011-05-27  
Resolution : 2.77 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.11  
buster-report : 1.1.7 (2018)  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.11


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

## *X-RAY DIFFRACTION*

The reported resolution of this entry is 2.77 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	4107 (2.80-2.76)
Clashscore	141614	4575 (2.80-2.76)
Ramachandran outliers	138981	4487 (2.80-2.76)
Sidechain outliers	138945	4489 (2.80-2.76)
RSRZ outliers	127900	4027 (2.80-2.76)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	888	

## 2 Entry composition [i](#)

There are 5 unique types of molecules in this entry. The entry contains 5346 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Ribonucleoside-diphosphate reductase large chain 1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	666	5254	3338	893	992	31	22	0	0

There is a discrepancy between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	293	ALA	ARG	ENGINEERED MUTATION	UNP P21524

- Molecule 2 is PHOSPHOAMINOPHOSPHONIC ACID-ADENYLATE ESTER (three-letter code: ANP) (formula:  $C_{10}H_{17}N_6O_{12}P_3$ ).




Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
			Total	C	N	O	P		
2	A	1	31	10	6	12	3	0	0

- Molecule 3 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	1	Total Mg 1 1	0	0

- Molecule 4 is CYTIDINE-5'-DIPHOSPHATE (three-letter code: CDP) (formula:  $C_9H_{15}N_3O_{11}P_2$ ).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
4	A	1	Total	C	N	O	P	0	0
			25	9	3	11	2		


- Molecule 5 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	A	35	Total O 35 35	0	0

### 3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $\text{RSRZ} > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Ribonucleoside-diphosphate reductase large chain 1



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 2	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	107.78Å 116.81Å 64.13Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	40.29 – 2.77 40.08 – 2.77	Depositor EDS
% Data completeness (in resolution range)	98.3 (40.29-2.77) 98.3 (40.08-2.77)	Depositor EDS
$R_{merge}$	0.14	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	3.08 (at 2.77Å)	Xtriage
Refinement program	REFMAC 5.2.0019	Depositor
R, $R_{free}$	0.196 , 0.256 0.200 , 0.257	Depositor DCC
$R_{free}$ test set	2086 reflections (10.03%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	52.1	Xtriage
Anisotropy	0.017	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.32 , 37.1	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.93	EDS
Total number of atoms	5346	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	57.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.52% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: CDP, MG, ANP

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.90	6/5373 (0.1%)	0.88	11/7284 (0.2%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	2

All (6) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	80	ARG	CD-NE	-12.15	1.25	1.46
1	A	620	CYS	CB-SG	-10.46	1.64	1.82
1	A	553	LYS	CB-CG	-9.83	1.26	1.52
1	A	376	LYS	CG-CD	7.94	1.79	1.52
1	A	578	LYS	CG-CD	-6.71	1.29	1.52
1	A	472	GLU	CG-CD	5.68	1.60	1.51

All (11) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	553	LYS	CA-CB-CG	8.41	131.91	113.40
1	A	584	ASP	CB-CG-OD1	7.17	124.76	118.30
1	A	578	LYS	CB-CG-CD	6.75	129.14	111.60
1	A	594	LYS	CG-CD-CE	-6.46	92.52	111.90
1	A	620	CYS	N-CA-CB	-6.36	99.15	110.60
1	A	594	LYS	CD-CE-NZ	-6.30	97.21	111.70
1	A	220	LEU	CA-CB-CG	6.23	129.63	115.30

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	80	ARG	CD-NE-CZ	-6.15	114.99	123.60
1	A	80	ARG	CG-CD-NE	6.15	124.71	111.80
1	A	383	ILE	CG1-CB-CG2	5.21	122.86	111.40
1	A	376	LYS	CB-CG-CD	-5.12	98.29	111.60

There are no chirality outliers.

All (2) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	619	GLU	Peptide
1	A	80	ARG	Sidechain

## 5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	5254	0	5133	96	0
2	A	31	0	13	1	0
3	A	1	0	0	0	0
4	A	25	0	12	1	0
5	A	35	0	0	2	0
All	All	5346	0	5158	98	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 9.

All (98) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:724:LEU:HA	1:A:727:MET:HE3	1.43	0.98
1:A:619:GLU:O	1:A:620:CYS:HB3	1.71	0.89
1:A:665:THR:HG23	1:A:666:GLN:HE21	1.43	0.84
1:A:270:ASN:HB3	1:A:274:PRO:HG2	1.62	0.80
1:A:459:ASP:OD1	1:A:460:GLY:N	2.19	0.76
1:A:530:LEU:O	1:A:534:GLN:HG3	1.88	0.74

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:417:LYS:HE3	1:A:574:MET:HE1	1.69	0.73
1:A:716:LEU:HD11	1:A:727:MET:HE2	1.72	0.72
1:A:724:LEU:HA	1:A:727:MET:CE	2.22	0.69
1:A:505:ILE:HG22	1:A:602:THR:HA	1.75	0.68
1:A:716:LEU:HD11	1:A:727:MET:CE	2.24	0.68
1:A:345:ASN:HD22	1:A:386:GLN:CD	1.98	0.66
1:A:619:GLU:O	1:A:620:CYS:CB	2.36	0.66
1:A:214:GLN:NE2	1:A:216:SER:H	1.94	0.66
1:A:560:PHE:CZ	1:A:596:GLY:HA2	2.30	0.64
1:A:106:ASN:HB3	1:A:109:THR:HG22	1.80	0.63
1:A:265:THR:HB	1:A:267:GLY:H	1.63	0.63
1:A:214:GLN:HE22	1:A:216:SER:H	1.48	0.62
1:A:109:THR:HG23	1:A:111:LYS:H	1.65	0.62
1:A:251:HIS:HB3	1:A:424:SER:HB3	1.82	0.61
1:A:139:ARG:O	1:A:142:GLN:HG2	2.01	0.60
1:A:383:ILE:HG22	5:A:913:HOH:O	2.01	0.60
1:A:522:PRO:HG2	1:A:525:SER:HB3	1.84	0.59
1:A:686:THR:HG23	1:A:688:TRP:HD1	1.68	0.59
1:A:508:GLY:HA3	1:A:606:MET:HE1	1.87	0.57
1:A:126:GLU:OE1	1:A:181:ARG:NH1	2.33	0.57
1:A:120:VAL:HG21	1:A:209:GLY:HA2	1.85	0.57
1:A:428:CYS:HB2	1:A:430:GLU:OE2	2.05	0.56
1:A:655:TRP:HA	1:A:659:MET:HG3	1.87	0.56
1:A:170:LEU:O	1:A:174:VAL:HG23	2.05	0.56
1:A:534:GLN:O	1:A:538:THR:HG23	2.07	0.55
1:A:447:SER:HB3	1:A:606:MET:CE	2.37	0.55
1:A:619:GLU:HB3	1:A:620:CYS:HB2	1.89	0.54
1:A:220:LEU:HD21	1:A:426:ASN:HB3	1.89	0.54
1:A:334:ILE:HD12	1:A:404:VAL:HG13	1.90	0.53
1:A:482:ASN:HD22	1:A:599:ASN:HD21	1.55	0.53
2:A:889:ANP:H8	2:A:889:ANP:O5'	2.09	0.53
1:A:106:ASN:OD1	1:A:109:THR:HG22	2.09	0.53
1:A:518:LEU:HD13	1:A:644:TYR:CE2	2.44	0.52
1:A:627:ASN:ND2	1:A:668:GLY:O	2.43	0.51
1:A:383:ILE:CG2	5:A:913:HOH:O	2.58	0.51
1:A:345:ASN:HD22	1:A:386:GLN:NE2	2.07	0.51
1:A:383:ILE:CD1	1:A:384:LYS:H	2.23	0.51
1:A:482:ASN:ND2	1:A:599:ASN:HD21	2.09	0.51
1:A:213:PRO:HD2	1:A:489:TYR:HB2	1.93	0.51
1:A:84:SER:O	1:A:88:LYS:HG3	2.10	0.51
1:A:456:THR:O	1:A:458:GLU:HG3	2.11	0.50

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:604:ALA:HB2	1:A:708:GLN:HB2	1.92	0.50
1:A:713:ASN:ND2	1:A:742:TYR:HB2	2.27	0.50
1:A:560:PHE:CE2	1:A:596:GLY:HA2	2.47	0.50
1:A:84:SER:HA	1:A:87:HIS:HD2	1.77	0.50
1:A:740:MET:SD	1:A:743:LEU:HB2	2.52	0.49
1:A:692:GLN:O	1:A:696:ILE:HG12	2.13	0.49
1:A:265:THR:HB	1:A:267:GLY:N	2.26	0.49
1:A:106:ASN:HB3	1:A:109:THR:CG2	2.43	0.49
1:A:686:THR:HG23	1:A:688:TRP:CD1	2.47	0.48
1:A:447:SER:HB3	1:A:606:MET:HE3	1.95	0.48
1:A:676:VAL:HB	1:A:681:LYS:HE3	1.96	0.48
1:A:627:ASN:H	1:A:668:GLY:HA3	1.77	0.48
1:A:686:THR:CG2	1:A:688:TRP:HD1	2.25	0.48
1:A:327:ASP:O	1:A:328:LEU:HD23	2.14	0.47
1:A:723:LYS:O	1:A:727:MET:HG3	2.14	0.47
1:A:83:ILE:O	1:A:86:LEU:HB3	2.15	0.47
1:A:217:SER:HB2	1:A:445:LEU:HD13	1.96	0.47
1:A:606:MET:HE3	1:A:608:THR:CG2	2.44	0.47
1:A:526:GLU:OE2	1:A:529:ARG:NH1	2.46	0.46
1:A:544:MET:CE	1:A:570:LEU:HD22	2.45	0.46
1:A:140:ASP:OD2	1:A:167:PRO:HB2	2.14	0.46
1:A:84:SER:HA	1:A:87:HIS:CD2	2.51	0.46
1:A:538:THR:HB	1:A:583:TRP:NE1	2.32	0.45
1:A:334:ILE:HA	1:A:335:PRO:HD2	1.85	0.45
1:A:456:THR:HA	1:A:463:SER:HA	1.99	0.45
1:A:458:GLU:OE1	1:A:462:THR:HB	2.17	0.44
1:A:534:GLN:O	1:A:538:THR:CG2	2.65	0.44
1:A:630:SER:HB2	1:A:639:GLN:NE2	2.32	0.44
1:A:362:CYS:HA	1:A:421:VAL:HG21	1.99	0.43
1:A:550:LEU:HA	1:A:550:LEU:HD23	1.89	0.43
4:A:891:CDP:H6	4:A:891:CDP:H5'2	1.83	0.43
1:A:556:PRO:HA	1:A:597:VAL:O	2.17	0.43
1:A:287:ASP:O	1:A:289:GLY:N	2.51	0.43
1:A:214:GLN:HE21	1:A:488:ASN:HD21	1.67	0.43
1:A:417:LYS:HE3	1:A:574:MET:CE	2.43	0.43
1:A:363:TYR:HB2	1:A:408:ASP:OD1	2.19	0.43
1:A:726:SER:O	1:A:730:TYR:HB2	2.19	0.42
1:A:128:LYS:HB3	1:A:128:LYS:HE2	1.81	0.42
1:A:447:SER:HB3	1:A:606:MET:HE1	2.01	0.42
1:A:716:LEU:HD11	1:A:727:MET:HE1	2.01	0.42
1:A:211:PRO:O	1:A:213:PRO:HD3	2.21	0.41

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:484:VAL:O	1:A:488:ASN:HB2	2.20	0.41
1:A:745:THR:HG23	1:A:746:GLN:H	1.85	0.41
1:A:259:GLY:HA2	1:A:268:THR:OG1	2.20	0.41
1:A:413:LYS:NZ	1:A:735:GLY:O	2.47	0.41
1:A:538:THR:HB	1:A:583:TRP:HE1	1.85	0.41
1:A:730:TYR:O	1:A:734:LYS:HG2	2.21	0.41
1:A:304:TRP:O	1:A:350:LEU:HA	2.21	0.40
1:A:628:MET:O	1:A:629:TYR:C	2.60	0.40
1:A:642:ASN:HA	1:A:643:PRO:HD3	1.96	0.40
1:A:606:MET:CE	1:A:608:THR:HG23	2.51	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles ⓘ

### 5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	662/888 (74%)	622 (94%)	31 (5%)	9 (1%)	11	31

All (9) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	457	SER
1	A	459	ASP
1	A	629	TYR
1	A	717	ARG
1	A	287	ASP
1	A	620	CYS
1	A	639	GLN
1	A	667	ASN
1	A	288	GLN

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	562/760 (74%)	516 (92%)	46 (8%)	11	30

All (46) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	76	THR
1	A	80	ARG
1	A	86	LEU
1	A	131	LEU
1	A	153	ARG
1	A	154	SER
1	A	160	ASN
1	A	176	LEU
1	A	195	LEU
1	A	214	GLN
1	A	216	SER
1	A	218	CYS
1	A	220	LEU
1	A	265	THR
1	A	277	ARG
1	A	301	LEU
1	A	337	LEU
1	A	368	GLU
1	A	383	ILE
1	A	388	LEU
1	A	394	GLU
1	A	456	THR
1	A	457	SER
1	A	462	THR
1	A	473	ILE
1	A	505	ILE
1	A	512	LEU
1	A	514	ASP
1	A	518	LEU
1	A	520	ARG

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	A	530	LEU
1	A	538	THR
1	A	597	VAL
1	A	606	MET
1	A	610	SER
1	A	624	VAL
1	A	626	SER
1	A	628	MET
1	A	639	GLN
1	A	647	ARG
1	A	667	ASN
1	A	673	LEU
1	A	686	THR
1	A	712	LEU
1	A	724	LEU
1	A	746	GLN

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (9) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	168	GLN
1	A	214	GLN
1	A	251	HIS
1	A	345	ASN
1	A	482	ASN
1	A	613	GLN
1	A	666	GLN
1	A	692	GLN
1	A	713	ASN

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates

There are no carbohydrates in this entry.

## 5.6 Ligand geometry

Of 3 ligands modelled in this entry, 1 is monoatomic - leaving 2 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
4	CDP	A	891	-	21,26,26	2.25	5 (23%)	28,40,40	1.77	5 (17%)
2	ANP	A	889	3	29,33,33	1.85	6 (20%)	31,52,52	2.19	6 (19%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	CDP	A	891	-	-	1/14/32/32	0/2/2/2
2	ANP	A	889	3	-	4/14/38/38	0/3/3/3

All (11) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	A	891	CDP	C6-N1	5.98	1.43	1.35
4	A	891	CDP	C4-N3	5.01	1.43	1.35
2	A	889	ANP	PB-N3B	4.76	1.75	1.63
4	A	891	CDP	PB-O1B	4.26	1.64	1.50
2	A	889	ANP	PG-N3B	4.13	1.74	1.63
2	A	889	ANP	PB-O3A	3.72	1.63	1.59
4	A	891	CDP	O4'-C1'	3.35	1.45	1.41
2	A	889	ANP	PG-O1G	3.34	1.51	1.46
2	A	889	ANP	PB-O1B	2.82	1.50	1.46
2	A	889	ANP	C5-C4	2.36	1.47	1.40
4	A	891	CDP	C2-N3	2.11	1.42	1.38

All (11) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	889	ANP	O1G-PG-N3B	-7.19	101.19	111.77
2	A	889	ANP	O1B-PB-N3B	-4.45	105.22	111.77
4	A	891	CDP	C2-N3-C4	4.42	120.82	116.34
2	A	889	ANP	C3'-C2'-C1'	4.22	107.33	100.98
2	A	889	ANP	O2B-PB-O1B	3.83	117.96	109.92
4	A	891	CDP	O3B-PB-O3A	3.34	115.84	104.64
4	A	891	CDP	O4'-C1'-C2'	-3.34	102.05	106.93
4	A	891	CDP	PA-O3A-PB	-3.16	122.00	132.83
2	A	889	ANP	PA-O3A-PB	-3.00	122.05	132.62
2	A	889	ANP	N3-C2-N1	-2.90	124.15	128.68
4	A	891	CDP	N4-C4-N3	2.28	120.09	116.49

There are no chirality outliers.

All (5) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
2	A	889	ANP	PG-N3B-PB-O1B
2	A	889	ANP	PA-O3A-PB-O1B
2	A	889	ANP	PA-O3A-PB-O2B
2	A	889	ANP	PB-O3A-PA-O2A
4	A	891	CDP	PB-O3A-PA-O1A

There are no ring outliers.

2 monomers are involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	A	891	CDP	1	0
2	A	889	ANP	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues

There are no chain breaks in this entry.

## 6 Fit of model and data ⓘ

### 6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	666/888 (75%)	0.09	27 (4%) 37 32	24, 54, 85, 100	7 (1%)

All (27) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	659	MET	5.4
1	A	638	PHE	4.9
1	A	146	PHE	4.5
1	A	165	GLU	3.9
1	A	457	SER	3.5
1	A	629	TYR	3.3
1	A	662	TYR	3.1
1	A	145	TYR	3.1
1	A	718	ALA	2.9
1	A	164	ALA	2.7
1	A	321	GLU	2.6
1	A	74	TYR	2.6
1	A	318	HIS	2.6
1	A	431	ILE	2.5
1	A	604	ALA	2.5
1	A	640	VAL	2.4
1	A	639	GLN	2.4
1	A	727	MET	2.3
1	A	183	ILE	2.2
1	A	461	LYS	2.2
1	A	163	VAL	2.2
1	A	107	ALA	2.2
1	A	630	SER	2.1
1	A	670	ILE	2.1
1	A	746	GLN	2.1
1	A	443	CYS	2.0
1	A	661	GLN	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 6.4 Ligands [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.


Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
3	MG	A	890	1/1	0.89	0.21	67,67,67,67	0
4	CDP	A	891	25/25	0.93	0.23	64,74,77,79	0
2	ANP	A	889	31/31	0.95	0.12	53,57,62,63	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

**Electron density around CDP A 891:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)





## 6.5 Other polymers [i](#)

There are no such residues in this entry.