



# Full wwPDB X-ray Structure Validation Report ⓘ

May 23, 2020 – 04:18 pm BST

PDB ID : 6S81  
Title : Crystal structure of methionine adenosyltransferase from *Pyrococcus furiosus*  
Authors : Degano, M.; Minici, C.; Porcelli, M.  
Deposited on : 2019-07-08  
Resolution : 1.78 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.11  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.11


# 1 Overall quality at a glance ⓘ

The following experimental techniques were used to determine the structure:

## *X-RAY DIFFRACTION*

The reported resolution of this entry is 1.78 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	9185 (1.80-1.76)
Clashscore	141614	10184 (1.80-1.76)
Ramachandran outliers	138981	10051 (1.80-1.76)
Sidechain outliers	138945	10050 (1.80-1.76)
RSRZ outliers	127900	9032 (1.80-1.76)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	401	<div> <div>5%</div> <div> <div></div> <div>93%</div> <div>6%</div> <div></div> </div> <div>• •</div> </div>
1	B	401	<div> <div>4%</div> <div> <div></div> <div>91%</div> <div>6%</div> <div></div> </div> <div>•</div> </div>
1	C	401	<div> <div>4%</div> <div> <div></div> <div>95%</div> <div>5%</div> <div></div> </div> <div></div> </div>
1	D	401	<div> <div>4%</div> <div> <div></div> <div>96%</div> <div></div> <div></div> </div> <div>•</div> </div>

## 2 Entry composition [i](#)

There are 5 unique types of molecules in this entry. The entry contains 13319 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called S-adenosylmethionine synthase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	389	Total	C	N	O	S	0	6	0
			3057	1938	528	583	8			
1	B	389	Total	C	N	O	S	0	9	0
			3067	1946	528	585	8			
1	C	400	Total	C	N	O	S	0	6	0
			3143	1990	542	604	7			
1	D	400	Total	C	N	O	S	0	5	0
			3132	1982	542	601	7			


- Molecule 2 is MANGANESE (II) ION (three-letter code: MN) (formula: Mn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	B	1	Total	Mn	0	0
			1	1		
2	A	1	Total	Mn	0	0
			1	1		
2	D	1	Total	Mn	0	0
			1	1		
2	C	1	Total	Mn	0	0
			1	1		

- Molecule 3 is CHLORIDE ION (three-letter code: CL) (formula: Cl).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	B	1	Total	Cl	0	0
			1	1		
3	A	2	Total	Cl	0	0
			2	2		
3	C	1	Total	Cl	0	0
			1	1		

- Molecule 4 is 2-(N-MORPHOLINO)-ETHANESULFONIC ACID (three-letter code: MES) (formula: C<sub>6</sub>H<sub>13</sub>NO<sub>4</sub>S).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
4	C	1	Total	C	N	O	S	0	0
			12	6	1	4	1		
4	D	1	Total	C	N	O	S	0	0
			12	6	1	4	1		


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	226	Total	O	0	0
			226	226		
5	B	233	Total	O	0	0
			233	233		
5	C	207	Total	O	0	0
			207	207		
5	D	222	Total	O	0	0
			222	222		


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


#### • Molecule 1: S-adenosylmethionine synthase



#### • Molecule 1: S-adenosylmethionine synthase




#### • Molecule 1: S-adenosylmethionine synthase



#### • Molecule 1: S-adenosylmethionine synthase





## 4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	61.11Å 133.90Å 213.10Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	106.55 – 1.78 113.38 – 1.78	Depositor EDS
% Data completeness (in resolution range)	99.1 (106.55-1.78) 99.4 (113.38-1.78)	Depositor EDS
$R_{merge}$	0.26	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.08 (at 1.78Å)	Xtriage
Refinement program	PHENIX	Depositor
R, $R_{free}$	0.209 , 0.231 0.217 , 0.238	Depositor DCC
$R_{free}$ test set	8269 reflections (4.98%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	26.3	Xtriage
Anisotropy	0.643	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.34 , 41.2	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.56$ , $\langle L^2 \rangle = 0.40$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.96	EDS
Total number of atoms	13319	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	35.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 51.13 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 5.8648e-05. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

<sup>1</sup> Intensities estimated from amplitudes.

<sup>2</sup> Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: MN, MES, CL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.31	0/3118	0.51	0/4217
1	B	0.32	0/3140	0.52	0/4249
1	C	0.31	0/3210	0.51	0/4346
1	D	0.31	0/3196	0.51	0/4328
All	All	0.31	0/12664	0.51	0/17140

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3057	0	3145	11	0
1	B	3067	0	3163	16	0
1	C	3143	0	3218	13	0
1	D	3132	0	3207	7	0
2	A	1	0	0	0	0
2	B	1	0	0	0	0
2	C	1	0	0	0	0
2	D	1	0	0	0	0
3	A	2	0	0	0	0

*Continued on next page...*



*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	B	1	0	0	0	0
3	C	1	0	0	0	0
4	C	12	0	13	0	0
4	D	12	0	13	0	0
5	A	226	0	0	6	0
5	B	233	0	0	5	0
5	C	207	0	0	5	1
5	D	222	0	0	1	1
All	All	13319	0	12759	46	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (46) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:D:77:GLY:O	5:D:601:HOH:O	2.00	0.79
1:B:127:ASN:OD1	5:B:601:HOH:O	2.08	0.72
1:B:228:GLU:OE2	5:B:602:HOH:O	2.08	0.72
1:C:171:GLU:OE2	5:C:602:HOH:O	2.09	0.70
1:A:238:GLU:OE1	5:A:601:HOH:O	2.12	0.67
1:A:345:GLN:OE1	5:A:602:HOH:O	2.17	0.60
1:B:395:GLU:OE1	1:B:397:LYS:NZ	2.37	0.57
1:A:308:LYS:HD2	1:A:313:HIS:CD2	2.40	0.56
1:B:121:ARG:NH2	1:B:186:GLU:OE1	2.39	0.55
1:B:134[A]:ARG:NH1	5:B:604:HOH:O	2.30	0.52
1:A:68:VAL:HG11	1:C:134[A]:ARG:HG2	1.90	0.51
1:B:95:GLU:HB3	1:B:137[B]:GLN:HG3	1.93	0.51
1:B:241:GLU:HG3	1:B:254:ILE:HD12	1.94	0.50
1:A:4:ASN:ND2	1:A:354:LEU:O	2.39	0.49
1:C:134[B]:ARG:NH2	5:C:615:HOH:O	2.45	0.49
1:A:241:GLU:HG3	1:A:254:ILE:HD12	1.95	0.48
1:C:198:ASP:HB2	1:C:274:THR:HA	1.96	0.47
1:D:324:LEU:HD11	1:D:392:MET:HE1	1.97	0.47
1:D:358:ILE:HD11	1:D:379:ALA:HB2	1.95	0.47
1:A:233[B]:LYS:HG2	5:A:789:HOH:O	2.15	0.47
1:B:198:ASP:HB2	1:B:274:THR:HA	1.96	0.47
1:B:98:ASP:O	1:B:99:GLN:HB2	2.14	0.47
1:C:388:LYS:HG3	1:C:391:LYS:HE3	1.97	0.46
1:C:152:ASN:ND2	5:C:608:HOH:O	2.35	0.46
1:D:162:PHE:HA	1:D:342:ILE:O	2.16	0.46

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:198:ASP:HB2	1:A:274:THR:HA	1.97	0.45
1:B:137[A]:GLN:HG2	5:B:788:HOH:O	2.17	0.45
1:A:154:ILE:N	5:A:614:HOH:O	2.50	0.45
1:C:162:PHE:HA	1:C:342:ILE:O	2.17	0.44
1:C:140:VAL:HG21	1:C:145:VAL:CG2	2.48	0.44
1:B:117:LYS:NZ	5:B:620:HOH:O	2.51	0.44
1:B:310:PRO:HG2	1:B:394:LEU:HD11	2.00	0.44
1:B:57:ILE:HD12	1:B:311:VAL:HG21	2.00	0.44
1:C:92:ARG:NH2	1:C:138:GLY:HA2	2.33	0.43
1:A:134[A]:ARG:NH1	5:A:607:HOH:O	2.37	0.43
1:D:198:ASP:HB2	1:D:274:THR:HA	2.00	0.43
1:A:28:ILE:HD11	5:A:737:HOH:O	2.19	0.42
1:B:158[B]:ASN:OD1	1:B:159:ASP:N	2.52	0.42
1:C:303:GLU:OE2	5:C:601:HOH:O	2.21	0.42
1:B:48:ARG:NH2	1:B:394:LEU:O	2.53	0.42
1:D:310:PRO:HB3	1:D:393:ILE:HD13	2.01	0.42
1:B:159:ASP:OD1	1:B:160:THR:N	2.52	0.42
1:C:48:ARG:NH1	5:C:609:HOH:O	2.37	0.42
1:C:75:ARG:O	1:C:220:ASP:HB2	2.20	0.41
1:D:60:HIS:O	1:D:309:ASN:HB2	2.21	0.41
1:C:144:SER:O	1:C:148:LYS:HG2	2.20	0.41

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
5:C:738:HOH:O	5:D:601:HOH:O[1_455]	2.14	0.06

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	391/401 (98%)	388 (99%)	3 (1%)	0	100	100
1	B	394/401 (98%)	390 (99%)	4 (1%)	0	100	100
1	C	404/401 (101%)	399 (99%)	5 (1%)	0	100	100
1	D	403/401 (100%)	398 (99%)	5 (1%)	0	100	100
All	All	1592/1604 (99%)	1575 (99%)	17 (1%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	334/339 (98%)	334 (100%)	0	100	100
1	B	337/339 (99%)	337 (100%)	0	100	100
1	C	344/339 (102%)	343 (100%)	1 (0%)	92	90
1	D	343/339 (101%)	342 (100%)	1 (0%)	92	90
All	All	1358/1356 (100%)	1356 (100%)	2 (0%)	93	91

All (2) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	C	328	ASP
1	D	266	ILE

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
1	B	127	ASN

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 5.6 Ligand geometry [i](#)

Of 10 ligands modelled in this entry, 8 are monoatomic - leaving 2 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
4	MES	D	501	-	12,12,12	2.12	1 (8%)	14,16,16	1.50	3 (21%)
4	MES	C	501	-	12,12,12	2.21	1 (8%)	14,16,16	1.44	3 (21%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	MES	D	501	-	-	0/6/14/14	0/1/1/1
4	MES	C	501	-	-	0/6/14/14	0/1/1/1

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	C	501	MES	C8-S	-7.44	1.66	1.77
4	D	501	MES	C8-S	-7.09	1.67	1.77

All (6) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	D	501	MES	O3S-S-C8	3.04	110.68	105.77
4	C	501	MES	O1S-S-C8	2.76	110.24	106.92
4	C	501	MES	C5-N4-C3	2.69	114.89	108.83
4	D	501	MES	O2S-S-C8	2.62	110.07	106.92
4	D	501	MES	C5-N4-C3	2.30	114.00	108.83
4	C	501	MES	O3S-S-C8	2.23	109.37	105.77

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	389/401 (97%)	0.37	19 (4%) 29 28	19, 32, 59, 91	0
1	B	389/401 (97%)	0.40	17 (4%) 34 32	18, 30, 54, 85	0
1	C	400/401 (99%)	0.36	17 (4%) 35 33	21, 31, 58, 88	0
1	D	400/401 (99%)	0.35	15 (3%) 40 39	20, 31, 57, 88	0
All	All	1578/1604 (98%)	0.37	68 (4%) 35 33	18, 31, 58, 91	0

All (68) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	158	ASN	4.6
1	A	156	LEU	4.3
1	A	154	ILE	4.3
1	A	73	TYR	4.2
1	B	263	GLU	4.1
1	A	100	GLU	4.0
1	B	99	GLN	4.0
1	C	263	GLU	3.8
1	C	99	GLN	3.6
1	A	186	GLU	3.6
1	B	264	ARG	3.6
1	A	141	ASP	3.5
1	A	99	GLN	3.5
1	D	143	VAL	3.3
1	B	346	ILE	3.3
1	B	265	GLY	3.3
1	D	151	GLU	3.3
1	A	263	GLU	3.3
1	D	263	GLU	3.2
1	A	142	LEU	3.1
1	D	393	ILE	3.1

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	D	101	LEU	3.0
1	D	75	ARG	2.9
1	D	148	LYS	2.9
1	D	99	GLN	2.9
1	D	146	PHE	2.8
1	B	137[A]	GLN	2.8
1	C	192	TYR	2.8
1	D	2	ALA	2.8
1	A	223	VAL	2.7
1	A	369	LYS	2.7
1	B	75	ARG	2.7
1	C	394	LEU	2.7
1	B	73	TYR	2.6
1	B	100	GLU	2.6
1	C	148	LYS	2.6
1	C	140	VAL	2.5
1	D	264	ARG	2.5
1	C	159	ASP	2.5
1	A	96	LEU	2.5
1	A	386	ILE	2.5
1	A	391	LYS	2.3
1	C	397	LYS	2.3
1	B	139	SER	2.3
1	D	100	GLU	2.3
1	B	369	LYS	2.3
1	C	143	VAL	2.3
1	A	101	LEU	2.3
1	C	381[A]	GLU	2.3
1	C	146	PHE	2.3
1	C	158	ASN	2.3
1	B	158[A]	ASN	2.2
1	C	150	ARG	2.2
1	C	395	GLU	2.2
1	D	346	ILE	2.2
1	A	75	ARG	2.2
1	D	391	LYS	2.2
1	A	2	ALA	2.1
1	C	249	SER	2.1
1	B	155	PRO	2.1
1	B	2	ALA	2.1
1	B	154	ILE	2.1
1	A	68	VAL	2.1

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	C	391	LYS	2.1
1	B	141	ASP	2.1
1	D	144	SER	2.0
1	C	92	ARG	2.0
1	B	230	LEU	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
3	CL	A	503	1/1	0.92	0.09	35,35,35,35	0
4	MES	C	501	12/12	0.92	0.12	33,34,62,74	0
4	MES	D	501	12/12	0.93	0.11	31,34,44,45	0
3	CL	C	503	1/1	0.98	0.24	36,36,36,36	0
3	CL	A	502	1/1	0.99	0.07	33,33,33,33	0
3	CL	B	502	1/1	0.99	0.04	32,32,32,32	0
2	MN	B	501	1/1	1.00	0.04	25,25,25,25	0
2	MN	D	502	1/1	1.00	0.06	24,24,24,24	0
2	MN	C	502	1/1	1.00	0.08	26,26,26,26	0
2	MN	A	501	1/1	1.00	0.07	29,29,29,29	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.